pain after the original insult has gone; they can also spread so that pain may be perceived as coming from a much wider area. Increased sensitisation of the dorsal horn may account for hyperpathia and allodynia. These changes could be responsible for the widespread sensitivity of the back and lower limbs to pressure commonly described by patients with chronic back pain. Descending impulses from the brain can also activate the sensitised dorsal horn so that a potential mechanism exists for integrating psychological distress and pain perception.¹² Patients who are given centrally acting analgesics before operation complain of less severe postoperative pain and wound hyperalgesia.¹³ The possibility of reducing this central sensitisation has implications for treating back pain.

Ultimately the study of any pain must focus on the brain.

We can now identify the anterior cingulate cortex, thalamus, and lenticular nucleus as areas within the brain that are activated by pain.¹⁴ Patients with atypical facial pain who have no demonstrable facial pathology have been shown to have abnormal activation in the anterior cingulate region of their brains, and decreased blood flow in their prefrontal cortex.¹⁵ Their pain is not imagined but is a real phenomenon, with morphological cerebral changes. We need to know whether there are comparable changes in patients with chronic back pain.

MALCOLM I V JAYSON
Professor of rheumatology

Manchester and Salford Back Pain Centre,
Department of Rheumatology,
University of Manchester, Hope Hospital,
Salford M6 8HD

- 1 Walsh K, Cruddas M, Cogan D. Low back pain in eight areas of Britain. *J Epidemiol Community Health* 1992;46:227-30.
- 2 Lawrence JS. Disc disorders. In: *Rheumatism in populations*. London: Heinemann, 1977: 68-97.
- 3 Jensen MC, Brant-Zawadzki MN, Obuchowski N, Modic MT, Malkasian D, Ross JS. Magnetic resonance imaging of the lumbar spine in people without back pain. *N Engl J Med* 1994;331:69-73.
- 4 Mixter WJ, Barr JS. Rupture of the intervertebral disc with involvement of the spinal canal. *N Engl J Med* 1934;211:210-5.
- 5 Colhoun E, McCall IW, Williams L, Cassar-Pullicino VN. Provocation discography as a guide to planning operations on the spine. *J Bone Joint Surg* 1988;70B:267-71.
- 6 Jackson RP, Jacobs RR, Montesano PX. Facet joint injection in low back pain. A prospective statistical study. *Spine* 1988;13: 966-71.
- 7 Porter RW. *Spinal stenosis of the central and root canal in the lumbar spine and back pain*. Edinburgh: Churchill-Livingstone, 1992:313-32.
- 8 Zimmerman M. Components of pain and pain-associated phenomena in the musculo-skeletal system. In: Jayson MIV, Swezey RL, Knoplich J, Hubault A, eds. *Back pain, painful syndromes and muscle spasms*. Carnforth: Parthenon, 1990:29-52.

- 9 Cooper RG, St Clair Forbes W, Jayson MIV. Demonstration of paraspinal muscle wasting in patients with chronic low back pain. *Br J Rheumatol* 1992;31:389-94.
- 10 Hoyland JA, Freemont AJ, Jayson MIV. Intervertebral foramen venous obstruction. A cause of periradicular fibrosis. *Spine* 1989; 14:558-68.
- 11 Lindrom L, Ohlund C, Eek C, Wallin L, Peterson L-E, Nachemson A. Mobility, strength and fitness after a graded activity programme for patients with subacute low back pain. *Spine* 1992;17:641-52.
- 12 Dubner R, Basbaum AI. Spinal dorsal horn plasticity following tissue or nerve injury. In: Wall PD, Melzack R, eds. *Textbook of pain*. Edinburgh: Churchill Livingstone, 1994:225-41.
- 13 Tverskoy M, Oz Y, Isakson A, Finger J, Bradley EL, Kissin I. Preemptive effect of fentanyl and ketamine on post-operative pain and wound hyperalgesia. *Anaesth Analg* 1994;78: 205-9.
- 14 Jones AKP, Brown WD, Friston KJ, Qi LY, Frackowiak RSJ. Cortical and subcortical localisation of response to pain in man using positron emission tomography. *Proc Roy Soc Lond (Biol)* 1991;244:39-44.
- 15 Derbyshire SWG, Jones AKP, Devani P, Friston KJ, Feinmann C, Harris M, et al. Cerebral responses to pain in atypical facial pain measured by positron emission tomography. *J Neurol Neurosurg Psychiatry* (in press).

Preventing blindness worldwide

Deserves a higher priority than some countries are prepared to give it

About 40 million people are blind¹ (defined as a corrected visual acuity of less than 3/60 in the better eye²). Cataract accounts for half the cases, and trachoma, glaucoma, diabetic retinopathy, onchocerciasis, blinding malnutrition (xerophthalmia), leprosy, and trauma account for most of the rest.³ With increasing longevity and population growth, age related visual loss from cataract and glaucoma is likely to increase because eye care services are overwhelmed in all but the most economically developed countries.

Geographically, patterns of blindness vary. In the industrialised countries of North America, Europe, and Australasia blindness is mainly due to disorders of the posterior segment of the eye (for example, macular degeneration and diabetic retinopathy). In Africa, Asia, and parts of South America disorders of the anterior segment of the eye (cataract, corneal scarring from various causes, and glaucoma) predominate.⁴ The prevalence of blindness varies from 0.2% in industrialised nations to 1% or more in poor areas of Africa and Asia. Even within a single country the prevalence of blindness can vary widely between different communities, depending on the availability of primary health care and specialised eye care services.³

Although four out of five cases of blindness occur in developing countries, the resources to combat blindness are concentrated in industrialised and urban areas of the world. Africa has about one ophthalmologist per million population; although 80% of the population live in rural areas, eye care services are almost entirely situated in the capital cities and large provincial towns.⁵

How can this burden of avoidable blindness be reduced, and how can those in need obtain access to services to prevent or cure blindness? The recent general assembly of the International Agency for the Prevention of Blindness addressed these questions in Berlin. The reasons why people do not have easy access to eye care include the paucity of trained personnel, distance to travel to obtain eye care (particularly in rural areas), the cost of treatment, lack of awareness that sight can be restored, and fear of hospitals, doctors, and surgery.

Because poor and rural areas have too few ophthalmologists to tackle the problem it makes sense to train other levels of health care workers to carry out specific tasks. Primary eye care can be integrated into primary health care. Blinding malnutrition from vitamin A deficiency can follow measles or a diarrhoeal disease, often in association with a poor diet. Through promotion of the expanded programme of immunisation, measles can be reduced and even eliminated. Through education, mothers can be encouraged to provide an improved diet for their children. Distribution of capsules of vitamin A, as a short term measure, not only prevents xerophthalmia but reduces mortality and morbidity.⁶ Cleaning the eyelids of all newborn babies and instilling 1% tetracycline ointment in their eyes can prevent ophthalmia neonatorum.⁷ Blinding trachoma could be reduced by face washing and improvements in sanitation.⁸ Distributing ivermectin tablets each year to communities with endemic onchocerciasis by the primary health care team could reduce ocular morbidity from onchocerciasis.⁹

In Africa, many patients blinded by cataract will never have access to an ophthalmologist. Eye nurses and ophthalmic assistants who have been trained to perform intracapsular cataract extraction safely are sent to work in district hospitals under the supervision of a visiting ophthalmologist; this has proved successful in East Africa. In India, where one and a half million cataract operations are performed each year, the burden of 5 million people blind from cataract could be reduced if the 7500 trained ophthalmologists could all be diverted to operating on cataracts.

Equipment and supplies account for up to 80% of the cost of surgery. Such costs can be reduced by appropriate technologies such as the local preparation of eye drops,¹⁰ pooled purchasing of standard equipment, and low cost, local production of eye sutures, aphakic spectacles, and intraocular lenses. The World Bank recognises cataract surgery and the distribution of vitamin A capsules as being among the most cost effective interventions in medicine today.¹¹

The cost of cataract surgery and the distance that patients must travel for surgery are not the only barriers to access: fear and lack of awareness may also be important. A survey in east Baltimore, in the United States, found that a fifth of patients with severe visual impairment had unoperated cataract.¹² Similar rates have been found in London.¹³ In Africa and Asia a fatalistic belief that poor sight is an inevitable consequence of old age is common. In India only between a third and a fifth of people blind from cataract accept the offer of free surgery.¹⁴ In Africa fewer than 5% of those who need cataract surgery receive it.¹⁵ There are cultural and communication barriers that need to be overcome. The patient whose vision has been successfully restored by operation is the best advertisement and advocate for promoting eye surgical services.¹⁴

The ophthalmological community believes that the case for the prevention of blindness is justified on economic

and, more importantly, humanitarian grounds. At present, this is poorly appreciated by many ministries of health in the developing world, where preventing blindness has a low priority, competing with major health issues such as malaria, tuberculosis, and AIDS. For most people, however, blindness is one of the most feared disabilities of life. Success in preventing blindness will depend on well researched and reasoned arguments presented to politicians and policy makers, reinforced by a groundswell of opinion from their constituents that everyone has the right to sight, even those at "the end of the track."

ANDREW R POTTER
Ophthalmologist

Centre Hospitalier Départemental,
BP 49,
Abomey,
Republic of Benin,
West Africa

- 1 Foster A, Johnson GJ. Blindness in the developing world. *Br J Ophthalmol* 1993;77:398-9.
- 2 World Health Organisation. *Guidelines for programmes for prevention of blindness*. Geneva: WHO, 1979:16.
- 3 Foster A. Worldwide blindness, increasing but avoidable. *Seminars in Ophthalmology* 1993;8:166-70.
- 4 Foster A. Patterns of blindness. In: Tasman W, Jaeger E, eds. *Duane's clinical ophthalmology* Vol 5. Philadelphia: Lippincott, 1991:1-7.
- 5 Johnson GJ, Foster A. Training in community ophthalmology. *Ophthalmology International* 1990;14:221-6.
- 6 West KP, Pokhrel RP, Katz J, Le Clerq S, Khattry SK, Tielsch JM, et al. Efficacy of vitamin A in reducing pre-school child mortality in Nepal. *Lancet* 1991;338:67-70.
- 7 World Health Organisation. *Conjunctivitis of the newborn*. Geneva: WHO, 1986.
- 8 Taylor HP, West SK, Mmbaga BBO, Katala SJ, Turner V, Lynch M, et al. Hygiene factors and increased risk of trachoma in central Tanzania. *Arch Ophthalmol* 1989;107:1821-5.
- 9 Abiose A, Jones BR, Cousens SN, Murdoch I, Cassels Brown A, Babaloia OE, et al. Reduction in incidence of optic nerve disease with annual ivermectin to control onchocerciasis. *Lancet* 1993;341:130-4.
- 10 World Health Organisation. *The local small scale preparation of eye drops*. Geneva: WHO, 1990. (WHO/PBL/90.20.)
- 11 World Bank. *Investing in health. World development report 1993*. New York: Oxford University Press, 1994.
- 12 Tielsch JM, Sommer A, Witt K, Katz J, Royall RM. Blindness and visual impairment in an American urban population. *Arch Ophthalmol* 1990;108:286-90.
- 13 Wormald RPL, Wright LA, Courtney P, Beaumont B, Haines AP. Visual problems in the elderly population and implications for services. *BMJ* 1992;304:1226-9.
- 14 Venkataswamy PG, Billant G. Social and economic barriers to cataract surgery in rural south India: a preliminary report. *Visual Impairment and Blindness* 1981;405-8.
- 15 Foster A. Who will operate on Africa's 3 million curably blind people? *Lancet* 1991;337:1267-9.

Acronymophilia

The exponential growth of the use of acronyms should be resisted

Eponyms such as Barlow, Carvallo, Master, and Wolff-Parkinson-White were until recently part of the sport of ward rounds. Times have changed. Now acronyms are in fashion. GUSTO, GISSI, ISIS, MRFIT, and TIMI are heard in hospital corridors and convention halls everywhere.

In cardiology acronyms for research trials have grown almost exponentially. In 1992 I compiled a list of acronyms of 244 major cardiological trials.¹ An updated version two years later saw the total nearly quadrupled to over 900.² As acronyms continue to be born almost every day they need to be made easily understandable by, and readily accessible to, clinicians, investigators, and editors of medical journals while their currency is at a peak.

Acronyms are useful because they simplify, facilitate, and accelerate communication. They have become the shorthand of medicine. Physicians, especially cardiologists, like to use or invent these medical timesavers—and are good at the task.

Acronyms of cardiological studies include many anatomical terms such as ARMS, ATLAS, BRAINS, CARDIA, CARDIAC, CAVA, EARS, FACET, HEART,

INTIMA, MAST, PROSTATE, RADIUS, and RALES.* Other acronyms show that their inventors must have food on their mind—APRICOT, BIG-MAC, DISH, GUSTO, MOCHA, SALAD, SALT, and TOAST. Still others represent places—MIAMI, NEVADA, PARIS, SIAM, THAMES, and TIBET. Some have a romantic touch—CASANOVA, M-HEART, and KISS—or are common women's names—ELSA, ERICA, EVA, GRACE, MONICA, NORA, PAMELA, PHYLLIS, and RITA. Finally, acronyms beget acronyms; there are beginning to be acronyms within acronyms—AIMS, AMPI, APSIM, ARMS, DPPS, LART, NHEFS, TAPS, and THIS.

Yet with this proliferation the rules should remain the same. When mentioned or used for the first time in a publication all acronyms must be fully explained to avoid confusion,³ avert misinterpretation,⁴ eliminate guesswork,⁵ and prevent aggravation.⁶ Specialists often take for granted that certain trade terms are so evident or self explanatory that they do not bother to define them. Not all readers of

*Explanation of all the acronyms cited in this editorial can be found in table 1 of reference 2.