with uptake rates for medical prescriptions in general.⁴ Those attending were not the "worried well" but patients referred by their general practitioners for help with a range of medical conditions such as obesity, hypertension, depression, and raised cholesterol concentrations.

The full evaluation report is available from the leisure services department of Stockport Metropolitan Borough Council (tel 061 474 4415). There is no doubt about the validity of the findings of the Stockport scheme, and we urge general practices not to look too long before leaping into a health promotion activity that could contribute towards improved cardiovascular and mental health.

FELICITY GREEN
Deputy director

Stockport Centre for Health Promotion,
Stockport,
Cheshire SK2 7AE

JANET LORD
Senior research officer

East Lancashire Public Health Resource and Intelligence Centre,
East Lancashire Health Authority,
Nelson,
Lancashire BB9 5SZ

1 Iliffe S, See Tai S, Gould M, Thorogood M, Hillsdon M. Prescribing exercise in general practice. *BMJ* 1994;309:494-5. (20-27 August.)

2 Goldberg DP. *Manual of the general health questionnaire*. Windsor: NFER Publishing, 1979.

3 Smith R. Where is the wisdom...? *BMJ* 1991;303:798-9.

4 Beardon P, McGilchrist M, McKendrick A, McDevitt D, MacDonald T. Primary non-compliance with prescribed medication in primary care. *BMJ* 1993;307:846-8.

Persistent post-traumatic stress disorder

Role of alcohol ignored

EDITOR,—The grand round presented by Brigitta C Bende and Robin M Philpott contains two major logical errors, and I am surprised that these were not mentioned in the discussion.¹ The patient was a heavy drinker with all the features of an organic mental disorder, including cognitive impairment. While he may have suffered a traumatic neurosis in the aftermath of his wartime experience, the continuing features that were the basis for the diagnosis of post-traumatic stress disorder should have been ascribed to alcohol unless they were still present after a lengthy period of abstinence. In fact, all symptoms cleared when he abstained. Frightening ideas and nightmares are common in heavy drinkers, and their content is determined by the previous experience of the sufferer.

Secondly, how can improvement be attributed to the various psychological treatments that the patient received when he simultaneously received a far more powerful treatment for his brain disorder—namely, abstinence from alcohol? Had his doctors waited to see the effects of this they would probably have found that no other treatment was necessary.

Forty per cent of all patients diagnosed as having post-traumatic stress disorder drink heavily,² and their symptoms subside when they abstain.

SAMUEL I COHEN
Emeritus professor of psychiatry

London NW11 7LT

1 Bende BC, Philpott RM, Saltissi S, Edwards RHT, Williams G, Wyatt GB, et al. Persistent post-traumatic stress disorder. *BMJ* 1994;309:526-8. (20-27 August.)

2 Wilson GT. Alcohol and anxiety. *Behav Res Ther* 1988;26:369-81.

May not need treatment in war veterans

EDITOR,—Brigitta C Bende and Robin M Philpott construe the current problems—depression and alcohol misuse—of a veteran of the second world war as being due to a 50 year old psychiatric

condition, persistent post-traumatic stress disorder.¹ When such veterans present now with disturbing wartime memories does it follow that they had post-traumatic stress disorder all along?

Firstly, an episode of psychiatric or physical disease arising for the first time may evoke emotive associations with the threat, horror, and loss of war. For example, recently a survivor of a Japanese concentration camp began to talk obsessively of his experiences, which his wife had never heard about before, in the aftermath of a near fatal cerebrovascular accident.

Secondly, the aging process may put pressure on physiological and psychological resources and thus on the adjustments and philosophies that have long characterised a person. At a time of diminishing roles and duties the past may be more vivid than the present, and many people look back over their lives in preparation for death. Extreme experiences, and the resonant questions about life and death that these threw up, may again loom large.

Thirdly, and fundamentally, a person processes trauma as a function of the social meaning assigned to it.² But such social constructions do not necessarily stand still: since 1945 Western culture has seen a dramatic rise in the power of psychological explanations for the world. Not just war but also peacetime disasters, criminal violence, childhood sexual abuse, and even bullying are now viewed as having potentially lifelong psychological consequences for victims and are much discussed in the media. This social climate influences and, indeed, may redefine what veterans think now about what happened to them during the war.

The features of post-traumatic stress disorder are not exclusive to survivors of war or other disasters; they are common in people stressed in more everyday ways, including those who have been bereaved. Our work with former combatants indicates that most of those in whom the disorder can be diagnosed do not need psychological help.³ Nor is the disorder necessarily the key to the problems of those who present in clinics. Moreover, whether the disorder is a clinical entity robust enough to have a natural cause traceable over decades is debatable. Few veterans say that war service left no mark on them, but most live unremarkably.

The authors of the paper state that post-traumatic stress disorder is often missed because doctors are not aware of it; in fact, it is now highly fashionable. As with other longstanding psychiatric diagnoses, we must take account of concerns about social labelling and the sick role. Human responses to war are multiply determined, and we must remember the inherent limitations of a narrowly medicalised framework in this complex arena.

DEREK SUMMERFIELD
Principal psychiatrist

Medical Foundation for the Care of
Victims of Torture,
London NW5 3EJ

FRANCESCA HUME
Senior psychologist

Department of Forensic Psychiatry,
St George's Hospital,
London SW17 0RG

1 Bende B, Philpott R. Persistent post-traumatic stress disorder. *BMJ* 1994;309:526-8. (20-27 August.)

2 Summerfield D, Hume F. War and post-traumatic stress disorder: the question of social context. *J Nerv Mental Dis* 1993;181:522.

3 Hume F, Summerfield D. After the war in Nicaragua: a psychosocial study of war wounded ex-combatants. *Medicine and War* 1994;10:4-25.

Former prisoners of war show symptoms

EDITOR,—In the discussion about an 80 year old man presenting with a primary diagnosis of post-traumatic stress disorder with comorbid alcohol misuse and depression the comment is made that similar symptoms have emerged at retirement in former prisoners of war seen at Liverpool School of

Tropical Medicine.¹ Over recent years people who were prisoners of war in the Far East have been screened for tropical diseases at Royal Air Force Hospital Wroughton. This has given us the opportunity to examine these veterans for psychological symptoms and to give at least an approximate answer to the implied question in the paper about their prevalence.

Goldstein *et al*, who studied 32 Americans who had been prisoners of war in the Far East, found a prevalence of psychological symptoms of 50%.² In our sample of 30 British people who had been prisoners of war in the Far East (mean age 75.4 years) we found that 30% fulfilled the criteria in the *Diagnostic and Statistical Manual of Mental Disorders, Third Edition, Revised* for a diagnosis of persistent post-traumatic stress disorder.³ Moreover, 90% had at least one post-traumatic psychological symptom. Our fuller results await publication. We have also noted appreciable comorbidity, and, as in other studies,⁴ major depression has predominated. The emergence of symptoms in later life, including at retirement, has been discussed,⁵ but in our sample (as in the case discussed¹) there was ample evidence that for many people the intervening years had been restricted and blighted by wartime trauma. The possibility of post-traumatic stress disorder in the present cohort of elderly people should not be overlooked.

J C HUGHES
Registrar
L A NEAL
Consultant

Department of Military Psychiatry,
Princess Alexandra's Hospital,
Royal Air Force Wroughton,
Swindon,
Wiltshire SN4 0QJ

1 Bende BC, Philpott RM. Persistent post-traumatic stress disorder. *BMJ* 1994;309:526-8. (20-27 August.)

2 Goldstein G, van Kammen W, Shelly C, Miller DJ, van Kammen DP. Survivors of imprisonment in the Pacific theatre during world war II. *Am J Psychiatry* 1987;144:1210-3.

3 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders, third edition, revised*. Washington, DC: APA, 1987.

4 McFarlane AC, Papay P. Multiple diagnoses in posttraumatic stress disorder in the victims of a natural disaster. *J Nerv Ment Dis* 1992;180:498-504.

5 Busuttill W, Aquilina C, Busuttill A. Early life trauma, infertility and late life paranoia: a relationship with post traumatic stress disorder? *Int J Geriatr Psychiatry* 1993;8:693-5.

One and two sided tests of significance

Statistical hypothesis should be brought into line with clinical hypothesis

EDITOR,—I believe that J Martin Bland and Douglas G Altman's strong preference for two sided significance tests results in one sided advice.¹ The wide acceptance of two sided statistical alternative hypotheses in the case of clearly one sided clinical or biological hypotheses has always amazed me. To my mind the statistical alternative hypothesis should be brought into line with the clinical hypothesis more often, even if this means a shift towards one sided tests. If on the basis of scientific considerations the clinical hypothesis is one sided then so should be the alternative statistical hypothesis, which then more precisely reflects the expectation of the clinical investigator.

Although statistical decisions depend on an outcome variable having or not having some critical value, in the case of an extreme result in an unexpected direction one faces the same practical problem regardless of the choice of a one sided or a two sided test and even regardless of the precise P value. The problem is to decide whether a new treatment is harmful or simply ineffective (if it is ineffective the unexpected outcome will be attributed to chance).

The authors' discussion of the outcome of the

study of Lund *et al* could also serve to stimulate the use of one sided tests in a consistent way instead of the use of two sided tests.

I am also unhappy with the authors' argument that a one sided test is appropriate when a large difference in one direction would lead to the same action as no difference at all, because again the alternative statistical hypothesis is determined not by expectation but primarily by the action to be taken afterwards. Depending on the (lack of) expectation of the direction of the outcome a two sided test may be required here.

As Bland and Altman say, in medicine things do not always work out as expected, but we may not need two sided tests as a reminder of this. In my view, allowing expectation to determine the formulation of the alternative hypothesis makes the process of scientific inference more consistent and transparent.

R WOLTERBEEK
Epidemiologist

3584 HL Utrecht,
Netherlands

1 Bland JM, Altman DH. One and two sided tests of significance. *BMJ* 1994;309:248. (23 July.)

One sided tests should be used more often

EDITOR,—It seems presumptuous for a non-statistician to question the almost ubiquitous use of two sided tests of significance in medical research. I am emboldened to do so by the fact that J Martin Bland and Douglas G Altman challenge conventional wisdom by giving at least some lukewarm support for the occasional use of a one sided test.¹ I suggest that one sided tests should be used far more often.

Randomised trials frequently compare two alternative forms of care, one of which is clearly more expensive or more invasive than the other. Use of the more expensive or more invasive treatment would be justified only if it was more effective than the other. In this case we are interested only in whether the experimental treatment is better than the less expensive or more invasive form. We are indifferent as to whether it is equal or worse.

If we choose to accept a 95% chance of avoiding a type I error (concluding that the treatment is more effective when in truth it is not) then a one tailed test is appropriate. By using a two tailed test we are demanding a 96.5% chance of avoiding a type I error and increase our risk of having a type II error (concluding that the treatment is not more effective when in truth it is). This fact may not be immediately apparent to casual readers.

I agree that the direction of the test and the reason for choosing to use a one sided test must be specified in advance. But I think that the one sided test could, and should, be used far more often than is presently the case.

MURRAY W ENKIN
Professor emeritus

Department of Obstetrics and Gynaecology,
Faculty of Health Sciences, McMaster University,
Hamilton,
Ontario, Canada L8N 3Z5

1 Bland JM, Altman DH. One and two sided tests of significance. *BMJ* 1994;309:248. (23 July.)

Authors' reply

EDITOR,—We cannot agree with R Wolterbeek. Significance testing should be approached from the null hypothesis, which is what is actually tested. In a one sided test the null hypothesis is that there is either no difference or a difference in a specified direction. We think that this rarely makes sense in clinical research. The research hypothesis may lead us to expect a difference in a particular direction, but it should not lead us to say that a

difference in the other direction is the same as no difference.

Murray W Enkin's approach is different. Rather than using the significance test as an indicator of the strength of the evidence to aid a scientific decision, he proposes to use it as a decision rule for an operational decision. This view is often taken in industrial applications of statistical methods, where answers are sought to questions such as whether increasing the temperature of a process increases the output. If there is no increase in output the temperature will be returned to the original level, whether there is a decrease in output or no change. If there is a decrease in output then manufacturers are not concerned with why, only that it should not happen again. If that was the situation we would agree with Enkin. We do not think that medical research proceeds in the same way. Typically, more than one trial will be carried out, and the effect of a single trial is not to change everyone's practice. Furthermore, if a treatment has harmful consequences we should ask why this is, not ignore it. Our statistical method must be able to detect such consequences when they occur.

In any case, we would prefer to present results in the form of confidence intervals, so avoiding the question. We shall pursue this further in future Statistics notes.

J MARTIN BLAND
Reader in medical statistics

Department of Public Health Sciences,
St George's Hospital Medical School,
London

DOUGLAS G ALTMAN
Head

Medical Statistics Laboratory,
Imperial Cancer Research Fund,
London WC2A 3PX

Microscopic haematuria

Role of renal biopsy is undervalued

EDITOR,—Fritz H Schröder emphasises the central importance of cystoscopy in the investigation of microscopic haematuria, regarding renal biopsy as required in only a minority of patients.¹ We have commented previously on the disadvantage of this approach, which is likely falsely to reassure some patients with glomerular disease, which may be progressive.²

Schröder relies on urinary red cell morphology to distinguish glomerular from non-glomerular haematuria and thus to direct the use of renal biopsy. Although both phase contrast microscopy and the alternative Coulter counter analysis of urinary red cells have been advocated enthusiastically, some authors have reported finding them unhelpful, particularly when the number of red cells is low³; the two techniques have not entered widely into British nephrological practice.⁴

We recently reported our experience in 165 patients with microscopic haematuria (aged 18-70) who underwent both renal biopsy and cystoscopy.⁵ We found evidence of glomerular disease in 73 patients, the commonest abnormalities being IgA nephropathy, other forms of mesangial proliferative glomerulonephritis, and thin membrane nephropathy. Age influenced this finding; the biopsy specimen was abnormal in nine of the 13 patients aged under 20 and even in six of the 13 patients in their seventh decade. By contrast an abnormality was found on cystoscopy in only five of the 165 patients (cystitis, three; urethral stricture, one; and radiolucent bladder stone, one). Furthermore, two of the patients with cystitis also had IgA nephropathy.

We accept the central importance of cystoscopy in excluding urothelial malignancy in older patients with microscopic haematuria. The incidence of such malignancy in people aged under 45, however, is extremely low, and a cause for haematuria will be found on renal biopsy in more than 40%. We

Algorithm for investigation of microscopic haematuria

therefore propose a modification of Schröder's algorithm (figure), in which younger patients undergo biopsy before cystoscopy is considered; this practice is already well accepted in children. We also recommend that older patients in whom cystoscopy gives normal results should be offered renal biopsy, although we do not recommend this for patients aged over 70 since glomerular disease found in this age group is unlikely to progress to a major clinical problem in the patient's lifetime.

This might be regarded as an invasive and cost ineffective approach. Schröder, however, recommends repeated imaging, cytological examination, and cystoscopy in patients in whom no cause has been found when his algorithm, which submits only a small minority of patients to renal biopsy, is used.

PETER TOPHAM
Registrar in nephrology

STEVEN HARPER
Wellcome research training fellow

JOHN FEEHALLY
Consultant nephrologist

Department of Nephrology,
Leicester General Hospital,
Leicester LE5 4PW

1 Schröder FH. Microscopic haematuria. *BMJ* 1994;309:70-2. (9 July.)

2 Feehally J, Walls J. Asymptomatic microscopic haematuria. *BMJ* 1986;292:1273.

3 Gibbs DD, Lynn KL. Red cell volume distribution curves in the diagnosis of glomerular and non-glomerular haematuria. *Clin Nephrol* 1990;33:143-7.

4 Feehally J, O'Donoghue DJ, Ballardie FW. Current nephrological practice in the investigation of microscopic haematuria. *J R Coll Physicians Lond* 1989;23:228-31.

5 Topham PS, Harper SJ, Furness PN, Harris KPG, Walls J, Feehally J. Glomerular disease as a cause of isolated microscopic haematuria. *Q J Med* 1994;87:329-36.

Author's reply

EDITOR,—I share Peter Topham and colleagues' views on differentiating between glomerular and non-glomerular haematuria by phase contrast microscopy or Coulter counter analysis. The points that the authors make are what led me to place this technology low in my algorithm. I am reluctant to give renal biopsy a more prominent