Guideline for colposcopic management of mildly abnormal results of cervical cytology (CIN=cervical intraepithelial neoplasia)

screening and the importance of not defaulting is vital. Most cases of cervical cancer occur in those who are unscreened or screened infrequently.⁴

If immediate referral for mild dyskaryosis is adopted, colposcopy must seek to detect and treat any significant abnormality (cervical intraepithelial neoplasia grade II or worse), avoid overtreating very mild lesions, and where possible expedite return to cytological surveillance. The correct management of cervical intraepithelial neoplasia grade I remains a grey area.

In 1991, 46% of referrals to the Oxford colposcopy unit were for persistent mild abnormalities (borderline nuclear change, human papillomavirus change, mild dyskaryosis). This prompted us to draw up guidelines (see figure) for the management of such cases and we have now reviewed the first 100 patients so managed. Our guidelines recommend biopsy rather than "see and treat" unless colposcopy suggests at least cervical intraepithelial neoplasia grade II.

Forty seven patients (mean age 33 years) were referred for mild dyskaryosis, 53 (mean age 39 years) for borderline nuclear or human papillomavirus changes. At first visit a colposcopic diagnosis of cervical intraepithelial neoplasia II or III was made in eight patients, seven of whom had been referred with mild dyskaryosis. After initial assessment 44 patients were discharged untreated for follow up cytology at one year, 22 were kept under review, and 34 were treated (23 at the first visit) by large loop excision of the transformation zone or cold coagulation.

After one year 56 of the 100 patients had been discharged untreated (15 of these were referred with mild dyskaryosis), 40 had been treated (29 referred with mild dyskaryosis, $P < 0.001$), and four remained under review (three referred with mild dyskaryosis). Histological examination for treated patients showed 15 cases of cervical intraepithelial neoplasia II or III, including 13 with mild dyskaryosis on referral ($P < 0.01$). Eleven patients may have been overtreated in that histology showed less than cervical intraepithelial neoplasia I.

These guidelines are helpful in a unit staffed by several doctors. They promote consistency of

management, facilitate audit, and provide sufficient flexibility to take account of patients' particular needs.

HONOR MERRIMAN
Clinical assistant
MARK CHARNOCK
Consultant gynaecologist
WINIFRED GRAY
Consultant pathologist

Oxford Radcliffe NHS Trust,
John Radcliffe Hospital,
Oxford OX3 9DU

Royal Infirmary of Edinburgh NHS Trust,
City Hospital,
Edinburgh EH10 5SB

NICHOLAS HALLAM
Consultant virologist

- 1 Flannely G, Anderson D, Kitchener HC, Mann EMF, Campbell M, Fisher P, et al. Management of woman with mild and moderate cervical dyskaryosis. *BMJ* 1994;308:1399-403. (28 May.)
- 2 Soutter WP, Fletcher A. Invasive cancer of the cervix in woman with mild dyskaryosis followed up cytologically. *BMJ* 1994;308:1421-3. (28 May.)
- 3 Hammond R. Management of woman with smears showing mild dyskaryosis. *BMJ* 1994;308:1383-4. (28 May.)
- 4 Macgregor JE, Campbell MK, Mann EMF, Swanson KY. Screening for cervical intraepithelial neoplasia in north east Scotland shows fall in incidence and mortality from invasive cancer with concomitant rise in preinvasive disease. *BMJ* 1994;308:1407-11. (28 May.)

Intrapartum care and cerebral palsy

EDITOR,—"This week in *BMJ*" for the paper by Geraldine Gaffney and colleagues referred to a retrospective investigation of 339 children aged 5 years, suffering from cerebral palsy.¹ The conclusion that only 6.8% of cases occurred in non-malformed term singleton babies following suboptimal obstetrical care, extrapolated to all patients with cerebral palsy in the region, is not supported by the data.

Nine subjects were born outside the area, 42 had abnormality or unidentified antepartum causes of cerebral palsy, 27 had postnatal causes, and the obstetrical antecedents of 125 were not investigated. There were therefore 205 children born in

the area who were investigated, of whom 138 (67%) may have suffered from intrapartum asphyxia. The obstetrical care of 128 was described in table V, which reveals 59 instances following suboptimal care. The exclusion of 10 mature asphyxiated infants without neonatal encephalopathy cannot be sustained: this assertion is close but not absolute, and it may be underdiagnosed.²

Suboptimal intrapartum care was recognised for 29% of all those investigated, and for either 46% or 49% of those in whom other identified causes were excluded. The criteria of suboptimal intrapartum care as defined in appendix 1 included delay in interpretation after the recognition of specified physical signs of "fetal distress" for more than 90 minutes, halved in the presence of meconium, but without any other qualification. These signs have been recognised as being ominous^{3,5}; the arbitrary time frame for response has been consistent delayed decelerations following three contractions for patients monitored by auscultation, and 10 to 15 minutes after ominous observations consistently recorded electronically.

Had stricter criteria been applied to the quality of intrapartum care there would almost certainly have been more instances of failure to respond to fetal distress. In Victoria, comprehensive data over many years have shown that fewer than 6% of infants are premature. The exclusion of 36% of patients with cerebral palsy from investigation because of prematurity, a significantly overrepresented group, further detracts from the validity of the study.

While the conclusions of this investigation of the contribution of suboptimal intrapartum care as a cause of cerebral palsy after either mature or all live births cannot be accepted, the quality and size of the epidemiological database is unique. A re-examination of the obstetrical antecedents of cerebral palsy, with the application of more appropriate standards for the response to fetal distress—especially relating to premature delivery, multiple pregnancy, and impaired fetal reserve before labour—would be invaluable.

E L G BEAVIS
Honorary consultant obstetrician

Alfred Hospital,
Melbourne,
Australia

- 1 Gaffney G, Sellers S, Flavell V, Squier M, Johnson A. Case-control study of intrapartum care, cerebral palsy, and perinatal death. *BMJ* 1994;308:743-50. (19 March.)
- 2 Chiswick ML. Intrapartum asphyxia and hypoxic-ischaemic encephalopathy. In: Spencer JAD, Ward RHT, eds. *Intrapartum fetal surveillance*. London: RCOG Press, 1993.
- 3 Haverkamp AD, Orleans M, Langendorfer S, McFee J, Murphy J, Thompson HE. A controlled trial of the differential effects of intrapartum fetal monitoring. *Am J Obstet Gynecol* 1979;134:399-408.
- 4 Haverkamp AD. Fetal monitoring in high risk labour. In: Spencer JAD, ed. *Fetal monitoring*. Oxford: Oxford University Press, 1991:207-10.
- 5 MacDonald D, Grant A, Sheridan-Pereira M, Boylan P, Chalmers I. The Dublin randomised trial of intrapartum fetal heart monitoring. *Am J Obstet Gynecol* 1985;152:524-39.

Authors' reply

EDITOR,—E L G Beavis's letter highlights the important fact that the term "cerebral palsy" is used to describe a group of conditions with different aetiological pathways.

As he correctly points out, among the recognised risk factors are preterm birth and multiple pregnancy. We, in collaboration with others, have been studying the factors which damage the developing brain in these groups of babies. Current evidence suggests that these factors can occur prenatally, during labour and delivery, or after birth. The extent to which "suboptimal care" during labour and delivery of the preterm baby contributes to the development of cerebral palsy is not known. As Beavis will be aware, there has been little systematic evaluation of different methods of management of preterm labour and delivery. Until current and new practice is rigorously tested for effectiveness