challenge to their professional omnipotence. In her classic paper Menzies notes how the ward environment and its procedures may be set up to reinforce defences that protect the staff working there from excessive anxiety to the detriment of patients' care.³

Individual professionals will find it difficult to work closely with dying patients unless they acknowledge their own emotional response to death and dying. Defences that leave their patients isolated will not be dismantled unless there is adequate and appropriate support provided. Until it is recognised that this support needs to be an integral part of both training and the weekly routine of the ward, patients dying in hospitals will die alone.

N RAMSAY
Senior registrar in psychiatry

Claybury Hospital,
Woodford Bridge,
Woodford Green,
Essex IG8 8BY

- 1 Mills M, Davies HTO, Macrae WA. Care of patients dying in hospital. *BMJ* 1994;309:583-6. (3 September.)
- 2 Ramsay N. Sitting close to death—a ward observation on a palliative care unit. *Group Analysis* (in press).
- 3 Menzies IEP. The functioning of social systems as a defence against anxiety. *Human Relations* 1960;13:95-121.

Technology foresight

EDITOR,—Richard Smith's interview with Sir Dai Rees draws attention to the technology foresight programme currently being run by the Office of Science and Technology and to its importance in determining the direction and future funding of medical research.¹ At the end of the interview Sir Dai states that the Medical Research Council has a portfolio of developments encompassing a range of biomedical and clinical sciences yet works with only a fairly narrow range of industries—pharmaceuticals and food. At least one other industry—the medical equipment industry—should have been mentioned in this context.

The contribution that the medical equipment industry makes to maintaining and improving the health and general quality of life of the British population is often overlooked and underrated. This is partly because the industry is dwarfed by the pharmaceutical and food industries and partly because it is composed of many small companies. Although the industry is generally profitable and maintains a net positive balance of trade, there is little slack to finance the development of the new, innovative products that are needed to maintain market position. The medical equipment industry needs to be revitalised if it is to take advantage of the anticipated outcomes of the technology foresight programme and so prevent the haemorrhage of British inventions to our competitors abroad. The development of mechanisms for achieving this will be an effective indicator of the success of the technology foresight programme.

PETER GREENAWAY
Member, health and life sciences panel,
technology foresight programme

Salisbury SP5 2AZ

- 1 Smith R. Developing foresight: an interview with Sir Dai Rees. *BMJ* 1994;309:324-6. (30 July.)

Use of fetal eggs for research

EDITOR,—In commenting on the public response to a consultation document published by the Human Fertilisation and Embryology Authority, Luisa Dillner correctly reports that 83% of responses were against the use of fetal eggs for infertility treatment but then says that "more than half the responses were in favour of using fetal eggs for research."¹ In fact, an overwhelming majority

was opposed to the use of fetal eggs in research: only 7.1% of the respondents were in favour and 58.1% said "No."² Dillner might also have mentioned that only 10.7% and 14.3% of the responses, respectively, were in favour of eggs from cadavers and live donors being used for research, while 29.2% and 24%, respectively, explicitly opposed this.

I am concerned that Dillner's inaccurate report may mislead readers about the strength of public feeling against these sinister developments. I am also concerned that the Human Fertilisation and Embryology Authority should "seek to increase the supply of eggs for embryo research" despite this massive vote against. Perhaps it will have cause to review this position in the light of the recommendation by the Council of Europe's steering committee on bioethics that "human embryos may not be created with the intention of conducting research on them,"³ although, as the authority notes, this is currently permitted (in Britain) under the Human Fertilisation and Embryology Act.

DAVID W EVANS
Retired consultant cardiologist

Cambridge CB3 9LN

- 1 Dillner L. Use of fetal eggs for infertility treatment is banned. *BMJ* 1994;309:289-90. (30 July.)
- 2 Human Fertilisation and Embryology Authority. *Donated ovarian tissue in embryo research and assisted conception*. London: HFEA, 1994.
- 3 Sheldon T. European experts produce draft on bioethics. *BMJ* 1994;309:221.

Results of prescription event monitoring study of salmeterol

EDITOR,—The prescription event monitoring study of salmeterol¹ is now complete and has shown the number of deaths due to asthma in patients treated with the drug to be lower than that predicted. Last year Castle *et al* reported on 16 787 patients treated with salmeterol for 16 weeks.² In these patients 54 deaths from all causes occurred (0.32%), including 12 from asthma (0.07%). The prescription event monitoring study has identified 1022 deaths (6.6%) in our cohort of 15 407 patients treated with salmeterol and observed for at least one year. Seventy three (7.1%) of the deaths (table I) were due to asthma, but only 39 of the 73 patients were taking salmeterol in the last month of life.

Our population was not only elderly (table II) but much more severely ill than that studied by Castle *et al*. For example, 37 of the 39 patients who died of asthma and were taking salmeterol in the last month of life were known to be receiving

TABLE II—Distribution of patients who died of asthma by age and sex. Figures are numbers (percentages)

Age (years)	Male patients	Female patients	All patients
<21	0	0	0
21-	1 (3)	1 (2)	2 (3)
31-	2 (6)	5 (12)	7 (10)
41-	4 (13)	3 (7)	7 (10)
51-	2 (6)	12 (29)	14 (19)
61-	12 (39)	12 (29)	24 (33)
71-	8 (26)	8 (19)	16 (22)
>80	2 (6)	1 (2)	3 (4)
Total	31	42	73
Average (SD) age	62.6 (14.6)	59.0 (14.2)	60.5 (14.4)

regular inhaled corticosteroids or oral corticosteroids, or both, before death.

We conclude that the death rate for asthma in our study (73/15 407; 0.47%) was greater than that found by Castle *et al* (12/16 787; 0.07%) because our study was at least three times longer than theirs and our population was both older and more severely ill. Nevertheless, the death rate from asthma in our study was substantially lower than that predicted¹ at the time of the report by Castle *et al*.

The overall death rate from asthma in the prescription event monitoring study reflects the age distribution of and severity of disease in the population that received salmeterol in the early years after its introduction. All of the deaths from respiratory causes in the study seemed on careful examination of the clinical records to have been due to natural causes, although in four of the 39 patients who took salmeterol in the last month of life there was a temporal relation between death and use of salmeterol; this relation does not, of course, define a causal association.

This study, which will be reported fully, elicited no evidence that salmeterol contributed to the deaths from asthma. One probable reason for the earlier overestimate of the number of deaths due to asthma that were likely to occur in this study¹ is the assumption that they would be evenly distributed throughout the study. In fact, 39 of the 73 deaths occurred in the first seven months of the study, which lasted for 25 months overall.

RONALD D MANN
Director

Drug Safety Research Unit,
Southampton SO13 1AA

- 1 Inman WHW. Bronchodilator treatment in asthma. *BMJ* 1993; 306:1610.
- 2 Castle W, Fuller R, Hall J, Palmer J. Serevent nationwide surveillance study: comparison of salmeterol with salbutamol in asthmatic patients who require regular bronchodilator treatment. *BMJ* 1993;306:1034-7.

TABLE I—Causes of death among 15 407 patients treated with salmeterol

Cause of death	Total No	Time from first prescription (months)			
		1-4	5-8	9-12	>12
Respiratory:					
Asthma	73	27	17	12	17
Chronic bronchitis	18	6	4	5	3
Bronchopneumonia	13	7	0	0	6
Chronic obstructive airways disease	362	92	71	82	117
Emphysema	32	11	8	8	5
Other	29	10	6	6	7
Cardiovascular:					
Aneurysm	19	4	8	4	3
Cardiac failure	17	5	4	6	2
Cerebrovascular accident	35	8	9	4	14
Ischaemic heart disease	177	48	34	35	60
Other	37	6	3	11	17
Malignancy:					
Carcinoma of the bronchus	62	10	6	23	23
Other	83	14	20	23	26
Gastrointestinal:					
Peptic ulcer	10	1	1	3	5
Other	13	2	4	2	5
Other:					
All other known causes	28	7	4	7	10
Cause uncertain	14	0	0	1	13
Total	1022	258	199	232	333