registers potentially places the psychiatrist and the key worker in an untenable position subject to a duty of care over a "patient" who probably does not want to see them and who may actively evade them. It may or may not be reasonably possible to know whether the patient is actually dangerous or at risk of suicide, but the duty of care may extend as long as the patient is on the register. As the guidelines cogently argue, the best predictor of the future is the past; but since many of the risk factors cited in the guidelines will remain in the patient's history, a decision to remove him or her from the register may be delayed notwithstanding a favourable change in circumstances.

See you in court

The key worker will find himself or herself under the spotlight alongside the consultant psychiatrist, and general practitioners also have their responsibilities for communication and effective action within the agreed care plan. The department may be right to claim that the introduction of supervision registers and detailed guidelines regarding their operation may not technically impose new legal or ethical responsibilities on these professionals, but they set comprehensive standards, such as the obligation to convene urgent multidisciplinary reassessments of the patient's status. Failure to meet the standards of care spelt out in the guidelines may well be cited in evidence in court.

New obligations are also placed on purchasers and providers of mental health care. For example, the guidelines specify that all staff "must be adequately trained in the care programme approach and in risk management and assessment." Having guidelines that require the provision of urgent multidisciplinary assessments is pointless if there are too few staff to implement them. The courts have held that a health authority providing a service is obliged to meet a reasonable standard of care in the provision of that service.⁶ Shortage of money will not be accepted as a defence by a health authority for failure to measure up to the required standard. How a court would decide these

matters when considering the complex issues surrounding duty of care toward vulnerable patients in the community is uncertain, but the registers (and the detailed guidance accompanying them) offer an invitation for aggrieved patients and relatives to find out.

The guidelines may still be modified after further discussions between the college and the Department of Health, but, whatever the final outcome, effective operational systems will be needed for the care of vulnerable people in the community. Pugh *et al* calculate that substantial numbers of psychiatric patients will need to be placed on registers according to the proposed criteria (p 611).⁷ These guidelines, together with those for the care programme approach, thus represent a substantial shift of resources within the mental health budget, and purchasers and providers will have to make the changes quickly and effectively. In some cases radical changes in working practices will be required to deliver the required standards of supervision, communication, and documentation. Harsh decisions may be required on priorities. Speed will be essential: civil litigation may soon ensure that any laggards catch up.

GLYNN HARRISON

Professor of community mental health

Department of Psychiatry,
University of Nottingham,
Mapperley Hospital,
Nottingham NG3 6AA

PETER BARTLETT

Lecturer in law

University of Nottingham,
University Park,
Nottingham

1 Supervision registers: the college's response. *Psychiatric Bulletin* 1994;18:385-6.

2 NHS Management Executive. *Introduction of supervision registers for mentally ill people from 1 April 1994*. Leeds: NHSME, 1994. (HSG(94)5.)

3 NHS Management Executive. *Guidance on the discharge of mentally disordered people and their continuing care in the community*. Leeds: NHSME, 1994. (HSG(94)27.)

4 Steering Committee of the Confidential Enquiry into Homicides and Suicides by Mentally Ill People. *A preliminary report on homicide*. London: SCCIHSMP, 1994.

5 Kingman S. Psychiatric patients who kill have often refused treatment. *BMJ* 1994;309:559.

6 *Wilsher v Essex Area Health Authority* [1986] 3 ALL ER 801 (CA).

7 Pugh R, Gardner J, Allen R. Implications of supervision registers in psychiatry. *BMJ* 1994;309:611.

Tiaprofenic acid and cystitis

Grounds for withdrawal?

Non-steroidal anti-inflammatory drugs are widely used and most of their adverse reactions are well known. The sites mainly affected by toxicity are the gut, the skin, and the kidney. Unusual toxic reactions, such as hepatotoxicity in the case of benoxaprofen or haematological toxicity in the case of phenylbutazone, have led to the drugs being withdrawn or their use severely restricted.^{1,2}

Clinical use of non-steroidal anti-inflammatory drugs should be based on their known adverse reactions, particularly their toxic effect on the gut. On this basis it is possible to construct a league table of risk: low dose ibuprofen is the safest drug, naproxen and diclofenac have intermediate risk, and piroxicam and azapropazone are the most toxic.³ Other adverse effects, such as the high rate of skin reactions with fenbufen, may also need to come into the equation.⁴

In 1991 Ahmed and Davison reported a case of cystitis in association with the non-steroidal anti-inflammatory

drug tiaprofenic acid.⁵ Cystitis has occasionally been reported through the Committee on Safety of Medicines' yellow card scheme with other non-steroidal anti-inflammatory drugs, but recent evidence from Australia has highlighted the risk with tiaprofenic acid.⁶ In three years' postmarketing surveillance in Australia 47 reports of cystitis associated with tiaprofenic acid were received out of a total of 71 urinary tract reactions to this drug. In contrast, only three reports of cystitis out of 74 urinary tract reactions to other non-steroidal anti-inflammatory drugs were received by the Australian adverse drug reaction scheme in its 25 year existence.

Similar evidence now comes from Britain in this week's *BMJ* in the form of two series of patients with cystitis induced by tiaprofenic acid—one from Manchester (eight patients), and one from Newcastle (three patients).^{7,8} Some of these patients had cystectomies, and, worryingly, two of the patients in the Manchester series had symptoms that

See articles on pp 574 and 599

did not resolve on stopping the drug. The Newcastle group includes data from the Committee on Safety of Medicines showing that cystitis and related symptoms are overrepresented among the British reports of adverse reaction to tiaprofenic acid compared with all other non-steroidal anti-inflammatory drugs.

Cystitis secondary to drugs other than cytotoxic drugs is an unusual problem. The cystitis associated with tiaprofenic acid seems likely to represent a direct toxic effect of the drug or its metabolites on the bladder. Tiaprofenic acid was being used for months or years in all the patients affected. Many of these patients were elderly, and their need for any long term treatment with a non-steroidal anti-inflammatory drug should be regularly reviewed. On present evidence long term treatment with tiaprofenic acid is potentially hazardous, and patients receiving this drug should be assessed for symptoms of cystitis, and if there are any the drug should be withdrawn.

The wider issue is whether a drug with such a potentially devastating effect should still be available, particularly

when its indications for use are shared by many other agents without this particular toxic effect. This must be an issue for drug regulatory authorities to consider. Once again spontaneous reporting by doctors on yellow card forms in Britain has been invaluable in assessing the toxicity of a drug. Doctors should be encouraged to send in reports of adverse reaction even when unsure of a causal association.

D N BATEMAN
Reader in therapeutics

Regional Drug and Therapeutics Centre,
Newcastle upon Tyne NE1 4LP

- 1 Smith A, Wright B. Benoxaprofen. *BMJ* 1982;285:459-60.
- 2 Phenylbutazone and oxyphenbutazone: time to call a halt. *Drug Ther Bull* 1984;22:5-6.
- 3 Bateman DN. NSAIDs: time to reevaluate drug toxicity. *Lancet* 1994;343:1051-2.
- 4 Brock PG, Jackson D. UK general practitioner experience of fenbrufen in elderly patients. *Eur J Rheumatol Inflamm* 1982;5:326-31.
- 5 Ahmed M, Davison ON. Severe cystitis associated with tiaprofenic acid. *BMJ* 1991;303:1376.
- 6 ADRAC. Update on tiaprofenic acid and urinary symptoms. *Australian Adverse Drug Reaction Bulletin* 1994;13:6.
- 7 Mayall FG, Blewitt RW, Staff WG. Tiaprofenic acid (Surgam) associated cystitis and ureteric obstruction. *BMJ* 1994;309:599-600.
- 8 Harrison WJ, Willis G, Neal DE. Adverse reactions to tiaprofenic acid (Surgam) can mimic interstitial cystitis. *BMJ* 1994;309:574.

The World Health Organisation in Africa

Too much politics, too little accountability

This month sees a crucial vote in international public health: the nomination of the World Health Organisation's next regional director for Africa. The appointment will affect how \$850m of public money is spent and what happens to the health of 550 million of the world's poorest people.

Many people are unhappy with what has been achieved since the current regional director, Dr Lobe Monekosso, took office 10 years ago. His three phase development plan for Africa has remained on the shelf, and only just over half of the region's \$136m budget for 1992-3 reached the region's 45 member states. The rest of the budget was spent at the regional office in Zaire, along with two thirds of the £112m donated for special programmes such as the AIDS and immunisation programmes.

Dr Monekosso has been unfortunate. Chronic difficulties caused by Africa's lack of infrastructure have been aggravated over the past year by civil unrest in Brazzaville, where the regional office is based. The office has been closed, and Dr Monekosso has had to decamp with some of his staff to his native Cameroon. The unrest prevented the WHO's auditors from evaluating biennial spending of \$122m, assets of \$20m, and liabilities of \$21m,¹ and left Dr Monekosso unable to counter suggestions of poor management voiced by, among others, Zambian delegates in an outspoken attack at this year's World Health Assembly.

Dr Monekosso's main rival for the post of regional director is Dr Ebrahim Samba, director of the onchocerciasis control programme. Dr Samba, from the Gambia, took over the ailing programme in 1980 and has turned it into a success. Funded by the World Bank and executed by the WHO, the programme has already achieved its 1995 target of eliminating onchocerciasis as a public health problem. From its base in Ouagadougou in Burkino Faso it has cut the prevalence of infection in 11 west African countries from 95% in 1974 to less than 5% today. Aerial pesticide spraying has allowed resettlement of 25 million hectares of fertile river basin that had been abandoned because of

infestation by blackfly. Donors' confidence has been maintained by clear results and tight financial management.

If the appointment of regional director was to be decided on a comparison of past performance Dr Monekosso would have reason to be worried. But it will be decided by the regional committee, which is made up of health ministers of the region's 45 member states, and observers say that, as in politics elsewhere, the sitting tenant will have the upper hand. According to a recent report, too much of regional directors' time is spent on regional politics.²

The fact that regional directors are answerable only to their regional committees and not to the WHO's governing body, the World Health Assembly, is increasingly seen as one of the WHO's major structural flaws. In theory the regional directors implement WHO policy. In practice they have complete autonomy within their domains, setting policy, appointing staff, and allocating resources.

Curtail regional directors' power

Observers say that, until recently, the regional directors' independence was tempered by the strong personality of the previous director general, Dr Halfden Mahler. His successor, Dr Hiroshi Nakajima, lacks his authority and communication skills. He is known to be in favour of limiting the regional directors' autonomy, and an internal United Nations report has now come to his aid.³ It recommends that regional directors' powers should be curtailed and that they should be selected by the director general after consultation with the regional committees. The European region, whose director, Dr Jo Asvall, is also up for reselection this month, has gone one step further. A search group has been set up to look for potential candidates and to assess those put forward by member states.

The internal report's recommendations will be discussed at the next assembly, but most observers doubt that the WHO will succeed in implementing them across all six regions. They think it highly unlikely that the Pan