intake of hindmilk and hence of vitamin K.⁴ If the baby needs extra vitamin K it can be given as a separate supplement.

Slattery makes some cogent points about the ethics of routine treatment of babies with vitamin K in the absence of controlled trials looking at possible side effects.^{1,5} Interpreting statistics on haemorrhagic disease of the newborn in relation to feeding and prophylactic treatment is extremely difficult. In the leaflets we have collected the incidence of haemorrhagic disease quoted to parents varies from 1 in 50 to 10 in 1.7 million babies. We welcome the developing debate on the size of the problem and the benefits and risks of the alternative preventive strategies. In particular, we value discussion about the appropriateness of a randomised controlled trial.

Despite the practical difficulties and the long wait for results² we believe that a trial is appropriate, given that the whole population of newborn babies is currently targeted. Careful attention should be paid to producing clear information based on research for all parents considering treatment for their child.

MARY NEWBURN
Head of policy research
ROSEMARY DODDS
Policy research officer

National Childbirth Trust,
London W3 8AR

- 1 Slattery J. Vitamin K for neonates. *BMJ* 1994;308:1635. (18 June.)
- 2 Draper G, McNinch A. Vitamin K for neonates: the controversy. *BMJ* 1994;308:867-8. (2 April.)
- 3 Von Kries R, Shearer PT, McCarthy M, Haug G, Harzer G, Gobel U. Vitamin K₁ content of maternal milk: influence of the stage of lactation, lipid composition, and vitamin K supplementation given to the mother. *Pediatr Res* 1987;22:513-7.
- 4 Woolridge MW, Baum JD. Infant appetite-control and the regulation of breast milk supply. *Children's Hospital Quarterly* 1991;3:113-9.
- 5 Slattery J. Why we need a clinical trial for vitamin K. *BMJ* 1994;308:908-10. (2 April.)

Treating hypertension after stroke

EDITOR,—Janice E O'Connell and Christopher S Gray's editorial highlights the uncertainty concerning the optimal management of hypertension in acute stroke.¹ I disagree with the authors over the treatment of hypertension after intracerebral haemorrhage. They advocate treating high systemic pressures (systolic >200 mm Hg; diastolic >120 mm Hg) and quote a paper by Lavin in support of this.² Lavin in fact suggested that intracranial pressure, rather than systemic blood pressure, should be reduced. Although he added that treatment of systemic hypertension may reduce cerebral oedema, he qualified this by suggesting that systemic hypertension should be reduced by no more than 20% and that ischaemia of the border zone may result. In response to a subsequent letter Lavin clarified that his personal bias was to lower intracranial pressure and not systemic blood pressure.³

A recent review of decisions on treatment in acute hypertension after stroke concluded that the role of hypotensive therapy in intracerebral haemorrhage remains unresolved and that, as with cerebral infarction, there are cogent reasons for believing that reduced systemic blood pressure may worsen cerebral blood flow, with resultant ischaemic damage.⁴

Although O'Connell and Gray state that no randomised trials looking specifically at antihypertensive treatment in acute stroke have been published, such a study has been carried out in ischaemic stroke.⁵ Although the study was small, no clinical benefit was observed, but it was suggested that nicardipine and possibly other calcium channel antagonists may impair cerebral blood flow.

I believe that, until robust evidence exists to support the use of antihypertensive drugs in acute stroke (either ischaemic or haemorrhagic) in adults, these drugs should be withheld initially unless hypertensive encephalopathy or aortic dissection is present.

RICHARD J DAVENPORT
Clinical research fellow

Neurosciences Trials Unit,
Department of Clinical Neurosciences,
University of Edinburgh,
Edinburgh EH4 2XU

- 1 O'Connell JE, Gray CS. Treating hypertension after stroke. *BMJ* 1994;308:1523-24. (11 June.)
- 2 Lavin P. Management of hypertension in patients with acute stroke. *Arch Intern Med* 1986;146:66-8.
- 3 Lavin P. Sodium nitropruside treatment in patients with acute strokes. *Arch Intern Med* 1986;146:1454.
- 4 Powers WJ. Acute hypertension after stroke: the scientific basis for treatment decisions. *Neurology* 1993;43:461-7.
- 5 Lisk DR, Grotta JC, Lamki LM, Tran HD, Taylor JW, Molony DA, et al. Should hypertension be treated after acute stroke? A randomized controlled trial using single photon emission computed tomography. *Arch Neurol* 1993;50:855-62.

Running an emergency helpline

The media may exploit doctors

EDITOR,—In their article on how to run an emergency helpline Cameron Stark and colleagues state, "If information can be adequately transmitted by the media a helpline should not be needed."¹ Our experience is that dealing with the media is not that simple.

We recently conducted a look back exercise after learning that a health care worker at a local hospital was positive for HIV. We had wanted to avoid unhelpful and prejudicial publicity caused by insensitive handling of the subject by the media, and so we avoided contact with the press for as long as possible. Patients who needed to be traced were approached personally through their general practitioners, and the tracing exercise was completed without press coverage.

The media inevitably heard of the incident. Unnecessary anxiety was generated among the general public, and a helpline was set up to answer concerns that the media raised. Over 2000 calls were received in the first 48 hours, and peaks of calls coincided with the broadcasting of news bulletins. The helpline proved a useful way of conveying information but was costly in terms of time and staff. There were also obvious conflicts of interest between us and the news hungry press, not the least of which concerned the confidentiality of the health care worker. Fortunately, we were able to maintain this.

The relationship between the health service and the press can be a valuable one, but once the media get hold of what they perceive to be a newsworthy health issue doctors need to make sure that they use the media to their advantage and are not exploited.

ALISON WALKER
Registrar in public health
NAUREEN BHATTI
Senior registrar in public health
MARK MCCARTHY
Director of public health

Camden and Islington Health Authority,
London NW1 2LJ

- 1 Stark C, Christie P, Marr AC. How to run an emergency helpline. *BMJ* 1994;309:44-6. (2 July.)

Guidance from NHS is required

EDITOR,—Cameron Stark and colleagues' article on how to run an emergency helpline described the actions necessary once the decision has been taken to establish such a helpline.¹ More pressing issues, however, are under what circumstances such

helplines are warranted and what constitutes such a serious risk to the patient that a "look back" exercise is needed. Over the past year such exercises have taken place in relation to helplines set up to provide advice on HIV and hepatitis B virus infection and unreliable results of cervical smear tests. In many cases the risk to patients may be remote, and the risk should be considered against the cost of such exercises and their tendency to provoke media alarm and public confusion.

The NHS Executive has no coherent overall view on look back exercises and no guidance for health authorities and trusts on the criteria for undertaking such initiatives and setting up helplines. Such guidance is needed to prevent unnecessary alarm among patients and to ensure that resources are directed towards such exercises only when there is clear risk to a large population of patients.

J B L HOWELL
Chairman

Board of Science and Education,
BMA,
London WC1H 9JP

- 1 Stark C, Christie P, Marr AC. How to run an emergency helpline. *BMJ* 1994;309:44-5. (2 July.)

Training in psychosexual medicine

EDITOR,—May Duddle's criticism of the Institute of Psychosexual Medicine's training is ill informed¹ as she has taken no part in the institute's training, research, and development, which have been carried out nationwide for 30 years. I wish to set the record straight.

Doctors accepted for basic training must expect to meet patients who expect examination of their sexual organs as well as emotional understanding; in general practice, patients come for family planning, genitourinary medicine, gynaecology, and andrology. Certain nurses, midwives, and obstetric physiotherapists also have to use intimate therapeutic techniques, and some join seminars with our accredited leaders. The clinical problems studied are thus wider than, though include, dysfunction in sexual performance. We expect our trainees to learn other techniques elsewhere: you would not expect to learn ear, nose, and throat surgery on an obstetric firm.

Training in seminars aims to convey a precise, extra psychosomatic skill (distinct from other non-directive verbal counselling and psychotherapy) to everyday consultations, whether they are for the overt presentation of sexual difficulty or marital discord; the fitting of a contraceptive or an intra-cavernosal injection; postmenopausal atrophy or a widower's impotence. In each case we examine less conscious aspects of the immediate doctor-patient interaction, considering what light they throw on other relationships. Secondly, we learn from the patients' reaction to, or anticipation of, genital examination something of their vision of their personal sexuality.

Duddle comments that we do not encourage partners to attend. Many young men with premature ejaculation or women with vaginismus are too anxious to be able to expose themselves to a partner, let alone to propose joining him or her in behaviour therapy. Furthermore, primary care doctors, unlike psychiatrists accepting referrals, must regard "the patient" as being whoever consults them. Much time in training concerns why doctors find themselves wishing to split up a couple who have elected to present together and, conversely, why in some cases doctors send for a partner they have never met, rather than noticing that they have a complaining or blaming patient before them.²

Finally, as an experienced psychotherapist Duddle will surely agree that the psychodynamics