We concluded that despite education and high motivation in an organised group of travellers simple preventive measures were not observed. Guidelines, however, are regularly updated,² and doctors advising travellers should be aware of current advice.

J H MILLER
Registrar

Military Wing,
Musgrave Park Hospital,
Belfast BFPO 801

D A WARRELL
Professor

Nuffield Department of Clinical Medicine,
John Radcliffe Hospital,
Oxford OX3 9DU

- 1 Miller JH, Byers M, Whiteoak R, Warrell DA. Imported falciparum malaria in British troops returning from Kenya. *J R Army Med Corps* (in press).
- 2 Bradley D. Prophylaxis against malaria for travellers from the United Kingdom. *BMJ* 1993;306:1247-52.

No documentary evidence of bribery by German hospital directors

EDITOR,—On behalf of the German Chamber of Physicians, I wish to comment on Helmut L Karcher's report on accusations of bribery against German hospital directors.¹ At the end of May the association of all German public health insurance companies said in a statement that directors of 50 out of 51 heart centres in western Germany were to blame for having accepted bribes from producers of artificial heart valves. The news magazine *Der Spiegel* published these accusations.²

In fact, no such nationwide scandal exists. When the public health insurance companies had to prove their accusations at a special meeting of the health committee of the German Bundestag on 1 June they claimed only 12 cases of bribery without naming the heart centres of hospital directors. Even three months later they were not able to support their accusations with documentary evidence.

There are no indications that bribery is common in German heart centres. According to the news magazine *Focus*, however, there are indications that the public health insurance companies themselves spent more than DM1m (£421 000) in an attempt to gain information about the alleged scandal from insiders.³ Karcher's article was written as though the story was true, but the story was a hoax.

ALEXANDER DÜCKERS
Head

Pressestelle der deutschen Ärzteschaft,
D-50931 Cologne,
Germany

- 1 Karcher HL. German hospital directors accused of bribery. *BMJ* 1994;309:1588-9. (18 June.)
- 2 Bares oder einen BMW. *Der Spiegel* 1994;22:92-8.
- 3 Kowalski M. Herzklappenaffäre: Flop für Krankenkassen. *Focus* 1994;31:18-21.

New treatments for benign prostatic hypertrophy

EDITOR,—Roger S Kirby and M C Bishop's examination of treatment of benign prostatic hypertrophy is timely,¹ given the growing number of methods of treatment that are being advocated and the emerging evidence that the volume of unmet need for treatment in the population is large.² We believe, however, that the arguments advanced by Kirby are not supported by the evidence. In the absence of data from randomised controlled trials his conclusions concerning alternative treatments can be only speculative. Unfortunately, despite continued uncertainty

about this issue, urologists in Britain have been reluctant to participate in a randomised controlled trial comparing transurethral resection of the prostate with open surgery.

Bishop argues that patients with mild lower urinary tract symptoms are best left alone, a view with which we concur.³ Apart from consideration of the balance of benefits and risks, the cost implications of advocating any of these innovative treatments for such men is enormous and the treatments should be undertaken only for good reason. There is therefore an urgent need to evaluate them properly. Before this is done, however, several issues that are insufficiently emphasised by Kirby and Bishop need to be resolved. Are the treatments seen as a substitute for transurethral resection of the prostate among patients currently offered treatment? Is it envisaged that they will be offered to patients with less severe disease than those currently being treated? What measure of outcome will be used, given the lack of concordance between symptoms and urinary flow?⁴ And as drug treatment may need to be continued for many years, what is the most appropriate time at which to measure outcome?

Kirby may be correct in saying that urologists unprepared to grasp the future will be left behind. The history of medicine is, however, littered with those who rushed to embrace new technologies only to abandon them when the initial enthusiasms proved unfounded.⁵ The question of who should pay also arises. At present health authorities and general practice fundholders should purchase these new treatments only if they are part of a randomised controlled trial.

MARTIN MCKEE
Senior lecturer

NICK BLACK
Reader

DUNCAN HUNTER
Research fellow

COLIN SANDERSON
Senior lecturer

Department of Public Health and Policy,
Health Services Research Unit,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

- 1 Kirby RS, Bishop MC. Are the days of transurethral resection of prostate for benign prostatic hyperplasia numbered? *BMJ* 1994;309:716-8. (17 September.)
- 2 Hunter DJW, McKee CM, Black NA, Sanderson CFB. Urinary symptoms: prevalence and severity in British men aged 55 and over. *J Epidemiol Community Health* (in press).
- 3 Doll HA, Black NA, McPherson K. Transurethral resection of the prostate for benign prostatic hypertrophy: factors associated with a successful outcome at one year. *Br J Urol* 1994;73:669-80.
- 4 Barry MJ, Cockett ATK, Holtgrewe HL, McConnell JD, Sihelnik SA, Winfield HN. Relationship of symptoms of prostatism to commonly used physiological and anatomic measures of the severity of benign prostatic hyperplasia. *J Urol* 1993;150:351-8.
- 5 Beeson P. Changes in medical therapy. *Medicine* 1980;59:79-84.

Supervised administration of methadone by pharmacists

EDITOR,—Supervision of the administration of methadone by community pharmacists has benefits beyond the prevention of "street leakage."¹ The risks of loss or theft of the drug, bingeing, injecting, and overdose are minimised; trust between doctors and patients is enhanced; and as a reward for progress in treatment patients may be given more than one day's supply to take home. A scheme similar to that in Glasgow¹ has run in West Glamorgan since 1991, with 25 pharmacists each supervising up to 15 patients. The anxieties expressed by pharmacists to Robert T A Scott and colleagues¹ have not been justified in practice.

A new scheme in Mid Glamorgan has been straightforward to implement. The local pharmaceutical committee agreed the protocol and

circulated it to all pharmacists. In both schemes scripts to be supervised are stamped to be easily identifiable. All patients newly prescribed methadone have their consumption supervised daily, Monday to Saturday. Pharmacists do not enforce the scheme but telephone the prescriber regarding non-compliance so that appropriate action can be taken.

P D Thomas's criticisms seem misplaced.² Both sets of guidelines referred to by Scott and colleagues emphasise the need for treatment and contact with a doctor over and above the prescription of methadone.^{3,4} It is perhaps surprising that neither refers to supervised consumption. The suggestion that prescribers should be responsible for supervision ignores both the unwillingness of many general practitioners even to prescribe and the number of patients per doctor seen by specialist services. Similar concerns over confidentiality and privacy were expressed when pharmacists first provided needle exchange, but pharmacists are now generally seen by drug misusers as "drug user friendly." The reasons for a prescription of methadone are understood by the pharmacist irrespective of where the drug is consumed. Three pharmacists in our schemes have rearranged their premises with privacy in mind, and others may follow. We would emphasise that pharmacists undertake supervision because they recognise the benefits for patients and the public; they receive only the dispensing fee.

Community pharmacists' supervision of consumption of methadone has many advantages. With collaboration between professional groups it can be straightforward to implement. It should be considered wherever centralised methadone maintenance clinics with supervised dispensing on site are impractical.

ANDREW J MCBRIDE
Consultant psychiatrist

IMAD MALI
Senior registrar

Llwyn yr Eos Clinic,
Church Village,
Mid Glamorgan CF38 1RN

ROSEMARY ATKINSON
Consultant psychiatrist

Central Clinic,
Swansea,
West Glamorgan

- 1 Scott RTA, Burnett SC, McNulty H. Supervised administration of methadone by pharmacists. *BMJ* 1994;308:1438. (28 May.)
- 2 Thomas PD. Doctors should supervise administration of methadone. *BMJ* 1994;309:53-4. (2 July.)
- 3 Advisory Council on the Misuse of Drugs. *AIDS and drug misuse update*. London: HMSO, 1993:42-9.
- 4 Department of Health, Scottish Home and Health Department, Welsh Office. *Drug misuse and dependence: guidelines on clinical management*. London: HMSO, 1991.

Computer training for doctors and students

EDITOR,—We agree with the comments by Ronald LaPorte and colleagues¹ and Ray Jones and Sue Kinn.² The information superhighway, an important technical development of immense potential impact in clinical medicine, will not fulfil its potential if doctors do not, will not, or cannot use it because they do not know how. Education is clearly the key—a fact recognised by the General Medical Council, which has stated that "a working knowledge of modern medical information technology will be essential to the doctor of the future."³

For some years our unit has run a course in information technology as part of the first year curriculum at Leeds University Medical School, teaching skills necessary to use computers and networks (accessing network information, word processing, use of spreadsheets and databases) and emphasising the use of computers as a source of information for help with practical problems.