The second constraint was the need for mobility, especially problematic with the poor housing market, and dual career households. Marriage was seen as a constraint for both men and women, to a much greater extent than were children. Seemingly, it is the effects of the working conditions on the family that affect choice of career rather than the family alone affecting it. This goes right down to the day to day detail of how hard it is to keep a nanny when you work such dreadful hours. Sadly, the women in the study see this as a problem for women rather than a problem for parents.

Senior doctors come in for their usual share of criticism in terms of the effects they have on women's careers in particular. In the words of one young woman: "I was very ambitious at first and it made me want to do gynaecology because I felt they needed more women in the specialty. But then I realised that I didn't want to spend the rest of my life on the defensive." So she became a general practitioner. Other quotes speak of humiliation right through medical school and on—for example, "Our consultant renal physician . . . said he couldn't see why women tried to do medicine as they'd only go off to 'breed some time, like an old cow' "; and "One of the professors said, 'Why are we wasting time on you when you'll end up washing nappies?' " Such an irrational remark, contrary to all the evidence, sounds more like wishful thinking to me. A locum in general practice noted, "They tended to favour women for junior posts because they're more conscientious and work harder. But they didn't see them as career doctors."

Clearly, throughout medical school and their early post-graduate years these doctors are inspired by good examples from seniors and appalled and driven running from those who humiliate and bully. Although this occurs for both men and women, for women the remarks tend to strike at their core in terms of gender—something that will inevitably be demoralising for all but those with the thickest of skins. Certainly women emerge as more self critical than men,⁶ and this in itself is likely to stop them applying

for more senior posts.

This bantering (at best) and abuse (at worst) may undermine the foundations of any career ladder, so nothing more than the first rungs are attempted. But many of those with the confidence to go for higher jobs are still met with selection processes, often blatantly illegal, that ensure that they climb no further.

It's a grim picture of little or no change, and in some cases an actual deterioration, in what might make things better for these young women and men. People have likened changing the NHS to changing course in an ocean liner; it seems to me that changing medicine is more akin to trying to turn an iceberg. So many of the blocks and dangers are hidden under the surface. There are the mystique and initiation rites that come secretly with any profession; the psyches of some consultants that suggest little better than misogyny; and the women who so often blame themselves when things go wrong and complain politely if at all. These factors and more need researching and addressing—and not just in medicine—if any fundamental change is to take place.

Allen produces a useful list of recommendations. For me the glints of light in the study come mainly from the young men, many of whom now see that what is happening to them and to women is wrong. The most hopeful trend that Allen's data show is the increasing recognition by these doctors of discrimination by some seniors. If they can just hold on to that, things might eventually change.

JENNY FIRTH-COZENS
Principal research fellow

Department of Psychology,
University of Leeds,
Leeds LS2 9JT

- 1 Skentelbery D. Junior doctor died of natural causes says coroner. *BMJ* 1994;309:1530.
- 2 Allen I. *Doctors and their careers: a new generation*. London: Policy Studies Institute, 1994.
- 3 Dillner L. Doctors are more miserable than ever, says report. *BMJ* 1994;309:1529.
- 4 Allen I. *Doctors and their careers*. London: Policy Studies Institute, 1988.
- 5 NHS Management Executive. *Women in the NHS: an implementation guide to opportunity 2000*. London: NHSME, 1992.
- 6 Firth-Cozens J. Women doctors. In: Firth-Cozens J, West M, eds. *Women at work: psychological and organizational perspectives*. Buckingham: Open University Press, 1991:131-42.

Diagnosing pulmonary embolism

D-dimer needs rigorous evaluation

All doctors with patients suspected of having an acute deep venous thrombosis or a pulmonary embolism would love a diagnostic magic bullet. They long for a test that would rule in or rule out the diagnosis of venous thromboembolism with a high degree of certainty. Ideally, such a test would be rapid, non-invasive, and widely available.

The search for a reliable test has continued because the clinical diagnosis of thromboembolism is neither sensitive nor specific. Neither symptoms nor examination of the legs are reliable; nor can clinical signs or symptoms define the presence or absence of pulmonary embolism. Clinical signs and symptoms can only suggest these diagnoses—and asymptomatic venous thrombosis and pulmonary embolism are both fairly common.

Not surprisingly, then, over the years a succession of simple, relatively non-invasive tests has been explored. Blood tests have included measurement of the "diagnostic" triad of lactic dehydrogenase, bilirubin, and glutamic oxaloacetic transaminase; immunoglobulin E¹; arterial oxygen and the alveoloarterial oxygen gradient; and

fibrinopeptide A²—and others have been proposed.^{3,4} Lung physiologists have suggested the use of measurements of dead space and tidal volume. Radionuclide venography enjoyed a brief career.

For various reasons each approach followed a parabolic course from the rapid ascendancy of initial excitement to rapid descent as its value failed to be validated. Still in play are techniques that use radiolabelled components of thrombus—such as platelets and monoclonal antibodies to fibrin. Radioactively labelled fibrinogen, which provided invaluable epidemiologic data, is no longer seen as the answer to the diagnosis of venous thrombosis.⁵ But computed tomography and magnetic resonance imaging are still being investigated.⁶

As this galaxy of non-invasive tests has moved across the horizon a new and promising entrant has appeared: measurement of D-dimer. D-dimer is a specific degradation product of crosslinked fibrin that is released when the endogenous fibrinolytic system attacks the fibrin matrix of fresh venous thromboemboli. Many recent research papers

have made claims for the value of measuring D-dimer concentrations in patients suspected of having venous thromboembolism.^{7,8}

One issue has been clarified: raised concentrations of D-dimer should not be interpreted as supporting the diagnosis of deep vein thrombosis or pulmonary embolism. The test is non-specific; raised values can be found in many other conditions and, indeed, in the postoperative period, a high risk time for thromboembolism. On the other hand, measurements of the concentration of D-dimer may be used to exclude the diagnosis—that is, the absence of a raised concentration implies that there is no fresh thromboembolic material undergoing dissolution in the deep veins or in the pulmonary arterial tree. Such a (nearly) exclusionary test would be valuable, especially when combined with other well studied non-invasive tests for deep vein thrombosis (impedance plethysmography, ultrasonography) and pulmonary embolism (lung scanning).

Will this promise be fulfilled? It is still too early to tell; indeed, we are some distance from knowing. Too many assays have been used in different studies—at least four enzyme linked immunosorbent assays (ELISAs) and four latex assays. Normal values differ for each: as Bounameaux *et al* recently said, standardisation and calibration are urgently needed.⁹ Furthermore, any test born in the tightly controlled incubator of research tends to slide a bit in performance when it comes into general use.

Other issues need to be addressed. Normal values tend to vary with age. How long does it take for a deep vein thrombosis or pulmonary embolism to become sufficiently organised that it no longer releases D-dimer? How severely will the value of the test be reduced by all of the competing causes for a raised D-dimer concentration in patients in hospital?

Venous thromboembolism is potentially lethal, so the margin of tolerable error is not wide. D-dimer has not yet been put to the ultimate test—a trial in which treatment is withheld from patients suspected of having deep vein

thrombosis or a pulmonary embolism with D-dimer concentrations below a specified cut off value. Such trials of outcome should take account of the existing diagnostic approaches that have already gone through such trials—impedance plethysmography and ultrasonography¹⁰ for detection of deep vein thrombosis and perfusion ventilation scanning^{11,12} and pulmonary angiography for exclusion of pulmonary embolism. In these instances, the available data indicate that treatment may be withheld without adverse outcomes.

What is the way ahead? After standardisation and calibration, assays for D-dimer must be subjected to the definitive study: they should be combined with other established tests to support decisions not to treat. Let us hope that D-dimer will not descend the parabola of other diagnostic blood tests, but only time and further investigations will tell.

KENNETH M MOSER
Professor of medicine

UCSO Medical Center,
San Diego, CA 92103-8372, USA

- 1 Takekawa H, Miyamoto K, Yamaguchi E, Munakata M, Kawakami Y. Acute rise in serum immunoglobulin E concentration in pulmonary thromboembolism. *Chest* 1993;104:61-4.
- 2 Tulchinsky M, Zeller JA, Reba RC. Urinary fibrinogen A in evaluation of patients with suspected acute pulmonary embolism. *Chest* 1991;100:394-8.
- 3 Walsh SJ. Diagnostic accuracy and deep venous thrombosis. *J Nucl Med* 1991;12:2328-31.
- 4 Moser KM. Venous thromboembolism: state of the art. *Am Rev Respir Dis* 1990;141:235-49.
- 5 Kanke M, Matsueda GR, Strauss HW, Yasuda T, Liau CS, Khaw BA, *et al*. Localization and visualization of pulmonary emboli with a radiolabeled fibrin-specific monoclonal antibody. *J Nucl Med* 1991;32:1254-69.
- 6 Spritzer CE, Norcup JJ Jr, Sostman HD, Coleman RE. Detection of deep venous thrombosis by magnetic resonance imaging. *Chest* 1993;104:54-60.
- 7 Bounameaux H, Cirafo P, de Moerloose P, Slosman D, Schneider PA, Reber G, *et al*. Measurements of D-dimer in plasma as diagnostic aid in suspected pulmonary embolism. *Lancet* 1991;337:196-200.
- 8 Goldhaber SZ, Simons GB, Elliot GG, Haire WD, Toltzis R. Quantitative plasma D-dimer levels among patients undergoing pulmonary angiography for suspected pulmonary embolism. *JAMA* 1993;270:2819-22.
- 9 Bounameaux H, de Moerloose P, Perrier A, Reber G. Plasma measurements of D-dimer as diagnostic aid in suspected venous thromboembolism: an overview. *Thromb Haemostas* 1994;71:1-6.
- 10 Heijboer H, Buller H, Lensing AWA, Turpie A, Colly LP, Wouter Ten Cate J. A comparison of real-time compression ultra sonography with impedance plethysmography for the diagnosis of deep-vein thrombosis in symptomatic outpatients. *N Engl J Med* 1993;329:1365-9.
- 11 Hull R, Raskob GE, Coates G. Clinical validity of a normal perfusion lung scan in patients with suspected pulmonary embolism. *Chest* 1990;97:23-6.
- 12 Kipper MS, Kortman KE, Ashburn WL, Moser KM. Long-term follow-up of patients with suspected pulmonary embolism and a normal scan. *Chest* 1982;82:411-5.

Botulinum toxin in the cerebral palsies

Grounds for cautious optimism

Botulinum toxin is now routinely used in adult focal dystonias,¹ including blepharospasm,² spasmodic torticollis,³ and hemifacial spasm, and also in squint and some types of tremor.⁴ When injected close to the main area of nerve arborisation within a muscle, the toxin is selectively taken up by cholinergic nerve terminals and blocks neuromuscular transmission for between two and four months until new terminals grow. Recent reports of its use in the cerebral palsies are encouraging but need to be placed in context.

The cerebral palsies are motor disorders arising from static brain abnormalities that may result from early embryological defects or damage in utero or early extra-uterine life. Cerebral palsy phenotypes often include primary cognitive, psychiatric, sensory, and seizure disorders and a wide range of secondary skeletal and general medical illnesses. The motor disorders vary greatly in severity and neurological signs. Each pathology may therefore cause a range of types of cerebral palsy without any necessary concordance between pathological and clinical categories.⁵

The process of progressive developmental revelation with age, the cascade of disabilities in the first two to three decades, and the propensity of the developing nervous system to compensate for early damage (albeit by its own efforts⁶ rather than those of doctors and therapists) have prompted a drive for early intervention. Very early definition of motor disorders and even of the population at risk may result in many babies with transient abnormalities being treated, who may then feature as "cured." Until now, however, no solid evidence of the medical cure of a cerebral palsy syndrome has been published, and the problems of running good randomised studies in such a heterogeneous population are clear.

As we move from global treatments (early management⁷) through regional approaches (such as dorsal rhizotomy⁸) to selective interventions in which one or two muscle groups have their tone reduced, we can more easily define problems in objective terms and conduct trials likely to yield statistically valid conclusions. That does not preclude benefit from less selective approaches; it merely means