ROLE OF SYSTEMIC TREATMENT FOR PRIMARY OPERABLE BREAST CANCER

M A Richards, I E Smith, J M Dixon

Survival without relapse of patients with operable breast cancer (data from Guy's Hospital, London).

Outline of options for systematic treatment of large, operable breast cancer.

Advantages and disadvantages of adjuvant and primary systemic treatment

Systemic treatment	Advantages	Disadvantages
Adjuvant	Proved efficacy Prognostic information available after surgery	Uncertainty whether treatment is effective in individual patients
Primary	Allows direct assessment of effectiveness Tumour shrinkage may allow breast conservation	Loss of prognostic information May treat in situ disease (if diagnosis made by fine needle aspiration cytology alone)

Over half of women with operable breast cancer who receive locoregional treatment alone (surgery with or without radiotherapy) will die from metastatic disease, indicating that for most women the cancer has already spread by the time of presentation. The major risk factors for development of metastases are axillary lymph nodes being involved, an adverse histological grade (indicating an undifferentiated cancer), and large tumour size. Combinations of these factors can be used to define groups with widely different risks of relapse: from less than 10% to more than 90% remaining free of disease after five years. The only way to improve the chance of survival for many of these women is to give them effective systemic treatment.

Systemic treatment may be given either as adjuvant treatment after locoregional treatment or as primary systemic treatment before locoregional treatment. The effectiveness of adjuvant treatment has been clearly shown in clinical trials, while primary systemic treatment is still being evaluated. A problem with adjuvant treatment is that its effectiveness in individual patients cannot be assessed as there is no overt disease to monitor. In contrast the effectiveness of primary medical treatment can be assessed by monitoring the size of the primary tumour. Primary medical treatment in operable breast cancer can result in shrinkage of a tumour; thus, a large tumour initially treatable only by mastectomy can be made suitable for breast conserving treatment.

One potential problem with primary systemic treatment is that, if the diagnosis of cancer is made by fine needle aspiration cytology alone, in situ disease could be treated by chemotherapy (cytology cannot differentiate invasive and in situ disease). For this reason most units perform core or incisional biopsy to obtain a histological diagnosis of invasive cancer before embarking on primary medical treatment. There is no evidence that leaving a primary tumour in the breast during primary systemic treatment increases patient's anxiety.

Improvements in 10 year survival of women with stage II breast cancer associated with different treatments*

Treatment	Proportional reduction in annual mortality (SD)	Absolute reduction in 10 year mortality per 100 women treated (SD)
Women aged ≥ 50		
Polychemotherapy† only	12% (4%)	5 (2)
Tamoxifen only (for median of 2 years)	20% (2%)	8 (1)
Polychemotherapy† and tamoxifen	30% (5%)	12 (2)
Women aged < 50		
Polychemotherapy† only	25% (5%)	10 (3)
Ovarian ablation only	28% (9%)	11 (4)
Polychemotherapy† and ovarian ablation‡	30-40%	12-16

*Absolute benefit may be about half as great for stage I cancer.

†For example, > 6 months of cyclophosphamide, methotrexate, and fluorouracil.

‡Estimates unreliable because of small amount of data.

Polychemotherapy, oophorectomy, and treatment with tamoxifen produce significant reductions in the annual rates of tumour recurrence and of death. These treatments have been shown to affect survival for at least 10 years and probably longer. The treatments are proportionally as effective for women at high risk of relapse (such as node positive patients) as for women at lower risk, but absolute reduction in mortality is much greater for patients at high risk.

Chemotherapy

From results of clinical trials it is clear that

- A combination of drugs is more effective than a single drug
- A single (perioperative) course of treatment is less effective than a more prolonged course of treatment (six months), but there seems to be no advantage in giving treatment for longer than six months
- The benefits of chemotherapy are greatest in women aged under 50; a smaller but still significant benefit is seen in older women.

Side effects of drugs used for adjuvant treatment

Chemotherapy

- Fatigue and lethargy
- Alopecia (temporary)
- Nausea and vomiting
- Induction of menopause
- Specific side effects of certain drugs
- Risk of infection
- Oral mucositis
- Diarrhoea
- Weight gain

Oophorectomy

- Induction of menopause
- Vaginal dryness
- Hot flushes
- Osteoporosis

Gonadotrophin releasing hormone analogues

- As for oophorectomy
- Pain and bruising at injection site

Tamoxifen

- Hot flushes
- Altered libido
- Gastrointestinal upset
- Visual disturbances—rare (stop drug if reported)
- Vaginal dryness
- Menstrual disturbance
- Weight gain

Antiemetic regimens during chemotherapy

Moderately emetogenic chemotherapy*

- Intravenous dexamethasone before and after chemotherapy, intravenous metoclopramide, and oral dexamethasone and metoclopramide to take home
- If previous vomiting when given chemotherapy or if patient aged < 45 consider intravenous ondansetron 8 mg or intravenous granisetron 3 mg before chemotherapy and intravenous dexamethasone 8-16 mg

Highly emetogenic chemotherapy†

- Intravenous ondansetron 8 mg or intravenous granisetron 3 mg, intravenous dexamethasone, and oral dexamethasone to take home

*For example, cyclophosphamide, methotrexate, and fluorouracil.

†Regimens containing high doses of doxorubicin and cisplatin.

Oophorectomy

- Is only of benefit in women aged under 50 (premenopausal)
- Produces survival benefits of similar magnitude to those obtained by polychemotherapy in such younger women
- Can be achieved surgically, by radiation, or by administration of analogues of gonadotrophin releasing hormone.

Tamoxifen

- 20 mg a day is as effective as higher doses
- One year of treatment is less effective than two or more years, and five years may be better than two years
- Reduces the risk of contralateral breast cancer by about 40%
- Benefits are greatest in patients with tumours rich in oestrogen receptors, but smaller reductions in mortality still occur in patients with tumours poor in oestrogen receptors
- Is effective in all age groups

Side effects

Although hair loss is the most common concern of patients before starting chemotherapy, 80% report fatigue and lethargy as the most troublesome side effect. The occurrence of alopecia with some chemotherapy regimens may be reduced by scalp cooling. Nausea and vomiting are unpleasant side effects but can be controlled in most patients by appropriate antiemetic drugs. Younger patients seem to be more at risk of nausea and vomiting and are more likely to suffer from extrapyramidal side effects from standard antiemetic regimens including metoclopramide. It is thus appropriate to give serotonin-3 (5-HT₃) antagonists to women aged under 45 as first line treatment even for moderately emetogenic chemotherapy.

The side effects of hormonal treatment are greatest in premenopausal patients. Only 3% of patients given tamoxifen stop taking the drug because of side effects, but vaginal dryness, loss of libido, and hot flushes can have a considerable impact on quality of life. First line treatment for vaginal dryness is with a non-hormonal cream (such as Replens), but if this is not effective creams containing an oestrogen (such as dienoestrol cream) should be tried. Clonidine is occasionally effective at relieving flushing. Evening primrose oil has been reported anecdotally to improve menopausal symptoms, but in controlled trials it has not been shown to be of benefit. Weight gain can be a problem with tamoxifen and with chemotherapy.

Selection of adjuvant treatment

Definitions of risk groups and associated risk of relapse

Risk group	Definition	Survival without relapse after 5 years
Node negative patients		
• Low risk	Tumour ≤ 1 cm in diameter	>90%
• Intermediate risk	Tumour >cm, grade I or II	75-80%
• High risk	Tumour >cm, grade III	50-60%
Node positive patients		
• Low and intermediate risk	1-3 axillary nodes involved	40-50%
• High risk	4-9 axillary nodes involved	20-30%
• Very high risk	≥ 10 axillary nodes involved	10-15%

Common adjuvant chemotherapy regimens

- Cyclophosphamide, methotrexate, and fluorouracil; repeated every 21-28 days
- Doxorubicin and cyclophosphamide; repeated every 21 days (fluorouracil may be added to regimen, and epirubicin may be substituted for doxorubicin)

Which chemotherapy regimen?

Cyclophosphamide, methotrexate, and fluorouracil is the most widely used regimen in the United Kingdom. Whether regimens containing anthracyclines (such as doxorubicin and epirubicin) are superior remains controversial. Some units give regimens containing doxorubicin to premenopausal women at high risk of relapse. Four cycles of doxorubicin and cyclophosphamide seem to be as effective as six cycles of cyclophosphamide, methotrexate, and fluorouracil.

Combinations of chemotherapy and hormonal treatment

For women aged over 50 the effects of tamoxifen and chemotherapy seem to be additive. Tamoxifen, however, has a considerably greater effect on mortality than does chemotherapy in these older, postmenopausal women and less toxicity. Whether the additional benefit justifies the use of chemotherapy in these patients is a matter for individual judgment by both patient and doctor. The effects of combining chemotherapy and hormonal treatment in women aged under 50 are unclear. In many centres patients are entered into trials of adjuvant treatment, which may therefore not follow that outlined here.

Patients at very high risk

Protocols of intensive treatment are being assessed in patients at very high risk. In the United States women with 10 or more positive axillary lymph nodes are offered high dose chemotherapy with autologous transplantation of bone marrow or harvesting and subsequent rescue of peripheral stem cells. Early results show mortality related to treatment is controllable (around 5%), but morbidity is high. Comparison with historical controls suggests a prolongation in time to relapse, and preliminary evidence indicates a survival benefit. Randomised controlled clinical trials are currently under way, and results from these studies are required before this treatment is introduced in all units. In Britain this approach is being piloted in a few specialist units.

Adjuvant treatment for patients with breast cancer

Risk group	Premenopausal patients	Postmenopausal patients
Node negative patients		
• Low risk	Tamoxifen or no treatment	Tamoxifen or no treatment
• Intermediate risk	Tamoxifen	Tamoxifen
• High risk	Consider chemotherapy* (with or without tamoxifen) or Ovarian ablation (with or without tamoxifen) if tumour is oestrogen receptor positive	Tamoxifen (with or without chemotherapy)
Node positive patients		
• Low and intermediate risk	Chemotherapy* (with or without tamoxifen) or Ovarian ablation (with or without tamoxifen) if tumour is oestrogen receptor positive or Chemotherapy* and ovarian ablation (with or without tamoxifen)	Tamoxifen with or without chemotherapy
• High and very high risk	Consider more intensive chemotherapy† (with or without tamoxifen)	Tamoxifen and chemotherapy if fit

*For example, cyclophosphamide, methotrexate, and fluorouracil.

†For example, regimen containing anthracycline. Some units are piloting use of intensive chemotherapy supported by rescue of bone marrow or peripheral stem cells for patients at very high risk.

Primary medical treatment

World Health Organisation's definition of objective response

- **Complete clinical response**
Disappearance of palpable disease
- **Partial response**
Decrease of $\geq 50\%$ in total size of tumour
- **No change**
Decrease of $< 50\%$ or increase of $< 25\%$ in total size of tumour
- **Progressive disease**
Increase of $\geq 25\%$ in total size of palpable lesion

Mammograms of breast showing primary cancer and involved axillary node (left) and after primary chemotherapy (right). At subsequent surgery patient was found to have no residual carcinoma in breast or axilla—complete pathological response.

Serial mammograms during primary treatment with systemic chemotherapy. Mass lesion disappeared, but microcalcification remained; subsequent mastectomy showed that all microcalcification was associated with residual carcinoma in situ.

Serial ultrasound scans at fortnightly intervals during primary treatment with chemotherapy.

J M Dixon acknowledges the support of the Cancer Research Campaign. Photographs were prepared by Mr D Dirom, Medical Illustration Unit, University of Edinburgh. The data presented in the box of improvements in survival associated with different treatments are from Early Breast Cancer Trialists' Collaborative Group, *Lancet* 1992;339:71-85 and are reproduced with permission of the journal.

The use of primary medical (neoadjuvant) treatment for operable breast cancer has increased over the past five years. The main advantages to this approach have already been outlined. Accurate assessment of response is important to determine whether the systemic treatment is effective. Clinical response is assessed according to criteria of the International Union Against Cancer (UICC); mammographic or ultrasonographic assessments of response may also be useful. Both the primary tumour and lymph node metastases can be shown to respond, and invasive cancer seems to be more sensitive to chemotherapy than in situ disease.

Chemotherapy

With conventional regimens for primary systemic chemotherapy, about 70% of patients show a partial response (primary cancer shrinking by $> 50\%$), 20-30% show a complete clinical response, and a small number (about 10-15%) achieve a complete pathological response. The regimens used for primary chemotherapy have generally been the same as those used for adjuvant treatment. There is preliminary evidence that continuous infusional chemotherapy with agents such as fluorouracil combined with intermittent agents such as epirubicin and cisplatin achieves higher response rates (over 90%) than bolus chemotherapy.

Hormonal treatments

Tamoxifen (20 mg a day by mouth) produces a partial response in 75% of elderly patients with hormone responsive (oestrogen receptor positive) tumours and a complete clinical response in 15%. The use of gonadotrophin releasing hormone analogues (goserelin 3.6 mg monthly given subcutaneously or leuporelin 3.75 mg monthly given subcutaneously or intramuscularly) as primary medical treatment for premenopausal women with oestrogen receptor positive tumours is under evaluation in some centres. Few patients show complete pathological responses after hormonal treatment, but the side effects are generally much less than with chemotherapy.

Outcome

Tumours will show either a useful response or further progression within 12 weeks of primary medical treatment. To ensure that women with progressive disease are detected as early as possible, clinical and ultrasonographic assessment of tumour volume should be performed at monthly intervals.

Between 50% and 70% of patients with large tumours will have sufficient tumour regression to avoid mastectomy. All patients require some form of local treatment (surgery or radiotherapy) after primary systemic treatment. It is not yet clear whether primary medical treatment prolongs survival, and a series of randomised trials are under way to assess this.

Selection of patients

Primary systemic treatment was initially given to patients with locally advanced (inoperable) breast cancers, and its use has now been extended to patients with large operable breast cancers in an attempt to avoid mastectomy. The use of primary systemic treatment for other groups of patients cannot be recommended (except in clinical trials) until results are available from trials comparing it with standard treatments.

M A Richards is reader in medical oncology at ICRF Clinical Oncology Unit, Guy's Hospital, London; I E Smith is consultant medical oncologist at Royal Marsden Hospital, London; and J M Dixon is senior lecturer in surgery at Edinburgh Royal Infirmary and honorary consultant surgeon at Western General Hospital, Edinburgh.

The ABC of Breast Diseases has been edited by J M Dixon.