symptoms when reintroduced into their diet. This practical approach to food intolerance supports earlier more rigorous studies in which double blind challenges with food instilled down nasogastric tubes were proved to reproduce symptoms.¹⁴

About 10% of all patients with the irritable bowel syndrome develop symptoms abruptly after an acute bout of diarrhoea.¹⁵ This postdysenteric irritable bowel syndrome has a good prognosis, most patients being free of symptoms within five years.¹⁵ The mechanism is uncertain, but in a small number of patients malabsorption of bile salts after salmonella ileitis may be responsible.¹⁶ The resulting bile salt catharsis responds well to cholestyramine, aluminium hydroxide, or, more palatably, loperamide.

With a clearer, more restrictive definition of the irritable bowel syndrome that excludes people with a predominantly psychiatric problem, the success of management has improved. Doctors treating such patients with reassurance and dietary advice (mainly to increase fibre content) together with bulking agents such as ispaghula and where necessary antispasmodics or antidiarrhoeal drugs can expect 85% of patients at six weeks and 68% after five years to be free of symptoms or have only minor symptoms.¹⁵ Only a few patients need the more intensive reassurance provided by hypnotherapy or other forms of psychotherapy.

1 Drossman DA, Thompson WG, Talley NJ, Funch-Jensen P, Janssens J, Whitehead WE. Identification of subgroups of functional gastrointestinal disorders. *Gastroenterol Int* 1990;3:159-72.

- 2 Sandler RS, Drossman DA, Nathan HP, McKee DC. Symptom complaints and health care seeking behavior in subjects with bowel dysfunction. *Gastroenterology* 1984;87:314-8.
- 3 Smith RC, Greenbaum DS, Vancouver JB, Henry RC, Reinhardt MA, Greenbaum RB, et al. Psychological factors are associated with health care seeking rather than diagnosis in irritable bowel syndrome. *Gastroenterology* 1990;98:293-301.
- 4 Drossman DA, McKee DC, Sandler RS, Mitchell CM, Cramer EM, Lowman BC, et al. Psychosocial factors in the irritable bowel syndrome. A multivariate study of patients and nonpatients with irritable bowel syndrome. *Gastroenterology* 1988;95:701-8.
- 5 Whitehead WE, Bosmajian L, Zonderman AB, Costa PT, Schuster MM. Symptoms of psychological distress associated with irritable bowel syndrome. Comparison of community and medical clinic samples. *Gastroenterology* 1988;95:709-14.
- 6 Thompson DG, Laidlow JM, Wingate DL. Abnormal small bowel motility demonstrated by radiotelemetry in a patient with irritable colon. *Lancet* 1979;ii:1321-3.
- 7 Kellow JE, Phillips SF. Altered small bowel motility in irritable bowel syndrome is correlated with symptoms. *Gastroenterology* 1987;92:1885-93.
- 8 Kellow JE, Gill RC, Wingate DL. Prolonged ambulant recordings of small bowel motility demonstrate abnormalities in the irritable bowel syndrome. *Gastroenterology* 1990;98:1208-18.
- 9 Gorard DA, Libby GW, Farthing MJG. Ambulatory small intestinal motility in "diarrhoea" predominant irritable bowel syndrome. *Gut* 1994;35:203-10.
- 10 Whitehead WE, Engel BT, Schuster MM. Irritable bowel syndrome. Physiological and psychological differences between diarrhea-predominant and constipation-predominant patients. *Dig Dis Sci* 1980;25:404-13.
- 11 Prior A, Maxton DG, Whorwell PJ. Anorectal manometry in irritable bowel syndrome: differences between diarrhoea and constipation predominant subjects. *Gut* 1990;31:458-62.
- 12 Kellow JE, Eckersley GM, Jones MP. Enhanced perception of physiological intestinal motility in the irritable bowel syndrome. *Gastroenterology* 1991; 101:1621-7.
- 13 Nanda R, James R, Smith H, Dudley CRK, Jewel DP. Food intolerance and the irritable bowel syndrome. *Gut* 1989;30:1099-104.
- 14 Jones VA, McLaughlan P, Shorthouse M, Workman E, Hunter JO. Food intolerance: a major factor in the pathogenesis of irritable bowel syndrome. *Lancet* 1982;ii:1115-7.
- 15 Harvey RF, Maudud EC, Brown AM. Prognosis in the irritable bowel syndrome: a 5 year prospective study. *Lancet* 1987;ii:963-5.
- 16 Sandragesaran K, Jones BJM. Chronic diarrhoea, bile salt malabsorption, and previous enteric infection. *Gut* 1991;32(suppl):S14.

Psychological treatment is essential for some

Francis Creed

Strong evidence exists that the irritable bowel syndrome has an important psychological component. Firstly, about half of patients with the irritable bowel syndrome in a hospital clinic have psychiatric disorder when assessed by research criteria.¹ This is two or three times greater than the prevalence among patients with organic gastrointestinal conditions such as peptic ulcers or inflammatory bowel disturbance and healthy controls (<20%),¹ suggesting a strong association between the irritable bowel syndrome and psychological disorder. Although anxiety, depression, and sexual problems might be the consequence of persistent abdominal symptoms, they are not found to the same extent in patients with organic diseases that also lead to persistent abdominal symptoms.² The anxiety and depression might explain attendance at the hospital rather than the occurrence of the symptoms,^{3,4} but people attending hospital also have much worse intestinal symptoms than people with the irritable bowel syndrome in the community, and the anxiety almost certainly exacerbates the pain and bowel dysfunction.⁵

Role of stress

Secondly, two thirds of patients with the irritable bowel syndrome have experienced a severe social stress such as bereavement, marital separation, or major argument leading to a broken family relationship before the onset of the abdominal symptoms. This compares with about a quarter of patients with organic disease and healthy controls.⁶ The pattern of social stress before the onset of the irritable bowel syndrome is similar to that preceding deliberate self harm, where precipitating stress is well recognised.

Even in patients with no overt anxiety or depression, onset of the irritable bowel syndrome is still preceded by stressful life events. Not all studies have confirmed this result, but these negative studies have measured life events before attending a clinic rather than before the onset of symptoms and have often used inadequate measures of life events.

Success of psychological treatment

Thirdly, patients with the irritable bowel syndrome attending the clinic for the first time have been randomly allocated to group stress management or pharmacological treatment.⁷ Although the short term response was similar, at six month follow up patients randomised to psychological treatment showed a significantly better response, with patients receiving drugs having returned to their initial symptom level. This result strongly suggests that the core problem is a psychological disturbance.

About 15% of patients with the irritable bowel syndrome do not respond to routine treatment with bulking agents, antispasmodics, and support. Most such patients respond to dynamic psychotherapy or hypnotherapy,⁸⁻¹⁰ and the reduction of bowel symptoms closely follows the reduction in psychological symptoms, indicating a clear link between the two.¹⁰

In their original description of the irritable colon, Chaudhury and Truelove observed that patients who had no history of dysentery before the onset of bowel symptoms but who did have evidence of psychological problem had the worst two year outcome.¹¹ Their case examples indicated that such patients might recover when their circumstances changed. A middle aged man

Department of Psychiatry,
Rawnsley Building,
Manchester Royal
Infirmary, Manchester
M13 9WL
Francis Creed, professor of
community psychiatry

developed symptoms during a marital separation; they cleared when he successfully remarried. A teenage girl with lifelong bowel problems improved greatly when she left home and the conflicts with her parents ceased.

Evidence of a motility disorder

The evidence for a motility disorder is less convincing.¹² The normal motility pattern of the gut is extremely variable, and abnormal motility patterns seem poorly related to a patient's experience of symptoms. In addition, the motility patterns of patients with the irritable bowel syndrome cannot satisfactorily be separated from those of psychoneurotic patients presenting to the psychiatrist,¹³ suggesting that psychological disturbance might cause the abnormal bowel motility patterns as well as the bowel symptoms. Although changed visceral sensitivity has been shown in the irritable bowel syndrome, the only treatment that has been shown to change the sensitivity and improve symptoms is hypnotherapy.¹⁴

Clinical implications

In the United States the irritable bowel syndrome is the commonest condition presenting to gastroenterologists. Half of the gastroenterologists working in academic settings use "personal psychosocial support," for at least three quarters of these patients. Yet there still seems to be a reluctance to recognise the importance of the role of psychological factors in the aetiology or exacerbation of the irritable bowel syndrome in Britain except at a few centres.

The reasons for this are currently being examined by a joint working group of the Royal Colleges of Physicians and Psychiatrists. Most gastroenterologists do not have training which enables them to feel confident in handling the emotional side of the patients they treat. In addition, the environment in medical outpatient clinics is oriented toward investigations for organic disease and not conducive to private interviews in which the doctor can explore psychological problems that might cause or exacerbate symptoms. Some gastroenterologists are prejudiced against acknowledging the importance of psychological factors in their patients, presumably as a result of the organic bias of medical training.

Increasing recognition of the importance of psychosocial factors in the aetiology of the irritable bowel syndrome would lead to more satisfactory treatment. Such treatment might follow that described by Rumsey,⁷ in which small groups of patients met for six weekly sessions to understand the nature of the disorder and the role of stress; to learn the techniques of progressive muscle relaxation and problem solving; and to be taught the importance of diet, fitness, and long term strategies. Although pharmacological treatment seems to help in the short term, any patient who has persistent symptoms should be considered for psychological treatment. This seems to produce better long term results,¹⁰⁻¹⁵ presumably because of the

Commentary: try dietary modification first

Reading these two articles it is difficult to believe that the authors are writing about the same condition. It seems clear that the irritable bowel syndrome is just what it says it is: a syndrome rather than a single pathophysiological entity. It would be interesting to know if those people with demonstrable abnormalities of gut motility also have a history of emotional stress or whether there are at least two essentially separate types of patient with the irritable bowel syndrome. Most people with these symptoms are going to present to general practitioners rather than gastroenterology specialists. Advising dietary modification is simple and inexpensive and can usefully be attempted in everybody. Those who do not respond and who seem to have a definite precipitating factor could then be considered for psychotherapeutic intervention.—PETER C RUBIN, *professor of therapeutics, University of Nottingham*

importance of underlying psychological problems in the syndrome.

- 1 Creed F. Relationship of non-organic abdominal pain to psychiatric disorder and life stress. In: Creed F, Mayo R, Hopkins A, eds. *Medical symptoms not explained by organic disease*. London: Royal Colleges of Psychiatrists and Physicians, 1992:9-16.
- 2 Guthrie E, Creed FH, Whorwell P. Severe sexual dysfunction in women with the irritable bowel syndrome: comparison with inflammatory bowel disease and duodenal ulceration. *BMJ* 1987;295:577-8.
- 3 Drossman DA, McKee DE, Sandler RS, Mitchell CM, Cramer EM, Lowman BC, et al. Psychosocial factors in the irritable bowel syndrome. A multivariate study of patients and nonpatients with irritable bowel syndrome. *Gastroenterology* 1988;95:701-8.
- 4 Whitehead WE, Bosmajian L, Zonderman AB, Costa PT Jr, Schuster MM. Symptoms of psychologic distress associated with irritable bowel syndrome. Comparison of community and medical clinical samples. *Gastroenterology* 1988;95:709-14.
- 5 Heaton K. What makes people with abdominal pain consult their doctor? In: Creed F, Mayo R, Hopkins A, eds. *Medical symptoms not explained by organic disease*. London: Royal Colleges of Psychiatrists and Physicians, 1992:1-8.
- 6 Creed F, Craig T, Farmer R. Functional abdominal pain, psychiatric illness and life events. *Gut* 1988;29:235-42.
- 7 Rumsey N. Group stress management programmes versus pharmacological treatment in the treatment of the irritable bowel syndrome. In: Heaton KW, Creed F, Goeting NLM, eds. *Towards confident management of irritable bowel syndrome*. Lyme Regis: Duphar, 1991:33-9.
- 8 Whorwell PJ, Prior A, Faragher EB. Controlled trial of hypnotherapy in the treatment of severe refractory irritable bowel syndrome. *Lancet* 1984;ii:1232-4.
- 9 Svedlund J. Psychotherapy in irritable bowel syndrome: a controlled outcome study. *Acta Psychiatr Scand* 1983;67(suppl 306):1-86.
- 10 Guthrie E, Creed F, Dawson D, Tomenson B. A controlled trial of psychological treatment for the irritable bowel syndrome. *Gastroenterology* 1991;100:450-7.
- 11 Chaudhury NA, Truelove SC. The irritable colon syndrome: a study of the clinical features, predisposing causes, and prognosis in 130 cases. *Q J Med* 1962;31:307-23.
- 12 Gorard SA, Libby GW, Farthing MJG. Ambulatory small intestinal motility in "diarrhoea" predominant irritable bowel syndrome. *Gut* 1994;35:203-10.
- 13 Latimer P, Sarna S, Campbell D, Latimer M, Waterfall W, Daniel EE. Colonic motor and myoelectrical activity: a comparative study of normal subjects, psychoneurotic patients, and patients with irritable bowel syndrome. *Gastroenterology* 1981;80:893-901.
- 14 Prior A, Colgan SM, Whorwell PJ. Changes in rectal sensitivity after hypnotherapy in patients with irritable bowel syndrome. *Gut* 1990;31:896-8.
- 15 Drossman DA, Thompson WG. The irritable bowel syndrome: review and graduated multicomponent treatment approach. *Ann Intern Med* 1992;116:1009-16.