These are not sufficient arguments for an editorial in the *BMJ* to recommend one class of β blockade and should be accompanied by the proviso that measurements of blood pressure are only a surrogate: it is the prevention of cerebrovascular accidents and myocardial infarctions that matters.

As a doctor treating patients with hypertension or ischaemic heart disease most days of the week, I tend to use low doses of the most β_1 selective lipophilic agent, bisoprolol, for hypertension and the highest affinity non-selective agent, timolol, for ischaemic heart disease. These drugs have been selected on the basis of theoretical arguments and surrogate measures on which Cleophas and Kalmansohn and I agree. As a clinical pharmacologist unhappy at such evidence, I set up the East Anglian comparison of the two classes of drug in ischaemic heart disease that I mentioned in my editorial and included them in a long term randomised comparison of antihypertensive drugs that started in 1986.

MORRIS J BROWN
Professor

Clinical Pharmacology Unit,
University of Cambridge Clinical School,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

Withdrawing artificial feeding from children with brain damage

EDITOR,—Ronald E Cranford's editorial on withdrawing artificial feeding from children with brain damage aims to encourage debate about basic medical and ethical principles.¹ Yet I believe that the central medical and ethical issues are not fairly addressed.

Firstly, Cranford states that so called artificial feeding is medical treatment but implies that normal eating and drinking are not. Surely this is specious. Tube feeding, whether by nasogastric tube or gastrostomy, is largely done to protect the patient's airway from soiling and for nursing expediency. Moreover, "it serves the same purpose that eating and drinking do, [to provide] the food any human being needs."² Tube feeding is better considered to be part of tender, loving care rather than to have any therapeutic benefit itself. I am sure that doctors would never want tender, loving care to be removed from a patient. This point is supported by my experience that nursing staff are generally extremely reluctant to countenance withdrawal of feeding because they have an instinctive belief that it is part of nursing care and explicitly not a medical treatment.

Secondly, Cranford concludes that allowing a child to die is not the same as euthanasia (or, presumably, infanticide). But in what way is it different? Both produce certain death. This sets the withdrawal of feeding apart from the withdrawal of other medical treatments, which merely increases the likelihood of death but does not guarantee it. The difference here is between, say, 99% and 100%—infinity. A spade should be called a spade. The proponents of withdrawal of feeding are, in truth, asking for permission to practise infanticide through privation. The next logical step is to give drugs to hasten the process.³ The Rubicon has been crossed.

Children with severe brain damage present immense challenges to everyone. If our only solution to the problem is deliberate killing what does that say about us? Can we see no intrinsic value in a person created in the image of God, or do we value only possession of abilities? Dyck at Harvard School of Public Health asks "whether we are the kind of persons who will care for [such children] without doubting their worth."⁴ He believes in the equality of life rather than the quality of life ethic. Koop, a former surgeon general of the Public Health Service in the United States, agrees with him.⁵ So do I. So did Hippocrates.⁶

Ethical principles exist to guide us with difficult issues despite the ever changing technology of medicine. We must recognise the difference between good medical practice and killing people.

P E SHANNON
Research registrar in anaesthetics

Royal Hallamshire Hospital,
Sheffield S10 2JR

- 1 Cranford RE. Withdrawing artificial feeding from children with brain damage. *BMJ* 1995;311:464-5. (19 August.)
- 2 Gormally L. Definitions of personhood: implications for the care of PVS patients. *Ethics and Medicine* 1993;3:44-8.
- 3 Koop CE. Life and death and the handicapped newborn. *Ethics and Medicine* 1987;3:39-44.

Informed consent for trial of elective ventilation will not be forthcoming

EDITOR,—J Fabre seeks the continuation and legalisation of the unlawful procedure of elective ventilation on the grounds that it is not harmful to patients.¹ Later Fabre contradicts this by admitting that elective ventilation might result in a persistent vegetative state and adds that the British Transplantation Society advocates "a carefully audited clinical trial." Surely such a trial, even if it was considered to be ethical, would require the explicitly informed prior consent of the patients concerned; it is difficult to see how this could be obtained.

A persistent vegetative state is not the only possible harm to a patient, as elective ventilation must prolong the process of dying. Once death is inevitable many people would prefer it to occur as quickly and naturally as possible and would wish to be able to trust their medical attendants to allow this. A badly damaged but artificially oxygenated brain (not necessarily brain stem dead) might also produce unpleasant and distressing subjective sensations, which the patient would not be able to communicate to the attendants.

Though I appreciate the great boon that successful transplantation confers on recipients and the natural desire of the British Transplantation Society to maximise this, we should never forget the legitimate interests of our many other patients.

EVA V TREGILLUS
Retired general practitioner

Rockmount,
Reeth, North Yorkshire

- 1 Fabre J. Elective ventilation of potential organ donors. *BMJ* 1995;311:950. (7 October.)

Misuse of dothiepin

EDITOR,—Recently we have become aware of a considerable population of intravenous drug misusers in the Greater Dublin area who are misusing the tricyclic antidepressant dothiepin (Prothiaden). The Drug Treatment Centre, Dublin, runs an evening programme of low dose methadone maintenance and harm reduction with needle exchange and health promotion advice but without urine testing. A self report questionnaire administered at this programme showed that 46% of clients (38/83) had misused this drug orally in the previous six months. Of all urine samples tested on 14 March of clients attending the centre by day who were not taking prescribed tricyclic or neuroleptic agents, 19% (19/99) were positive for tricyclics. The Emit tox test (Syva, San Jose, CA) was originally designed for serum samples, and the presence of phenothiazines is a known cause of false positive results. We conclude that this percentage is an indication of misuse of tricyclic drugs in those tested.

Patients who misuse dothiepin report euphoria and sedation with complex visual and auditory

hallucinations. The experiences are pleasant and seem to occur in clouded consciousness. The drug was taken orally in all cases, and the total amount taken varied from 150 mg to 600 mg each day.

We know of no previous report of dothiepin misuse, although there have been reports of abuse of another tricyclic, amitriptyline.^{1,2} Dothiepin is the leading antidepressant in Ireland, with over 140 000 prescriptions each year, representing 24% of total prescriptions for antidepressants. A recent review concluded that the major tranquillisers and tricyclic antidepressants had no dependence liability and no abuse potential of any clinical importance.³ Given our experience we must disagree. Concerns about the abuse potential of any centrally active or mood altering drug are not new,⁴ and clinicians should be aware of the potential of dothiepin and other similar compounds for misuse.

ARTHUR DORMAN
Senior registrar

PETER BYRNE
Registrar

The Drug Treatment Centre Board,
Dublin 2,
Republic of Ireland

DYMPNA TALBOT
Principal biochemist

JOHN O'CONNOR
Clinical director

- 1 Cohen MJ, Hanbury R, Stimmel B. Abuse of amitriptyline. *JAMA* 1978;240:1372-3.
- 2 Cantor R. Methadone maintenance and amitriptyline. *JAMA* 1979;241:2378.
- 3 Lichtigfeld FG, Gillman MA. The possible abuse of and dependence on major tranquillisers and tricyclic antidepressants. *SA Med J* 1994;84:5-6.
- 4 Sheppard M. The use and abuses of drugs in psychiatry. *Lancet* 1970;ii:31-3.

Palliative care in general practice

Should GPs do it at the expense of commoner problems?

EDITOR,—As a former general practitioner who is currently undergoing higher specialist training in palliative medicine, I am concerned at Zoe Kenyon's vision of a future in which a diploma in palliative medicine is a criterion for entry to general practice.¹ My concern arises not least because to get the diploma one must attend a part time course lasting at least a year, which would distract focus away from vocational training.

The editorial raises a larger question: is palliative care delivered by general practitioners for all patients a realistic option? The general practice palliative care facilitator project described in Kenyon's editorial was designed to enhance the quality of palliative care delivered by primary care teams. As the evaluation makes clear, the educational activities directed towards generic skills such as teamwork and communication were well received.² There was a poor response, however, to initiatives focused on more specific aspects of palliative medicine, such as control of symptoms, as these subjects are not considered "mainstream" by general practitioners. As the report states, "There is little evidence to suggest that the visits [of facilitators to practices] were able to . . . influence the 'peripheral' nature of palliative care as part of the workload of the primary care team."

While many aspects of good palliative care—multidisciplinary, holistic care, centred on the patient and family, that values good communication—are directly comparable with good general practice, the average general practitioner will see only about five patients dying of cancer each year and probably fewer with other end stage diagnoses but similar needs. Other problems seen more frequently, such as asthma, are bound to take priority. Criteria for entry to general practice would be better focused on the attainment of skills in good general practice, including communication skills, team working, holistic care, and a basic knowledge of first line control of symptoms.

One of the most important attributes of a good general practitioner is the wisdom to know when to refer to specialist services. This also depends, however, on such services being accessible to all in need.

DAVID BROOKS
CRC lecturer in palliative medicine

Department of Surgical and Anaesthetic Sciences,
Palliative Medicine Section,
University of Sheffield,
Royal Hallamshire Hospital,
Sheffield S10 2JF

1 Kenyon Z. Palliative care in general practice. *BMJ* 1995;311: 888-9. (7 October.)

2 Royal College of General Practitioners. *General practice palliative care facilitator project. Report of an evaluation.* London: RCGP, 1995.

Palliative care is integral to practice

EDITOR,—Zoe Kenyon's editorial on palliative care in general practice makes wide generalisations about the skills in and enthusiasm for palliative care of "many general practitioners" in Britain.¹ Few general practitioners shy away from palliative care, which, after all, is a large and satisfying part of their practice. Nor do general practitioners limit their definition of palliative care and its application to patients with cancer, as many hospitals do. On the contrary, most general practitioners expose themselves daily, if not hourly, to "painful emotions." Granted, there are exceptions, but the conclusions reported must be based on systematic research rather than potentially biased observations.

Fortunately, misconceptions about ceiling doses of opiates are disappearing and modern day general practice is not preoccupied with the science of controlling symptoms. Rather, it has advanced in both its teaching and practice to reduce suffering; pain control is only one element in this multi-dimensional approach. In other words, provision is made to prepare the "whole person" for a good death, which is probably preferable to a specialist pharmacologically controlled "biological death."

Both generalists and specialists have a role in palliative care, and they should work together, aware of each other's skills and emphasising the long held vision that "It is better to help a colleague with a difficult case than to tell him he is wrong and that he should make way for the expert."²

Rather than palliative care joining the ever growing list of subjects in which one can obtain a diploma, it should become an underpinning element of vocational training in general practice. This is particularly so with the shift in emphasis to community care, in which the general practitioner acts as the focal point for the primary health care team.

RODGER CHARLTON
Senior lecturer

Centre for Primary Health Care,
Keele University School of Postgraduate Medicine,
North Staffordshire Hospital,
Stoke on Trent ST4 7QB

1 Kenyon Z. Palliative care in general practice. *BMJ* 1995;311: 888-9. (7 October.)

2 Pugsley R, Pardoe J. The specialist contribution to the care of the terminally ill patient: support or substitution? *J R Coll Gen Pract* 1986;36:347-8.

Significant social class gradient in menstrual disorders

EDITOR,—Diana Kuh and Susan Stirling report significant educational gradients for risk of admission for diseases of the female genital system and breast, and in particular for menstrual disorders: 1% of highly educated women were admitted to hospital for menstrual problems compared with 19% of those with minimal qualifications.¹ Social gradients were also observed for hysterectomies

and dilatation and curettage: 37% and 44% of these procedures, respectively, were related to menstrual problems. The authors refer to a slight increase in rates of hysterectomy since the 1970s. They were unable, however, to establish whether increased risks of admission and surgery among less educated women and women of lower social class reflected the prevalence of disease or a different pattern of primary and secondary care.

Evidence for social differences in menstrual disorders, not affected by use of health services, is available from the 1958 British birth cohort study (table). At age 33, women reported whether they had ever had a menstrual problem. Eighteen per cent reported that they had, of whom 14% had had such a problem during the past year. There were strong associations between social position and menstrual problems.

Social class distribution of menstrual problems among women in 1958 British birth cohort study

	At birth		At age 33	
	% (No)	Odds ratio	% (No)	Odds ratio
Social class:				
I	10.5 (24)	1.00	10.9 (15)	1.00
II	13.1 (92)	1.28	16.8 (265)	1.66
IIINM	16.1 (84)	1.64*	16.5 (318)	1.63
IIIM	19.0 (508)	2.01*	16.3 (63)	1.59
IV	22.1 (144)	2.43*	21.8 (208)	2.28*
V	18.5 (83)	1.94*	23.0 (67)	2.45*
χ^2 For trend	23.2		17.9	
P value	<0.0001		<0.0001	
Educational qualifications:				
Degree			14.4 (211)	1.00
A level			18.1 (106)	1.32*
O level			17.6 (362)	1.27*
< O level			18.7 (179)	1.37*
None			26.4 (161)	2.13*
χ^2 For trend			32.5	
P value			<0.0001	

*P<0.05.

Evidence suggests an association between menstrual problems and obesity, with overweight and obese women having a higher risk of menstrual disorders (JL, CP, TC, unpublished observations).³ Women in lower social groups are also at a greater risk of obesity than those of higher socioeconomic status.² It is therefore important to consider whether the social distribution of obesity could account for the distribution of menstrual disorders highlighted in Kuh and Stirling's paper. The recent increase in obesity² may also be partly responsible for the increase in menstrual problems (and possibly hysterectomies and dilatation and curettage). If this hypothesis is confirmed then a reduction in the socioeconomic gradient in admissions for menstrual disorders (and possibly hysterectomies and dilatation and curettage) may be achieved by reducing the social gradient of obesity.

JULIE LAKE
Research assistant
Department of Epidemiology and Biostatistics,
Institute of Child Health,
London WC1N 1EH

CHRIS POWER
Senior lecturer

MRC Dunn Nutrition Unit,
Cambridge CB4 1XJ

TIM COLE
Statistician

1 Kuh D, Stirling S. Socioeconomic variation in admission for diseases of female genital system and breast in a national cohort aged 15-43. *BMJ* 1995;311:840-3. (30 September.)

2 Bennett N, Dodd T, Flatley J, Freeth S, Bolling K. *The health of our nation: health survey for England 1993.* London: Office of Population Censuses and Surveys, 1995.

3 Hartz AJ, Barboriak PN, Wong A, Katayama KP, Rimm AA. The association of obesity with infertility and related menstrual abnormalities in women. *International Journal of Obesity* 1979; 3:57-63.

Reflex sympathetic dystrophy in children

Dystrophic changes are less likely

EDITOR,—Having read Adrian R Lloyd Thomas and Gillian Lauder's article on reflex sympathetic dystrophy in childhood,¹ we would like to raise some points that reflect our experience of this problem. Over two years we have seen 11 children (10 girls and one boy) under the age of 15 with reflex sympathetic dystrophy.

The authors state that most patients with childhood reflex sympathetic dystrophy do not have a history of trauma. Only three of our patients could not recall a traumatic precipitant for their symptoms, although in three cases there was an asymptomatic period (of up to four months) between the traumatic episode that may have been responsible and the onset of symptoms. In the remaining five cases there was continuity between the symptoms directly attributable to trauma and the onset of the reflex sympathetic dystrophy. Whereas dystrophic changes are seen in the adult form of the condition, our experience suggests that these are relatively rare in children (only two patients had changes that could be construed as dystrophic (quadriceps and calf muscle wasting)).

The authors suggest the use of radiography to help diagnose this condition. Although osteoporosis is a well recognised late finding in the adult form of the disease, we have found it less common in children. Only four of our patients had possible radiological evidence of osteoporosis, and in all four it was mild. Additionally, osteoporosis cannot be reliably diagnosed in plain radiographs in the absence of fractures. Most reports suggest that the prognosis for childhood reflex sympathetic dystrophy is good.^{2,4} Our findings support this, although in one of our patients the condition had a protracted course, symptoms still being present after two years.

R KAVANAGH
Rheumatology research registrar
A J CRISP
Consultant rheumatologist
B L HAZLEMAN
Consultant rheumatologist

Rheumatology Research Unit,
PO Box 194,
Addenbrooke's Hospital,
Cambridge CB1 4TH

R J COUGHLAN
Consultant rheumatologist

Department of Rheumatology,
University College Hospital,
Galway,
Republic of Ireland

1 Lloyd-Thomas AR, Lauder G. Reflex sympathetic dystrophy in children. *BMJ* 1995;310:1648-9. (24 June.)

2 Bernstein BH, Singen BH, Kent JT, Kornreich H, King K, Hicks R, et al. Reflex neurovascular dystrophy in childhood. *J Pediatr* 1978;93:211-5.

3 Ruggeri SB, Athreya BH, Dougherty R, Greig JR, Das MM. Reflex sympathetic dystrophy in children. *Clin Orthop* 1982;163:225-30.

4 Kesler RW, Saulsbury FT, Miller LT, Rowlingson JC. Reflex sympathetic dystrophy in children: treatment with transcutaneous electric nerve stimulation. *Pediatrics* 1988;82: 728-32.

Active early physiotherapy is the key to prevention

EDITOR,—The article on reflex sympathetic dystrophy in children is pertinent to the practice of paediatric accident and emergency care.¹ While the cases that the authors describe are extreme, they are by no means unusual. The authors, however, concentrate more on treatment than on prevention, and they state that the average time from onset of symptoms to presentation is 10 days.

Those of us who work in accident and emergency care, and especially those of us who see large numbers of children, are familiar with peri-adolescent girls presenting to the accident and