Published in final edited form as:

Org Lett. 2006 March 30; 8(7): 1283–1286. doi:10.1021/ol052948+.

Synthesis of (+)-Sch 642305 by a Biomimetic Transannular Michael Reaction

Barry B. Snider* and Jingye Zhou

Dept of Chemistry MS 015, Brandeis University, Waltham Massachusetts 02454-9110

Abstract

The synthesis of (+)-Sch 642305 (**1**) has been completed in 17 steps in 1.6% overall yield. Transannular Michael reaction of **2b** with NaH in THF provided cyclohexenone **23** stereospecifically. Heating **23** in TFA/CDCl₃ provided a 3:1 equilibrium mixture of **23** and **25**, which were hydrolyzed to give (+)-6-*epi*-Sch 642305 (**24**) and (+)-Sch 642305 (**1**), respectively.

In late 2003, Schering-Plough scientists reported the isolation and structure elucidation of the bicyclic macrolide Sch 642305 (**1**) from *Penicillium verrucosum* (culture ILF-16214) isolated from soil collected near Tucson, AZ.¹ The structure was determined spectroscopically and the absolute stereochemistry was assigned by X-ray crystallographic analysis of the *p*-bromobenzoate ester. Macrolide **1** inhibits bacterial DNA primase with an EC₅₀ of 70 μM.¹ In 2005, Merck scientists reported the isolation of **1** from the fungus *Septofusidium* sp. (JP3241) isolated from leaf litter collected in Puerto Rico.² Macrolide **1** inhibits HIV-1 Tat, a regulatory protein required for viral replication, with an IC₅₀ value of 1 μM.² Mehta and Shinde recently reported the syntheses of (+)-Sch 642305 (**1**)³ and (+)-11-*epi*-Sch 642305⁴ using a ring closing metathesis to elaborate a 10-membered ring lactone onto a highly functionalized cyclohexene.⁵

We thought that Sch 642305 (**1**) might be biosynthesized by a transannular Michael reaction of the stereochemically simpler hydroxyenone **2a** (see Scheme 1). The close relationship of the macrolides nigrosporolide (**7**)⁶ and mutolide (**8**)⁷ to the hypothetical biosynthetic intermediate **2a** supports this hypothesis (see Figure 1). If **1** is biosynthesized this way, it might also be possible to synthesize it selectively by cyclization of **2a** or a protected derivative such as **2b**. The cyclization of **2** to **1** introduces two new stereocenters so that four isomers are possible. However, related macrolides are often fairly rigid with their conformation controlling the stereochemistry of reactions.⁸ Although there are a few reports of synthetic applications of transannular Michael⁹ and aldol¹⁰ reactions, the potential of these reactions have not been fully developed.

snider@brandeis.edu.

Supporting Information Available. Full experimental details and copies of ¹H and ¹³C NMR

Macrolide **2** should be readily available by macrolactonization of hydroxyacid **3**, followed by hydrolysis of the dioxolane. Hydroxyacid **3** can be prepared by Lindlar reduction of the alkyne of hydroxy ester **4** followed by protecting group modification. Addition of the acetylide anion formed by deprotonation of **6** to commercially available aldehyde **5** will provide **4** with the complete carbon skeleton of **1**.

Addition of $\text{LiC}\equiv\text{CTMS}$ to 7-octenal (**9**)¹¹ in THF at $-40\text{ }^{\circ}\text{C}$ for 2 h afforded the propargylic alcohol, which was oxidized with PCC in CH_2Cl_2 for 12 h to give ynone **10** in 83% overall yield (see Scheme 2). Formation of the dioxolane with $\text{HOCH}_2\text{CH}_2\text{OH}$, $\text{HC}(\text{OMe})_3$, and TsOH in benzene at reflux for 8 h, followed by cleavage of the TMS group with K_2CO_3 in MeOH for 12 h at $25\text{ }^{\circ}\text{C}$ afforded **11** in 89% yield. Epoxidation of the terminal double bond of **11** proceeded cleanly with 2 equiv of mCPBA in CH_2Cl_2 for 12 h. 2-Methyl-2-butene was added and reaction was continued for 4 h to consume excess mCPBA, giving racemic **12** in 99% yield. Jacobsen kinetic resolution¹² using 0.01 mol% of the oligomeric (salen)Co(III) catalyst¹³ prepared from the (*R,R*)-diamine and 0.5 equiv of H_2O in MeCN for 48 h afforded 46% of (*R*)-**12** in 92.5% ee as determined by chiral HPLC analysis of the 2-mercaptobenzothiazole derivative.^{12b} Reduction of the epoxide with NaBH_4 in EtOH at reflux for 8 h afforded (*S*)-**13** in 99% yield.¹⁴

Initially we had carried out the kinetic resolution of **12** with 1 mol% of the monomeric (salen)Co(II) catalyst and 4 mol% of HOAc, which gave **12** in 43% yield and 97% ee. Reduction of this material with NaBH_4 in EtOH for 4 h at $50\text{ }^{\circ}\text{C}$ afforded a 3:1 mixture of **13** and the corresponding alkenol resulting from partial reduction of the triple bond. This byproduct was not formed during reduction of racemic **12**. We speculated that cobalt impurities present even in chromatographically purified kinetically resolved **12** reacted with NaBH_4 to form a catalyst that reduced the triple bond.¹⁵ This was confirmed by reaction of 1-pentadecyne with 1 mol % monomeric (salen)Co(II) catalyst and excess NaBH_4 in EtOH at $55\text{ }^{\circ}\text{C}$ to afford 15% of 1-pentadecene. Under these conditions, 1-octyn-3-ol was reduced more rapidly to give mainly 1-octen-3-ol. This side reaction is minimized with the oligomeric (salen)Co(III) catalyst, which can be used at only 0.01 mol% loading.

The alcohol of **13** was protected as the TBS ether with TBSCl and imidazole in DMF for 12 h at $25\text{ }^{\circ}\text{C}$ to give **6** in 92% yield. Deprotonation of **6** with *n*-BuLi in THF at $-40\text{ }^{\circ}\text{C}$ for 12 h followed by addition of aldehyde **5** and stirring for 30 min at $-40\text{ }^{\circ}\text{C}$ gave 74% of **14** as an inseparable 1:1 mixture of diastereomers. Numerous asymmetric additions of acetylide anions to aldehydes have been reported recently including those of Carriera,¹⁶ Pu,¹⁷ and Shibasaki.¹⁸ In our hands these procedures worked well on the reported substrates, but not with alkyne **6** and aldehyde **5**. We also oxidized **14** to give an enynone and investigated CBS-catalyzed asymmetric reduction,¹⁹ which has been used extensively on enynones. However, the reduction proceeded in only 30–60% yield with up to 70% ee. We therefore continued with **14** as a diastereomeric mixture of isomers, with the expectation that they would be readily separated after formation of the macrolide. The rationale for our approach to **1** was that the conformation and reactivity of macrolide **2** would be controlled by the stereochemistry of the substituents. In a sense, the mixture of diastereomers of **14** is advantageous because it should allow us to easily investigate the transannular Michael reaction of both **2** and its diastereomer.

Hydrogenation of the triple bond of **14** over 5% Pd/poisoned with lead (1 mol%) in EtOAc containing 0.5 equiv of quinoline under 1 atm H_2 for 6 h gave alkene **15** in 91% yield (see Scheme 3). The seven adjacent functionalized carbons in **15** made progress toward the macrolide challenging. For instance, treatment with TBAF in THF isomerized the allylic alcohol to give ketone **16**. Cleavage of the TBS ether with PPTS in EtOH at $50\text{ }^{\circ}\text{C}$ also cleaved the dioxolane giving a γ -hydroxy-*cis*-enone that cyclized and lost water to give furan **17**. Hydrolysis of the TBS ether with pyr/HF gave a diol, which underwent basic hydrolysis with

LiOH to give the desired dihydroxy acid. However, attempted macrolactonization under Yamaguchi conditions with $\text{C}_6\text{H}_2\text{Cl}_3\text{COCl}$, Et_3N , and DMAP failed; the conjugated doubly allylic alcohol is not stable under the reaction conditions. We hoped that protecting the doubly allylic alcohol with a bulky TBDPS ether would allow the macrolactonization to proceed successfully.

Reaction of **15** with excess TBDPSCl and imidazole in DMF for 12 h formed the TBDPS ether. MeOH was added and the solution was stirred for 4 h resulting in cleavage of the TBS ether to give **18** in 73% yield (see Scheme 4). Either imidazolium chloride or the HCl generated by reaction of excess TBDPSCl with MeOH catalyzed the hydrolysis of the TBS ether without cleaving the dioxolane. Hydrolysis of the ethyl ester with excess LiOH in 2:1:1 THF/MeOH/ H_2O for 20 h at 25 °C provided carboxylic acid **19** in 90% yield. Yamaguchi lactonization was effected by slow addition of the mixed anhydride (formed with $\text{C}_6\text{H}_2\text{Cl}_3\text{COCl}$ and Et_3N) to DMAP in toluene at 25 °C to give **20** as a difficultly separable mixture of diastereomers in 66% yield. Heating **20** with catalytic TsOH in 10:1 acetone/ H_2O at 50 °C for 16 h cleaved the dioxolane to give a readily separable mixture of the desired keto lactone **2b** (44%, 88% from the desired diastereomer of **20**), the diastereomeric ketone lactone **21** (6%, 12% from the undesired diastereomer of **20**) and the isomerized lactone **22** (18%). The stereochemistry of **2b** was assigned by its eventual conversion to **1**. The stereochemistry of the enol ether double bond of **22** was established by an NOE between the alkene and phenyl protons.

Pure macrolide **21** is readily isomerized to **22** by TsOH in acetone/ H_2O at 50 °C, while the desired isomer **2b** is stable. The stereochemistry of the OTBDPS group has a profound effect on the stability of **2b** and **21**. Enol formation from **21** is more rapid, suggesting that conformers in which the doubly allylic hydrogen is properly aligned with the enone double bond are more accessible for **21** than for **2b**.

Treatment of **2b** with 1.2 equiv of NaH in THF at 0 °C for 30 min afforded the transannular Michael adduct **23** as a single isomer in 80% yield (See Scheme 5). Deprotection of **23** with 1:1 TBAF/ HOAc^3 in THF for 3 h at 25 °C provided (+)-6-*epi*-Sch 642305 (**24**) in 90% yield. The *cis* stereochemistry of **24** was tentatively assigned based on the 3.2 Hz coupling constant between H_5 and H_6 . An NOE between H_4 and H_6 completed the assignment of the cyclohexene stereochemistry. An NOE between H_{11} and the H_7 that is anti-periplanar to H_6 suggested that the relative stereochemistry at C_4 , C_5 , and C_{11} is the same as in Sch 642305 (**1**). Cyclization of **2b** under a variety of other basic conditions gave predominantly **23** in lower yields. Attempted formation of the silyl enol ether from **2b** led mainly to **22**. Hydrolysis of **2b** with HF/pyr cleaved the silyl ether to give alcohol **2a**, which gave complex mixtures on treatment with base.

6-*epi*-Sch 642305 (**24**) differs from **1** in the stereochemistry at C_6 , which is adjacent to a ketone and therefore epimerizable. We were hopeful because MM2 calculations suggested that **1** is about 1 kcal/mol more stable than **24**. Treatment of **24** under basic conditions resulted in no reaction or decomposition. Treatment of **24** with 3% TFA in CDCl_3 for 1 d at 55 °C effected dehydration to give only phenol **26** in 65% yield. Treatment of **24** under less acidic conditions with 0.4% TFA in CDCl_3 for 14 d at 55 °C gave an inseparable 1.5:1:2 mixture of **24**, **1**, and **26**, respectively. Formation of phenol **26** was minimized by carrying out the isomerization on TBDPS ether **23**. The best results were obtained by microwave irradiation of **23** in 1.5% TFA in CDCl_3 for 3 h at 120 °C, which gave a readily separable 1:3 mixture of **25** (23% isolated, 58% based on recovered **23**) and **23** (60% isolated) containing only a trace of phenol **26**. Similar mixtures are obtained starting with **25** indicating that this is an equilibrium mixture.²⁰ The spectral data of **25** are identical to those reported by Mehta.³ Hydrolysis of the TBDPS ether of **25** with TBAF/ HOAc^3 in THF afforded (+)-Sch 642305 (**1**) in 86% yield with ^1H and ^{13}C NMR spectral data identical to those previously reported.^{1,3}

The carbons at δ 18.5, 22.6, 23.0, and 23.2 are broadened in the ^{13}C NMR spectrum of **24** and one absorbs as a poorly defined broad peak from δ 29.6–30.8 suggesting that a conformational equilibrium is slow on the NMR time scale. This was confirmed by a ^{13}C NMR spectrum of **24** at 50 °C that showed sharp peaks. This broadening is similar, but less pronounced, in the ^{13}C NMR spectrum of **1** at 25 °C.

In conclusion, we have completed the synthesis of (+)-Sch 642305 (**1**) in 17 steps in 1.6% overall yield. Transannular Michael reaction of **2b** with NaH in THF afforded cyclohexenone **23** stereospecifically. TFA catalyzed equilibration provided a 3:1 mixture of **23** and **25**, which were hydrolyzed to give (+)-6-*epi*-Sch 642305 (**24**) and (+)-Sch 642305 (**1**), respectively.

Supplemental Materials

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank the NIH (GM50151) for generous financial support. We thank Prof. Eric Jacobsen for a sample of the oligomeric (salen)Co(III) catalyst.

References

1. Chu M, Mierzwa R, Xu L, He L, Terracciano J, Patel M, Gullo V, Black T, Zhao W, Chan TM, McPhail AT. *J Nat Prod* 2003;66:1527–1530. [PubMed: 14695789]
2. Jayasuriya H, Zink DL, Polishook JD, Bills GF, Dombrowski AW, Genilloud O, Pelaez FF, Herranz L, Quamina D, Lingham RB, Danzeisen R, Graham PL, Tomassini JE, Singh SB. *Chem Biodiversity* 2005;2:112–122.
3. Mehta G, Shinde HM. *Chem Commun* 2005:3703–3705.
4. Mehta G, Shinde HM. *Tetrahedron Lett* 2005;46:6633–6636.
5. The numbering scheme used in reference 1 is retained for clarity.
6. Harwood JS, Cutler HG, Jacyno JM. *Nat Prod Lett* 1995;6:181–185.
7. Bode HB, Walker M, Zeeck A. *Eur J Org Chem* 2000:1451–1456.
8. Kaisalo L, Hase T. *Synthesis* 2001:1619–1622.
9. (a) Shimizu I, Nakagawa H. *Tetrahedron Lett* 1992;33:4957–4958. (b) Matsuura T, Yamamura S. *Tetrahedron Lett* 2000;41:4805–4809.
10. Karim MR, Sampson P. *Tetrahedron Lett* 1988;29:6897–6900.
11. Prepared in 84% yield by PCC oxidation of the commercially available alcohol.
12. (a) Tokunaga M, Larrow JF, Kakiuchi F, Jacobsen EN. *Science* 1997;277:936–938. [PubMed: 9252321] (b) Schaus SE, Brandes BD, Larrow JF, Tokunaga M, Hansen KB, Gould AE, Furrow ME, Jacobsen EN. *J Am Chem Soc* 2002;124:1307–1315. [PubMed: 11841300] (c) Ready JM, Jacobsen EN. *Angew Chem Int Ed* 2002;41:1374–1377. (d) Nielsen LPC, Stevenson CP, Blackmond DG, Jacobsen EN. *J Am Chem Soc* 2004;126:1360–1362. [PubMed: 14759192]
13. White DE, Jacobsen EN. *Tetrahedron: Asymmetry* 2003;14:3633–3638. Catalyst 2a was used
14. For the preparation of analogous alcohols see: (a) Chow S, Kitching W. *Tetrahedron: Asymmetry* 2002;13:779–793. (b) Fürstner A, Thiel OR, Kindler N, Bartkowska B. *J Org Chem* 2000;65:7990–7995. [PubMed: 11073608] Lower yields of **13** were obtained with LiBEt_3H
15. (a) Chung SK. *J Org Chem* 1979;44:1014–1016. (b) Heinzman SW, Ganem B. *J Am Chem Soc* 1982;104:6801–6802. (c) Satyanarayana N, Periasamy M. *Tetrahedron Lett* 1984;25:2501–2504.
16. (a) Frantz DE, Fässler R, Carreira EM. *J Am Chem Soc* 2000;122:1806–1807. (b) Boyall D, López F, Sasaki H, Frantz D, Carreira EM. *Org Lett* 2000;2:4233–4236. [PubMed: 11150207]
17. Pu L. *Tetrahedron* 2003;59:9873–9886.
18. Takita R, Yakura K, Ohshima T, Shibasaki M. *J Am Chem Soc* 2005;127:13760–13761. [PubMed: 16201775]

19. (a) Corey EJ, Helal CJ. *Angew Chem Int Ed* 1998;37:1986–2012. (b) McDonald FE, Reddy KS, Díaz Y. *J Am Chem Soc* 2000;122:4304–4309. (c) Garcia J, López M, Romeu J. *Synlett* 1999:429–431. (d) Yun H, Danishefsky SJ. *J Org Chem* 2003;68:4519–4522. [PubMed: 12762760] (e) Parker KA, Katsoulis IA. *Org Lett* 2004;6:1413–1416. [PubMed: 15101755] (f) Garcia J, López M, Romeu J. *Tetrahedron: Asymmetry* 1999;10:2617–2626.
20. Although MM2 calculations suggested that 25 is about 1 kcal/mol more stable than 23, the equilibration studies indicate that it is less stable. This could be due to inaccuracies in the MM2 calculations or entropic effects.

nigrosporolide (**7**)

mutolide (**8**)

Figure 1.
Structures of nigrosporolide (**7**) and mutolide (**8**)

Scheme 1.
Retrosynthesis of Sch 642305 (**1**)

Scheme 2.
 Synthesis of **14**

Scheme 3.

Scheme 4.
Preparation of Keto Lactone **2b**

Scheme 5.
Preparation Sch 642305 (**1**)