right for a patient to be managed in a unit which sees only occasional cases. Would it not be better for the patients to be referred to one of the five or six centres in Britain with a special interest in this difficult and disabling disease?

W J JEFFCOATE

Consultant Physician,
City Hospital,
Nottingham NG5 1PB

- Plotz CM, Knowlton AI, Ragan C. Natural history of Cushing's syndrome. *Am J Med* 1952;13:597-614.
- Scott HW, Liddle GW, Mulherin JL, McKenna TJ, Stroup SL, Rhamy RK. Surgical experience with Cushing's disease. *Ann Surg* 1977;185:524-34.
- Jeffcoate WJ, Rees LH, Tomlin S, Jones AE, Edwards CRW, Besser GM. Metyrapone in long term management of Cushing's disease. *Br Med J* 1977;iii:215-7.
- Fishman LM, Liddle GW, Island DP, Fleischer N, Kuchel O. Effects of amino-glutethimide on adrenal function in men. *J Clin Endocrinol Metab* 1967;27:481-90.
- Nelson DH, Meakins JW, Dealy JB, Matson DD, Emerson K, Thorn GW. ACTH-producing tumor of the pituitary gland. *N Engl J Med* 1958;259:161-4.
- Kelly WF, Macfarlane IA, Longson D, Davies D, Sutcliffe H. Cushing's disease treated by total adrenalectomy: long term observations of 43 patients. *Q J Med* 1983;52:224-31.
- Moore TJ, Dluhy RG, Williams GH, Cain JP. Nelson's syndrome: frequency, prognosis and effect of prior pituitary irradiation. *Ann Intern Med* 1976;85:731-4.
- Cushing H. The Weir Mitchell: surgical experiences with pituitary disorders. *JAMA* 1914;93:1515-25.
- Cushing H. The basophil adenomas of the pituitary body and their clinical manifestations (pituitary basophilism). *Bulletin of the Johns Hopkins Hospital* 1932;50:127-95.
- Blair Bell W. *The pituitary gland*. London: Ballière, Tindall, and Cox, 1919.
- Hardy J. Transsphenoidal hypophysectomy. *J Neurosurg* 1971;34:582-94.
- Fahlbusch R, Buchfelder M, Muller DA. Transsphenoidal surgery for Cushing's disease. *J R Soc Med* 1986;79:262-9.
- Howlett TA, Rees LH, Besser GM. Cushing's syndrome. *Clin Endocrinol Metab* 1985;14:911-45.
- Hardy J. Cushing's disease: 50 years later. *Can J Neurol Sci* 1982;9:375-80.
- Thomas JP, Richards SH. Long term results of radical hypophysectomy for Cushing's disease. *Clin Endocrinol* 1983;19:629-36.
- Brand IR, Dalton GA, Fletcher RF. Long-term follow up of trans-sphenoidal hypophysectomy for Cushing's disease. *J R Soc Med* 1985;78:291-3.
- Sharpe GF, Kendall-Taylor P, Precott RWG, et al. Pituitary function following megavoltage therapy for Cushing's disease: long term follow up. *Clin Endocrinol* 1985;22:169-77.
- Orth DN, Liddle GW. Results of treatment in 108 patients with Cushing's syndrome. *N Engl J Med* 1971;285:243-7.
- Ahmed SR, Shalet SM, Beardwell CG, Sutton ML. Treatment of Cushing's disease with low dose radiation therapy. *Br Med J* 1984;289:643-6.
- Lawrence JH, Tobias CA, Linfoot JA, Born JL, Chong CY. Heavy particle therapy in acromegaly and Cushing disease. *JAMA* 1976;235:2307-10.
- Jennings AS, Liddle GW, Orth DN. Results of treating childhood Cushing's disease with pituitary irradiation. *N Engl J Med* 1977;297:957-62.
- Burch W. A survey of results with transsphenoidal surgery in Cushing's disease. *N Engl J Med* 1983;308:103-4.
- Jeffcoate WJ, Dauncey S, Selby C. Restoration of dexamethasone suppression by incomplete adenomectomy in Cushing's disease. *Clin Endocrinol* 1985;23:193-9.
- Lamberts SWJ, Klijn JGM, De Jong FH. The definition of the recurrence of pituitary dependent Cushing's syndrome after transsphenoidal operation. *Clin Endocrinol* 1987;26:707-12.
- Sandler LM, Richards NT, Carr DH, Mashiter K, Joplin GF. Long term follow up of Cushing's disease treated by interstitial irradiation. *J Clin Endocrinol Metab* (in press).
- Krieger DT, Amorosa L, Linick F. Cyproheptadine-induced remission of Cushing's disease. *N Engl J Med* 1975;293:893-6.
- Jones MT, Gillham B, Beckford U, et al. Effect of treatment with sodium valproate and diazepam on plasma corticotrophin in Nelson's syndrome. *Lancet* 1981;ii:1179-81.
- Edwards CRW, Jeffcoate WJ. Bromocriptine and the adrenal cortex. In: *Pharmacological and clinical aspects of bromocriptine*. Southwick, Sussex: Grange Press, 1977:43-9.

Acanthamoeba keratitis

In 1975 a new infective agent that failed to respond to any treatment—*acanthamoeba*—was reported in patients with persistent keratitis.^{1,2}

Many micro-organisms capable of inducing infective keratitis had already been described, and ophthalmologists were surprised to learn that a hitherto obscure organism could cause serious eye infection. Since 1975 further reports have appeared.³⁻⁵

Acanthamoebas are free living amoebas, with trophozoites about 30 µm in diameter. They produce acanthopodia, which project from the surface of their pseudopodia and form double walled cysts. The organisms are found in brackish water and sea water and they have caused fatal granulomatous amoebic encephalitis.

In the eye *acanthamoebas* induce a persistent keratitis and uveitis. The infection occurs most commonly after trauma, the wearing of contact lenses, or ulcerative herpes simplex keratitis. The keratitis may be misdiagnosed as herpes simplex keratitis; yet the clinical appearance is characteristic (figure). Usually there is a central stromal opacity with an epithelial deficit and a pronounced surrounding ring of inflammatory infiltration.

Corneal ulcer in a contact lens wearer due to *acanthamoeba* infection.

Patients have severe recurrent attacks of keratitis that are at their worst when topical steroids are stopped. One of the most unpleasant aspects of the keratitis is the severe pain and photophobia, and on p 287 Dr David Seddon describes his harrowing personal experience of *acanthamoeba* keratitis.

Often there is considerable delay before the diagnosis is made. Many of the patients have had corneal grafts, and sometimes the diagnosis has been made from the cysts seen in the recipient button when it is examined histologically. But the diagnosis can be made more easily by taking biopsy specimens and examining them by immunofluorescent techniques⁶ or by inoculating them into Culbertson's medium for *acanthamoeba*. Treatment should begin with propamidine and neomycin drops. This may obviate the need for a corneal graft⁷; these are not always successful because organisms persist in the host tissue.

Many patients with *acanthamoeba* keratitis have recently worn contact lenses, and the single most important risk factor in case control studies has been using home made saline preparations to clean contact lenses. Therefore contact lens wearers should use properly prepared and sterilised clean solutions, a message that must be emphasised.

D L EASTY

Professor of Ophthalmology,
Bristol Eye Hospital,
Bristol BS1 2LX

- Jones DB, Visvesvara GS, Robinson NM. *Acanthamoeba* polyphage keratitis and *acanthamoeba* uveitis associated with fatal meningoencephalitis. *Trans Ophthalmol Soc UK* 1975;95:221-32.
- Watson PG. Amoebic infection of the eye. *Trans Ophthalmol Soc UK* 1975;95:204-6.
- Hirst LW, Green RW, Merz W, et al. *Ophthalmology* 1984;91:1105-11.
- Mannis MJ, Tamaru R, Roth AM, et al. *Acanthamoeba* sclerokeratitis: determining diagnostic criteria. *Arch Ophthalmol* 1986;104:1313-7.
- Yeoh R, Warhurst DC, Falcon MG. *Acanthamoeba* keratitis. *Br J Ophthalmol* 1987;71:500-3.
- Epstein RJ, Wilson LA, Visvesvara GS. Rapid diagnosis of *acanthamoeba* keratitis from corneal scrapings using indirect fluorescent antibody staining. *Arch Ophthalmol* 1986;104:1318-21.
- Wright P, Warhurst D, Jones BR. *Acanthamoeba* keratitis successfully treated medically. *Br J Ophthalmol* 1985;69:778-81.