Antigen detection in primary HIV infection

M VON SYDOW, H GAINES, A SÖNNERBORG, M FORSGREN, P O PEHRSON, Ö STRANNEGÅRD

Abstract

Serial blood samples were obtained from 21 homosexuals who had developed symptomatic primary infection with human immunodeficiency virus (HIV) after a median incubation time of 14 days. During the first two weeks after the onset of illness HIV antigen (p24) was detected in the blood by enzyme linked immunosorbent assay (ELISA). During the second and third weeks after the onset of illness p24 antibody was detected by Western blot assay and antigen concentrations rapidly decreased to undetectable values. Dissociation of antigen-antibody complexes showed complexed antigen during the phase of declining concentrations of free antigen. Neither free nor complexed antigen was detected in any serum samples for several months thereafter, which suggested that failure to detect HIV antigen reflected low or absent synthesis of viral protein rather than masking of antigen by antibodies. Reappearance of HIV antigen with a fall in p24 antibody concentration was observed in a few patients six months or more after the onset of disease.

The combined use of antigen and antibody assays made it possible to obtain evidence of infection with HIV in all of the 95 serum samples tested, illustrating the usefulness of these assays for diagnosing infection with HIV in its early stages.

Introduction

Enzyme immunoassays to detect antigens of the human immunodeficiency virus (HIV) in body fluids or the supernatants of cell cultures have recently been developed.^{1,2} HIV antigen has been detected in the serum of asymptomatic carriers of the virus and patients with persistent generalised lymphadenopathy, the acquired immune deficiency syndrome (AIDS), and AIDS related complex with a prevalence that is correlated to the stage of infection with HIV.^{2,3} Concentrations of HIV antigen in the blood have been reported to correlate inversely with the concentrations of antibodies to the core protein p24.^{2,3} Though the presence of the antigen during various late stages of infection with HIV is quite well known, its appearance during the early stages has been incompletely studied. HIV antigen has been detected during the early stages of infection in haemophiliacs⁴ and male homosexuals.^{1,3} In these patients no history of clinical symptoms or disease associated with seroconversion was reported. Recently HIV antigen was detected in serum during the early stages of the disease in a few patients with clinical evidence of primary infection with HIV.⁵⁻⁷ We were able to obtain serial blood samples from 21 homosexual men for whom the time of onset of symptomatic primary infection with HIV was known. We report the appearance of HIV antigen and its relation to the antibody response in these patients.

Department of Virology, Central Microbiological Laboratory of Stockholm County Council, PO Box 70470, S-107 26 Stockholm, Sweden

M VON SYDOW, MD, clinical virologist
H GAINES, MD, registrar in clinical virology
A SÖNNERBORG, MD, registrar in clinical virology
M FORSGREN, MD, PHD, senior lecturer
Ö STRANNEGÅRD, MD, professor

Department of Infectious Diseases, Roslagstull Hospital, Karolinska Institute, Stockholm

P O PEHRSON, MD, internist

Correspondence to: Dr von Sydow.

Patients and methods

Altogether 95 serum samples were obtained from 21 male homosexuals with clinical illness associated with seroconversion of HIV (established with different enzyme linked immunosorbent assays (ELISA) and immunofluorescence, Western blotting, and radioimmunoprecipitation assays.) Primary infection with HIV was suggested by fever in all 21 patients, sore throat and lymphadenopathy in 20, rash in 18, leucopenia in 17, and thrombocytopenia in 13. Vomiting was noted in 10, myalgia in nine, and diarrhoea in eight; two had signs of meningoencephalitis and polyradiculitis. The incubation time for infection with HIV ranged from 11 to 28 days (median 14 days) in nine patients in whom the date of infection could be established with reasonable certainty.

Antibodies could be detected by radioimmunoprecipitation (gp160) and Western blotting (p24) at the very earliest on the 11th day and 13th day, respectively, after the onset of illness. Details of the serological methods and patterns have been reported.⁸ The antibody determinations described below were obtained by Western blot assay (Biotech), Abbott human T cell lymphotropic virus type III (HTLV-III) confirmatory enzyme immunoassay, and a second generation ELISA (Abbott recombinant HIV-1 enzyme immunoassay).

HIV antigen was assayed by an antigen capture ELISA (Abbott Laboratories) as described previously.^{1,2} This antigen test primarily detects the core protein p24. Briefly, polystyrene beads coated with human polyclonal anti-HIV IgG were incubated overnight at room temperature with 200 µl of undiluted serum sample. After washing rabbit anti-HIV IgG was added and the beads incubated for four hours at 40°C. After a second wash goat antirabbit IgG labelled with horseradish peroxidase was added and incubated at 40°C for two hours. Orthophenylenediamine was the substrate for the enzyme reaction, and the absorbance was read at 492 nm.

The method of Griswold was used to detect antigen complexed with antibodies, which entailed acidifying then neutralising the serum samples as follows.⁹ A mixture of 100 µl serum and 25 µl 0.5 M hydrochloric acid was incubated at room temperature for 90 minutes. After neutralisation with 25 µl 0.5 M sodium hydroxide the incubation mixture was diluted to a final volume of 200 µl with phosphate buffered saline and then immediately added to beads coated with HIV antibody. Thereafter the antigen capture ELISA was performed as described above. Control samples were treated identically except that phosphate buffered saline was used instead of hydrochloric acid and sodium hydroxide. The amounts of HIV antigen were determined after comparison with standard curves obtained from reference samples of HIV antigen treated with buffer or hydrochloric acid.

Results

DETECTION OF HIV ANTIGEN

Antigen was detectable on one or several occasions in all of the 13 patients for whom serum samples obtained one to 18 days after the onset of illness were available (fig 1). From the spectrophotometric absorbance values HIV antigen concentrations were in most cases high in the early serum samples, thereafter declining rapidly to undetectable values. A few of the serum samples, however, gave borderline absorbance values—for example, those taken on day 5 from two patients gave absorbance values that were slightly below the cut off value recommended by the manufacturer when tested initially but were above that value when the test was repeated. All 15 serum samples (from 11 patients) taken one to 12 days after the onset of disease contained detectable antigen; four out of eight samples (from eight patients) taken 13-18 days and none of seven (from seven patients) taken 21-33 days after the onset of disease.

RELATION BETWEEN OCCURRENCE OF P24 ANTIGEN AND ANTIBODY

During the first week of illness p24 antigen was detectable in the absence of antibodies in all 12 serum samples obtained from 11 patients (table I).

During the second and third weeks antigen was detected in seven out of 12 serum samples. In four of the patients p24 antigen and antibody were both detected; and in another four patients antibodies but not antigen were

detectable. In one patient neither p24 antibodies nor antigen could be detected in a serum sample taken 14 days after the onset of illness, although a sample drawn on day 5 contained detectable amounts of antigen; the second generation antibody ELISA, however, was reactive with a sample taken on day 14 from this patient.

FIG 1—Amounts of HIV antigen in serial serum samples from 14 patients during early stages of infection with HIV. Closed circles represent values above and open circles values below cut off point; asterisks indicate values for two serum samples that gave borderline results in initial tests but in subsequent tests gave values above cut off point.

TABLE 1—Occurrence of p24 antigen and antibody at different stages of primary infection with HIV. Figures are numbers (percentages) of serum samples showing specified reactivity

Pattern of reactivity	Weeks after onset of symptoms					Total (n=95)
	1 (n=12)	2-3 (n=12)	4-13 (n=30)	14-52 (n=23)	>52 (n=18)	
Positive for antigen	12 (100)	3 (25)			2 (11)	17 (18)
Negative for antibody						
Positive for antigen		4 (33)		4 (17)	3 (17)	11 (12)
Positive for antibody						
Negative for antigen		4 (33)	30 (100)	19 (83)	13 (72)	66 (69)
Positive for antibody						
Negative for antigen		1 (8)†				1 (1)†
Negative for antibody						
Positive for antigen or antibody, * or both	12 (100)	11 (92)	30 (100)	23 (100)	18 (100)	94 (99)

*By Western blot assay.

† Found to be reactive in a second generation ELISA.

During the latter part of the first year of disease p24 antibodies were always detectable, whereas antigen was undetectable during weeks 4-13 and only occasionally (in four out of 23 samples) during weeks 14-52. Samples taken later than one year after the onset of illness were positive for antigen (five out of 18), while p24 antibodies at that time were undetectable in two samples.

All 95 serum samples showed serological evidence of HIV infection when the assays used to detect HIV antigen and antibodies were combined.

REAPPEARANCE OF HIV ANTIGEN

In one patient, who was positive for antigen during the early stages of primary infection but then became negative, the antigen reappeared about

six months after the onset of infection and then persisted (table II). Antibodies to p24 disappeared simultaneously, whereas antibodies to the envelope protein (gp41) were detectable throughout by a competitive immunoassay using HIV core and envelope proteins produced from recombinant DNA (Abbott confirmatory enzyme immunoassay). Western blotting, however, detected p24 antibodies even though the bands gradually waned in later samples. During an observation time of two years the patient was subjectively well, although he showed signs of a fairly pronounced immunological disturbance: counts of T cells bearing the CD4 molecule two weeks after the onset of illness were $0.2 \times 10^9/l$ and two years later $0.28 \times 10^9/l$ (mean value in nine patients with primary HIV infection $0.52 \times 10^9/l$ on both occasions).

TABLE II—Time course of appearance of different serological markers of infection with HIV in patient with primary asymptomatic infection

Serological marker	Days after onset of disease							
	13	28	35	56	166	280	490	700
HIV antigen by ELISA	++	—	—	—	+	+	+	+
p24 Antibody by competitive ELISA	—	+	+	+	—	—	—	—
gp41 Antibody by competitive ELISA	—	+	+	+	+	+	+	+
p24 Antibody by Western blot assay	—	+	+	+	+	+	±	—

FIG 2—Appearance of free (hatched) and complexed (white) antigen in 21 serum samples from 10 patients with primary infection with HIV. Antigen concentrations are expressed in units, one unit corresponding to about 0.2 pg of p24 antigen. Results for seven serum samples obtained during days 44-100, all with undetectable concentrations of free and complexed antigen, are not shown.

Another patient who was positive for HIV antigen six and 12 months after the onset of illness (serum samples from the time of early infection were not available for determination of antigen) also had counts of T cells bearing the CD4 molecule of around $0.20-0.30 \times 10^9$. Concentrations of p24 antibody decreased appreciably during the period of follow up to values that were barely detectable by Western blotting. In a third patient who was initially antigenaemic antigen disappeared and then reappeared between three and four years later. At that time he had a progressive neurological disease, which was taken to be a manifestation of HIV infection. In the early stages of his disease T cells were not investigated, but after four years the counts of T cells bearing the CD4 molecule were low (less than $0.10 \times 10^9/l$). The appearance of HIV antigen was also associated with an appreciable decrease in p24 antibody to undetectable concentrations.

Antigen was not detected in eight other patients followed up for more than one year after their primary infection. None of these patients developed clinical signs of persistent HIV infection or profound changes in counts of T cells bearing the CD4 molecule.

HIV antigen was not detected in serum samples taken from six patients between nine and one and a half months before the onset of symptoms.

DETECTION OF COMPLEXED HIV ANTIGEN

Twenty one serum samples obtained from 10 patients at different times during the first six months after the onset of disease were tested for concentration of free and complexed antigen after dissociation of antigen-

antibody complexes. Complexed antigen was detected in samples taken during the phase of declining concentrations of free antigen but not in those drawn four and five days after the onset of illness (fig 2). Twelve samples drawn between days 18 and 100 showed no free or complexed antigen, except for one on day 25, which seemed to contain small amounts of complexed antigen. Samples from two different patients drawn on days 124 and 166 contained both free and complexed antigen. Four out of 15 samples from 11 patients drawn between seven months and three years after the onset of disease were positive for antigen (data not shown). Two of these four samples were drawn from one patient nine months and two years after the onset of disease and contained more complexed than free antigen, whereas the other two from another patient contained only free antigen.

Discussion

During the early phase of primary infection with HIV infectious virus can be detected in plasma.¹⁰ We showed previously that this early viraemia is commonly associated with the presence of detectable HIV antigen in the blood.⁵ In other studies HIV antigenaemia has been found to precede seroconversion in haemophiliacs⁴ and homosexuals³ infected with HIV and was also recently shown in patients with clinical primary HIV infection.^{6,7} The prevalence and time course of HIV antigenaemia in early stages of infection have, however, not been elucidated previously; the availability of serial blood samples from 21 homosexual men with symptomatic primary HIV infection was a good opportunity for us to study different aspects of HIV antigenaemia. As we were able to show HIV antigen in all of the 13 patients for whom serum samples were available during the first 18 days of illness we conclude that HIV antigenaemia consistently occurs during the early phase of symptomatic primary infection with HIV.

By using the antigen assay in combination with sensitive tests for HIV antibodies we were able to diagnose HIV infection in all the patients at any stage of the infection after onset of symptoms. Thus the existence of a "window"—that is, a time after infection when neither antigen nor antibody can be detected—may be of minor importance in symptomatic primary infection provided that sensitive assays are being used. Further studies are needed to establish whether this conclusion is also valid in cases of asymptomatic primary HIV infection. We emphasise, however, that to minimise the importance of the window antibody assays must be able to detect the early antibody responses. Currently used first generation commercially available tests often do not detect antibody responses until two to seven weeks after the onset of symptoms of infection,⁸ but with second generation ELISAs earlier detection is possible.¹¹ Also, as serum concentrations of HIV antigen may be low even during the acute stage of infection accurate diagnosis of early HIV infection requires the use of antigen tests with high sensitivity.

The rapid decline in antigen concentration observed about two weeks after the onset of disease could be due to the formation of immune complexes. In fact, evidence of such formation was obtained after serum samples had been subjected to a low pH to split up the complexes, a method successfully used by Mathieson to show complexed antigen and antibodies in patients infected with HIV (Mathieson T, second international conference of neuroimmunology, Philadelphia 1987). The time of appearance of detectable immune complexes coincided with the early antibody response as determined by assays of IgM and sensitive assays for IgG (H Gaines *et al*, unpublished data). As complexed antigen was not detectable in appreciable amounts during the first few months after the acute stages of infection it seems that the lack of detectable antigen at that time reflected low or absent synthesis of viral antigens rather than masking of antigen by antibodies. Failure to detect antigens, however, does not necessarily reflect the complete disappearance of infectious virus from the circulation as in some cases infectious virus may persist in the absence of detectable antigenaemia.⁵

In other studies circulating immune complexes, at least partly consisting of HIV antigen and antibodies, have been shown during later stages of HIV infection.¹²⁻¹⁴ The pathogenetic implications of the appearance of circulating immune complexes during the early and late phases of infection with HIV remain to be established.

Our study clearly shows that HIV antigenaemia wanes about two weeks after the onset of illness in symptomatic primary infection

with HIV. In other studies HIV antigen has been shown to occur during the later stages of the disease.^{1,3} Thus HIV antigenaemia seems to follow a diphasic course, initially appearing during the acute stage and then after varying periods of time reappearing in some patients; two of our patients showed this pattern. Most of our patients were followed up for only a limited period, and the prognostic importance of the reappearance of antigen remains to be established in further, prospective studies. A correlation exists, however, between antigenaemia and stage of disease,^{2,3} and the occurrence of HIV antigen in the blood seems to be an unfavourable prognostic sign.¹⁵

References

- Goudsmit J, Paul DA, Lange JMA, *et al*. Expression of human immunodeficiency virus antigen (HIV-Ag) in serum and cerebrospinal fluid during acute and chronic infection. *Lancet* 1986;ii:177-80.
- Paul DA, Falk LA, Kessler HA, *et al*. Correlation of serum HIV antigen and antibody with clinical status in HIV-infected patients. *J Med Virol* 1987;22:357-63.
- Lange JMA, Paul DA, Huisman HG, *et al*. Persistent HIV antigenaemia and decline of HIV core antibodies associated with transition to AIDS. *Br Med J* 1986;293:1459-62.
- Allain JP, Laurian Y, Paul DA, Senn D. Serological markers in early stages of human immunodeficiency virus infection in haemophiliacs. *Lancet* 1986;ii:1233-6.
- Gaines H, Albert J, von Sydow M, *et al*. HIV antigenaemia and virus isolation from plasma during primary HIV infection. *Lancet* 1987;i:1317-8.
- Wall RA, Denning DW, Amos A. HIV antigenaemia in acute HIV infection. *Lancet* 1987;i:566.
- Kessler HH, Blaauw B, Spear J, *et al*. Diagnosis of human immunodeficiency virus infection in seronegative homosexuals presenting with an acute viral syndrome. *JAMA* 1987;258:1196-9.
- Gaines H, von Sydow M, Sönnnerborg A, *et al*. Antibody response in primary human immunodeficiency virus infection. *Lancet* 1987;i:1249-53.
- Griswold WR. A modified Farr technique for the detection of antibody in the presence of antigen-antibody complexes. *J Immunol Methods* 1981;40:73-8.
- Albert J, Gaines H, Sönnnerborg A, *et al*. Isolation of human immunodeficiency virus (HIV) from plasma during primary HIV infection. *J Med Virol* 1987;23:67-73.
- Lelie PN, Reesink HW, Huisman JG. Earlier detection of HIV and second generation antibody assays. *Lancet* 1987;iii:343.
- Morrow WJW, Wharton M, Stricker RB, *et al*. Circulating immune complexes in patients with acquired immune deficiency syndrome contain the AIDS associated retrovirus. *Clin Immunol Immunopathol* 1986;40:515-24.
- Lange JMA, Paul DA, de Wolf F, *et al*. Viral gene expression, antibody production and immune complex formation in human immunodeficiency virus infection. *AIDS* 1987;1:15-20.
- Carini C, Mezzaroma I, Scano G, *et al*. Characterization of specific immune complexes in HIV-related disorders. *Scand J Immunol* 1987;26:21-8.
- de Wolf F, Goudsmit J, Paul DA, *et al*. Risk of AIDS related complex and AIDS in homosexual men with persistent HIV antigenaemia. *Br Med J* 1987;295:569-72.

(Accepted 30 November 1987)

ONE HUNDRED YEARS AGO

Cold weather and Christmas cheer are invigorating to most of us at this time of year, but it is a sad time for underfed children. The committee for supplying vegetarian dinners to school children in London have been actively prosecuting their good work this winter. We are glad to hear that their dinners have proved palatable and acceptable to the children. Their experiment is a fair one; but we should like to know whether the vegetarian dinners aided growth in the children as much as the mixed diets supplied, at equally low rates, to the school children in Birmingham. It does not appear that vegetarian dinners are more economical as to prime cost; it may also be doubted whether they are more economical physiologically. No doubt the ingredients necessary for the nutrition of the body are to be found in vegetarian food; but evidence is not forthcoming that the digestive organs are able to extract and assimilate nitrogenous matters as well from a purely vegetable diet as from a mixed one. Plants get their supplies of nitrogen from compounds in the soil, not from the atmosphere. The feeding of children is at least as necessary to their development as teaching, and the cost is trifling in comparison to some of the results obtained: 19,673 dinners were supplied at £68 12s., or a little over three farthings per head. We may safely say that this admirable movement will do good, and that it is economically worked; still, in any plan for distributing food to the poor, there remains the necessity of avoiding pauperisation. The independence of the poor is best respected by enabling them to help themselves; and we are glad to see the suggestion made that senior girls in the school should be employed in cooking and serving the food. Teachers in poor schools are worked hard enough, and must not be expected to do much beyond their paid duties. It is to be hoped that well cooked and comfortably served dinners are served in the middle of the day for the teachers in the schools.

(*British Medical Journal* 1888;i:33)