confident that we can attract the private funding element once our investment needs are acknowledged and a generous response from government is forthcoming.

In the meantime, we will be working continually to improve the physical conditions in which the people of Brixton, Camberwell, Dulwich, and Peckham receive their emergency treatment.

CROSSLEY SUNDERLAND
General Manager

EDWARD GLUCKSMAN
Accident and Emergency Consultant

King's College Hospital,
London SE5 9RS

Micrometastases in bone marrow in patients with breast cancer

SIR,—Having read the letter from Dr R W Blamey and others (5 December, p 1487) on the paper from Dr Janine L Mansi and her colleagues (31 October, p 1093), we write on the basis of our own currently available results.

Our group has also been studying the development of bone metastases in patients with breast cancer after detection of tumour cells in bone marrow at the time of primary treatment. Although the number of patients in our study is smaller (68 patients; T₁₋₃ N₀₋₂ M₀₋₁) and our median follow up time is 16 months (maximum 36, minimum 3), our observations support the hypothesis that the presence of micrometastases is a predictor for the subsequent development of bone metastases (table I).

Our method differs slightly from that of Dr Mansi and others in that we make use of a cocktail of monoclonal antibodies consisting of anti-EMA and anticytokeratin. Bone marrow sampling (in conjunction with immunocytochemistry) has allowed us to detect bone micrometastases in 36.7% of our patients (compared with 26.4% in the above study). Similarly, our results indicate a

TABLE I—Relation between presence of micrometastases in bone marrow of patients with primary breast cancer and site of first distant relapse

Site of relapse	Micro-metastases present (n=25)	Micro-metastases absent (n=43)
Bone	7	0
Visceral	1	2
Soft tissue (lymph nodes, skin)	2	1
Mixed (viscera and bone)	2	0

TABLE II—Comparison of occurrence of micrometastases with other clinical prognostic factors. Results are numbers (and percentages)

	No of patients	Micro-metastases present (n=25)	Micro-metastases absent (n=43)
T ₁	30	10 (34)	20 (66)
T ₂	26	9 (35)	17 (65)
T ₃	3	1 (33)	2 (67)
T ₄	5	2 (40)	3 (60)
T _x	4	3 (75)	1 (25)
No nodes	26	12 (46)	14 (54)
1-3 Nodes	22	7 (32)	15 (68)
>3 Nodes	20	6 (30)	14 (70)
Oestrogen receptor:			
Positive	33	7 (21)	26 (79)
Negative	24	11 (46)	13 (54)
Not tested	11	7 (64)	4 (36)
Premenopausal	29	10 (34)	19 (66)
Postmenopausal	39	15 (38)	24 (62)
Died	7	5 (71)	2 (29)

correlation between positive patients and T stage and no correlation with the menopausal state. Unlike Dr Mansi and her colleagues, however, we found a correlation with the presence of oestrogen receptors and did not find a correlation with spread to lymph nodes (table II).

To determine whether the differences in our observations are significant we need to continue and extend our study. In any case, it appears likely that the methods used by Dr Mansi and others and by our group may provide an additional prognostic factor (though not completely independent) to identify a high risk group for bone metastases at the time of primary treatment.

M UNTCH
NADIA HARBECK
W EIERMANN

Department of Gynaecology and Obstetrics,
Grosshadern Hospital,
Munich, West Germany

Charging patients for eye tests

SIR,—Several issues stemming from the government's white paper on primary care have been discussed in your columns, but their fuller implications for eye health have yet to be debated.

The general ophthalmic service has an important screening function and has regularly been a major source of referral of eye disorders to the hospital eye service. A good example of this preventive role is found in chronic simple glaucoma, which is responsible for about 12% of the registered blind in England and Wales and affects 150 000 to 300 000 of the population.¹ About 80% of those whose diagnosis is confirmed in an eye department are referred by opticians or ophthalmic medical practitioners.² This screening function might be lost if the system of free sight tests were to be abandoned.

Mrs Currie has suggested that commercial pressures would stimulate opticians to offer free sight tests so that the status quo would be unaltered. If the government is confident of this effect why does it offer exemptions to children, young people in full time education, and those on low incomes? Old age pensioners, who carry the greatest risk of blinding eye disease, are not included in these exemptions.

The effect of the new proposals on ophthalmic medical practitioners does not appear to have been considered. Ophthalmic medical practitioners have no financial interest in the sale of spectacles and therefore would be obliged to charge for a sight test. If opticians do not charge then a financial differential between the ophthalmic medical practitioner and optician could arise which might deprive the public of the practitioner's medical skills.

If, on the other hand, opticians do charge for sight testing there is a real possibility that patients whose only problem is an error of refraction will be diverted into the hospital eye service to make demands on already overburdened resources, causing further delays for those suffering from organic eye diseases.

Another, and most unwelcome, differential arises from paragraph 5.4 of the white paper, which states: "Without the constraints imposed by the NHS contract, opticians will be free to offer either a standard sight test or a fuller eye examination to meet their patients' needs and preferences." It is difficult to avoid the conclusion that a two tier, fee based system is being proposed with the most comprehensive assessments available for those who can afford it.

Although the white paper emphasised that its proposals were the fruit of wide ranging discussions and consultation, the proposals were not put to the Faculty of Ophthalmologists. If the government really wants to promote better health it should stop

regarding its provision as a marketing exercise, regarding patients as "consumers," and confusing legitimate propaganda with a public relations job. It also needs to learn to take advice and to take it from the right quarters.

ANTHONY J BRON

Oxford Eye Hospital,
Oxford OX2 6AW

- 1 Crick RP. Computerised clinical data base for glaucoma—ten years' experience. *Res Clin Forums* 1980;2:29-39.
- 2 Elkington AR, Lewry J, MacKean J, Sargent P. A collaborative hospital glaucoma survey. *Research and Clinical Forums* 1982;4(5):32-40.

SIR,—While full of good intentions Dr John S Yudkin's letter (2 January, p 64) indicates some degree of misunderstanding of the general ophthalmic services sight test. This is a test designed to see if a person (who is not at that stage a patient) has a defect of sight which can be corrected by an optical appliance. "Defect of sight" appears to be parliamentary jargon for a refractive error and has nothing to do with disease—in fact, it is specifically stated elsewhere in the act that the defect referred to is of an anatomical or physiological nature.

Thus the referral of a patient to the general optical services for the specific purpose of excluding diabetic retinopathy would be an inappropriate use of this particular service.

The other aspect of the matter, as to whether in a routine refraction an optician is likely to pick up diabetes or, more specifically, diabetic retinopathy, is rather different. The suggestion here, as with glaucoma, is that in the course of a sight test a medical screening will also occur.

This is a dangerous assumption. It is true that opticians, via their Eye Care Information Bureau, are not above advertising their self appointed quasi medical role with such eye catching headlines as "Thirty minutes that can add years to your life" (a recent advertisement in *The Times* in which diabetes is specifically mentioned), but such advertising is for commercial purposes and has little to do with genuine screening as envisaged by Dr Yudkin.

In my experience remarkably few new cases of diabetic retinopathy are picked up by opticians. I cannot remember one in the past two years. (I am responsible for reading 3500 referral letters to five consultants a year.) Nevertheless, many patients with various suspected conditions, mostly glaucoma, are referred via rather a wide bore general practitioner filter with the result that our outpatient departments are becoming overloaded. Waiting times are over a year in some centres and in most are over three months.

Something has to be done to stem the flood of false positive opticians' referrals and perhaps this is the point at which some thought might be given to using the ophthalmic medical practitioner in a more useful role. If ophthalmic medical practitioners were used to supervise opticians' referrals in some way instead of carrying out routine refractions (and getting paid less than opticians for it) they could possibly help to bring some order to an increasingly chaotic situation.

REDMOND J H SMITH

London W1N 1AA

SIR,—We share Dr John S Yudkin's concern (2 January, p 64) about the screening of diabetic patients to detect early retinopathy.

However, he makes a number of assumptions which flaw his argument in favour of ophthalmic opticians taking part in the primary care of patients with diabetes.

Opticians do indeed detect fundal abnormalities