CORRESPONDENCE

Cost of limiting abortion Madeleine Simms, MA.....	355	Secretor state of patients with insulin dependent or non-insulin dependent diabetes mellitus C C Blackwell, PHD	357	Light on seasonal affective disorders? R A Weale, PHD	359
Diagnosis of cystic fibrosis P S Baxter, MRCP, and others	355	Logic in medicine: the role of leads D E H Llewelyn, MRCP.....	357	Trusting patients with case notes Diana Elbourne and Alice Lovell	360
Which treatments work for drinking problems? G Milner, FRCPsych	356	Recent theories on the cause of cot death A M Stewart, FRCP	358	Plastic surgery J M Porter, FRCS	360
The future of medical microbiology N A Simmons, FRCPath	356	Computerised advice on malaria prevention and immunisation J H Cossar, MB, and others	358	Nursing grievances A J Byrne, BPharm	360
Response to deoxycytosine in T cell leukaemias D Catovsky, FRCPath, and Claire Dearden, MRCP	356	Rare tumour system J J Bolger, FRCR	358	Hospital and community health service costs: England and Scotland compared A Maynard, BPhil, and S Birch, DPhil	360
Gonadotrophin hormone releasing analogues open new doors in cancer treatment J Waxman, MRCP	356	Intervention by general practitioners to reduce smoking M G Goldstein, MD, and R Niaura, PHD.....	358	Points The malignant neuroleptic syndrome (A M P Kellam); The unquiet 1960s (B D M Williams); Public Health Laboratory Service pay (B Grotrex); The Royal College of Psychiatrists and South Africa (A King); Testing paternity (P A Webb and A J Jeffreys); Lactose absorption and milk drinking (J J Segall); Mattress contamination (P Lowthian); Sexism in medical writing (Francis Sandor); Hemlock poisoning (A Hollman)	361
Obstetricians on the labour ward M J Hare, MRCOG	356	Cocktail stick injuries W R Grandsen, MRCPath, and Susannah Eykyn, FRCPath	359	Correction: Response to deoxycytosine in mature T cell malignancies (Ho)	361
Diagnosis of diabetes mellitus: pitfalls in the glucose tolerance test Hilary King, MD, and others	357	Embryo research P Mallett; R M L Winston, FRCOG	359		

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.

Cost of limiting abortion

SIR,—Ms Caroline Woodroffe and Dr Valerie Beral (2 January, p 62) have done a public service by drawing attention to the large public health and social service cost implications of Mr David Alton's Abortion (Amendment) Bill. This important issue has hitherto been shrouded in taboo because it is so easy for those who calculate the cost implications of health legislation to be accused of heartlessness or worse.

The writers admit that their figures are conservative in that the costs of medical care and of family support are both excluded. Moreover, their calculations are based on 1983 figures, which have since risen. They do not touch on the emotional costs.

In 1985 I interviewed at length a small group of mothers in east London and Essex who had cared for a lifetime for severely handicapped children born just before the Abortion Act was passed and now aged between 19 and 25. Many of these mothers had developed strong views about abortion.¹ Two thirds said that with hindsight they would certainly have opted for abortion had this been available to them at that time. Several had given up all hopes of ever returning to work or resuming a career. The costs of this to the families concerned, the loss of a lifetime's earnings, are of course substantial and are also not included in Ms Woodroffe's and Dr Beral's calculations.

Among their observations are the following:

"At the time we wanted him to live, preferred that he would, but now realise this was a mistake. . . . The strain of caring has been overwhelming."

"I really had no idea what I had let myself in for. . . . Now we are imprisoned for life. . . . There is no future for us."

"The do gooders on TV make my blood boil. They have no idea what a child like this involves."

Shepperdson's pioneering study found that half the parents of children with Down's syndrome

in her sample believed that "not all handicapped children should be kept alive at all costs" and that this view increased with age.² Pahl and Quine wrote: "In an attempt to illuminate the overall reality of caring for these children, parents were asked what they would do if they knew an expected child would be severely mentally handicapped. Seventy eight per cent of parents answered that they would want the pregnancy terminated."³

The general public has always overwhelmingly supported late abortion on grounds of handicap: 87% favoured this according to the latest poll.⁴ If most parents who have themselves had to carry the heavy burdens of dependency would with hindsight have preferred to have had an abortion then the devastating emotional costs need to be taken into account as well as the economic costs in assessing the true implications of the Alton bill.

MADELEINE SIMMS

London NW11 8AG

1 Simms M. Informed dissent: the views of some mothers of severely handicapped young adults. *J Med Ethics* 1986;12:72-4.

2 Shepperdson B. Abortion and euthanasia of Down's syndrome children—the parents' views. *J Med Ethics* 1983;9:152-7.

3 Pahl J, Quine L. *Families with mentally handicapped children*. Canterbury: University of Kent, 1984.

4 Anonymous. *Report on British social attitudes*. London: Social and Community Planning Research, 1986.

Diagnosis of cystic fibrosis

SIR,—The interesting and complex case reported by Dr C H Skeoch and others (5 December, p 1445) raises several points about the diagnosis of cystic fibrosis. Their conclusion, however, that children with symptoms of cystic fibrosis and atypical features need to be checked for human

immunodeficiency virus (HIV) is too simple, as all causes of abnormal immune function, including HIV infection, should be considered. In addition, in cystic fibrosis diarrhoea is usually due to steatorrhea, which was apparently not present in this child, and this should also lead to a review of the diagnosis. Abnormal sweat chloride results are described in deprivation¹ and malnutrition,² but unfortunately these papers do not give sweat sodium concentrations. Malnutrition might have been the cause of the abnormal results of sweat tests in this girl.

Measurement of both chloride and sodium values is performed routinely on sweat in our laboratory, and our experience agrees with that of others^{3,4} that the chloride concentration is almost always higher than the sodium concentration in sweat from patients with cystic fibrosis, while healthy people have chloride values lower than sodium values. It would be interesting to know whether this is the experience in Glasgow. If so, the results in this girl should also make one reconsider the diagnosis of cystic fibrosis.

This report illustrates the pitfalls in diagnosing cystic fibrosis and re-emphasises the need for the diagnosis to be confirmed in a centre dealing with cystic fibrosis, as happened with this child. If failure to thrive is being investigated by jejunal biopsy studies of chloride secretion may also help.⁵

P S BAXTER
C J TAYLOR
E WORTHY

Departments of Paediatrics
and Chemical Pathology,
Children's Hospital,
Sheffield S10 2TH

1 Christoffel KS, Lloyd-Still JD, Brown G, Shwachman H. Environmental deprivation and transient elevation of sweat electrolytes. *J Pediatr* 1985;107:231-4.

- 2 Beck R, Durie PR, Hill JG, Levison H. Malnutrition: a cause of elevated sweat chloride concentration. *Acta Paediatr Scand* 1986;75:639-44.
- 3 Green A, Dodds P, Pennock C. A study of sweat sodium and chloride; criteria for the diagnosis of cystic fibrosis. *Ann Clin Biochem* 1985;22:171-6.
- 4 Shwachman H, Mahmoodian A, Neff RK. The sweat test: sodium and chloride values. *J Pediatr* 1981;98:576.
- 5 Taylor CJ, Baxter PS, Hardcastle J, Hardcastle PT. Absence of secretory response in jejunal biopsy samples from children with cystic fibrosis. *Lancet* 1987;ii:107-8.

Which treatments work for drinking problems?

SIR,—Professor Griffith Edwards's leader (2 January, p 4) summarises in a concise way current views on the effectiveness of treatment of drinking problems. Some of the statements in this review, however, are apt to be misinterpreted by cost conscious managers and others, who could come to the conclusion that they should leave the responsibility for treating this group of people to the voluntary and primary care agencies and decide not only to deprive mental health units of necessary funding to establish alcohol and drug services but also to consider closing these where they exist—a course of action contemplated in this district some time ago.

As has been pointed out,¹ primary care workers do not, by and large, possess the necessary skills, nor have they the time or the willingness, to treat problem drinkers. Moreover, if they were to acquire these skills it would inevitably increase the number of patients detected with higher degrees of disabilities who perhaps require specialist help.

Case register data from the Worcester Health District, where a substance abuse service has operated for some years, have indicated that the annual prevalence of substance abuse problems was 118 per 100 000 population in 1984 (105 alcohol and 13 drugs) and 151 per 100 000 in 1985 (112 alcohol and 39 drugs)² whereas the corresponding rates per 100 000 population in 1984 were 133 for schizophrenia and 147 for affective psychosis. Thus patients with alcohol and drug problems constitute a sizable and still growing proportion of the total psychiatric case load in a semirural district.

Furthermore, these patients have considerable medical, social, and legal problems, which had been previously attended to by a variety of professionals, who in turn often had not recognised that these were related to alcohol.

Many districts still do not provide specialist treatment services for this large group of patients with considerable handicaps. Capital outlay to set up a comprehensive community substance abuse service needs to be minimal. The revenue costs in the Worcester Health District are in the region of £100 000. This forms a small part of the total budget of the mental health unit, which at the moment is £7m a year.

A team of this type, integrating the voluntary and statutory sectors into one continuum of care, can effectively respond to a variety of client or patient needs, give training and support to primary care workers, and stimulate and coordinate local alcohol and drug abuse prevention strategies. I would suggest that it is also a good buy.

G MILNER

Substance Abuse Service,
Worcester Royal Infirmary,
Worcester WR5 1JG

- 1 Milner G. *Proceedings of regional symposium on alcohol and other dependencies. Warsaw, 1987.* World Psychiatric Association, 1987.
- 2 Glatt MM. Comments on "The pattern and range of services for problem drinkers." A report by the advisory committee on alcoholism to the DHSS and Welsh Office 1978. *Br J Addict* 1979;74:115-32.

The future of medical microbiology

SIR,—From representations being made to me by consultant microbiologists from all over Britain it is apparent that many are deeply concerned that some recent events could undermine not only the service that they provide but also the future of medical microbiology in Britain.

Firstly, the role of medical microbiologists in infection control is being questioned. For over 30 years they have been responsible for developing and implementing policies and procedures for controlling infection in hospitals, and they have not been unsuccessful. Hospital infection control is now an integral part of the practice of medical microbiology, and Britain leads the world in this subject. Consultant microbiologists have built up a body of knowledge and experience, which is being passed on to doctors in training, who must face questions on the subject in their examinations for membership of the Royal College of Pathologists. Unfortunately, at a time when microbiologists should be directing their undivided attention to solving such serious problems as those presented by the spread of multiple antibiotic resistant organisms and the acquired immune deficiency syndrome they are being diverted from the task.

Since the introduction of Griffiths management some district medical officers have shown a desire to assume an increased responsibility for hospital infection control. Unfortunately, although they may have statutory obligations and duties, few have the knowledge, training, and experience necessary for anything other than a managerial role. Yet microbiologists can find their authority eroded, their effectiveness reduced, and their time consumed by the new administrative arrangements.

The second area of concern results from a growing tendency to regard clinical pathology, including microbiology, as a purely technical service with laboratories looked on as mere processing units which, like slot machines, will issue results if they are fed with specimens. In this scenario the professional role of medical microbiologists working at the interface of clinical and laboratory medicine is disregarded, either because it is not appreciated or because the sort of bucket shop service which could be provided without them looks cheap. The concept of a medical microbiology service without doctors should be as unacceptable as a surgical service without surgeons. Medical microbiologists know that most of their clinical colleagues recognise the importance of the service provided. I hope that they will help us to defend it.

N A SIMMONS

Department of Clinical Bacteriology and Virology,
Guy's Hospital,
London SE1 9RT

Response to deoxycoformycin in T cell leukaemias

SIR,—Dr A D Ho (5 December, p 1483) comments on our results on the treatment of mature T cell leukaemias with deoxycoformycin (10 October, p 873). His experience does not appear to confirm a correlation between the membrane phenotype of the neoplastic T cells and the response to deoxycoformycin. We have now extended our results to 28 patients treated by deoxycoformycin (table) and these show even more clearly than before that no meaningful responses were observed in cases with a phenotype other than CD4+, CD8—.

One possible explanation for the differences observed is that the largest group of our patients,

Response to deoxycoformycin in post-thymic T cell leukaemias

Phenotype	No	Responders*	No remission
CD4+, CD8—	15	10 (67%)	5
CD4+, CD8—	8	0	8
CD4—, CD8+	5	0	5
All cases	28	10 (36%)	18

*Partial and complete remission.

11, had T polymorphocytic leukaemia,¹ a disease represented by one case in Dr Ho's study. Possibly the type of T cell disease may be just as important in relation to the effect of deoxycoformycin as the cell phenotype. For example, eight of the T polymorphocytic cases were CD4+, CD8— and six (75%) experienced a good response. In contrast only one out of six patients with adult T cell leukaemia/lymphoma (four were CD4+, CD8—) experienced a remission. We do not agree with Dr Ho that different definitions of remission could account for our results as some of our complete responders had prolonged periods (more than 20 months) free of disease without requiring further treatment.

DANIEL CATOVSKY
CLAIRE DEARDEN

MRC Leukaemia Unit,
Royal Postgraduate Medical School,
London W12 0HS

- 1 Matutes E, Talavera JG, O'Brien M, Catovsky D. The morphological spectrum of T-polymorphocytic leukaemia. *Br J Haematol* 1986;64:111-24.

Gonadotrophin hormone releasing analogues open new doors in cancer treatment

SIR,—In reply to my leading article (31 October, p 1084) Messrs A J Arnold and A D Desmond (12 December, p 1565) advocate the use of subcapsular orchidectomy for patients with prostatic cancer and suggest that "patient acceptability in elderly men is high."

At present three monthly depot injections of gonadotrophin releasing hormone analogues are under clinical investigation and have been shown to suppress testosterone over this period. If the patient is given a choice by his clinician of an injection once every three months or an operation I suggest that in almost every case he will prefer injection treatment. The cost at present is relatively high but no more expensive than that of cyproterone acetate treatment. Most drugs when introduced are expensive. This reflects development costs, and after an initial period the cost inevitably comes down.

Messrs Arnold and Desmond are advocating the use of a primitive and mutilating procedure and ignoring a clinical advance of advantage to the patient, who has enough, dealing with the idea of cancer, without having to experience surgical orchidectomy as well.

JONATHAN WAXMAN

Department of Clinical Oncology,
Royal Postgraduate Medical School,
London W2 0HS

Obstetricians on the labour ward

SIR,—Mr J S Samra and others (12 December, p 1566) express surprise at the findings of Ms Coupland and others (24 October, p 1077) and highlight directly and by implication the fears of many hospital doctors over two tier working. As a consultant obstetrician who elected to move from three tier to two tier working, may I seek to allay some of these fears.