lines which happens in medicine, the prevalence and incidence of the differential diagnoses having an important bearing on the reasoning process.⁵ The convention of substantiating the "knowledge" used by pointing to actual observations is linked closely to this type of reasoning.⁴ Perhaps "witch-doctors" (31 October, p 1108) are those who persistently scuttle away from such illuminating observations and open logical thought processes.

When we are actually talking to patients or examining them, we tend to interpret information quickly using subconscious thought processes. If we wish to explain our reasoning to patients or other doctors, however, we must use explicit, logical reasoning which often begins with "leads", and be prepared to substantiate the knowledge used. Perhaps this dual nature of the logic of medicine should have been emphasised.

DE H LLEWELYN

Department of Medicine,
King's College School of Medicine and Dentistry,
London SE5 8RX

- 1 Llewelyn DEH. Mathematical analysis of the diagnostic relevance of clinical findings. *Clin Sci* 1979;57:477-9.
- 2 Llewelyn DEH. Accuracy and value of the haemoccult test. *Br Med J* 1983;285:1144-5.
- 3 Eddy DM, Clanton CH. The art of diagnosis. *N Engl J Med* 1981;36:1263-8.
- 4 Llewelyn DEH. Applying the principle of logical elimination to probabilistic diagnosis. *Med Inf (Lond)* 1981;6:25-32.
- 5 Llewelyn DEH. The mathematics of clinical diagnosis. In: Rodbard D, Forti G, eds. *Computers in endocrinology*. New York: Raven Press, 1984:299-331.

Recent theories on the cause of cot death

SIR,—In his review (28 November, p 1366) Professor A D Milner divides all the hypotheses on the cause of cot deaths introduced in the past 10 years into four categories but fails to mention one which clearly belongs to the group which he has labelled "maturation failure."¹

This theory—which assumes that a high proportion of childhood cancers are initiated during embryogenesis²—ascribes maturation failure to a latent period effect of one of these embryomas (juvenile myeloid leukaemia). It was originally introduced 12 years ago to explain (a) why there was overrepresentation of summer and autumn births among cot deaths and underrepresentation of these births among equally young patients with leukaemia, and (b) why there are so few myeloid cases among childhood leukaemias.³ But it was reintroduced a few months ago to explain why cot deaths as well as juvenile myeloid leukaemia have associations with exceptionally high levels of fetal haemoglobin.⁴

A high proportion of embryomas among juvenile leukaemias would allow these cases to have associations with faulty maturation of blood corpuscles and consequent difficulty in replacing either passive with active immunity (faulty leucopoiesis) or fetal with adult haemoglobin (faulty erythropoiesis). The first difficulty would affect lymphatic as well as myeloid embryomas, but the second would affect only the myeloid subgroup. Furthermore, infants who had survived the neonatal period in spite of one or both difficulties would be at high risk of dying either from a minor infection (myeloid and lymphatic embryomas) or from tissue anoxia (myeloid embryomas). Finally, this difference between otherwise similar neoplasms would be more noticeable today than it was before antibiotics had reduced the risk of death from infection without affecting sudden deaths of apparently healthy infants.

In short, besides providing a reason why cot deaths are concentrated between 1 and 6 months of age, this "maturation failure" hypothesis would account for the high incidence of lymphatic

leukaemia among 2 to 4 year olds and for the emergence of relatively high mortality from leukaemia after the discovery of drugs which compensate for defective immune responses. It is also a hypothesis which invites further testing of HbF and HbA concentrations in cases of unexplained stillbirths, cot deaths, and deaths from respiratory infection in young infants.

A M STEWART

Cancer Epidemiology Research Unit,
Queen Elizabeth Hospital,
Birmingham B15 2TH

- 1 Stewart AM. Competing causes of death in young cases of myeloid leukaemia. *Paediatr Perinat Epidemiol* 1987;1:197-8.
- 2 Knox EG, Marshall T, Barling RT. Leukaemia and childhood cancer in twins. *J Epidemiol Community Health* 1984;38:12-6.
- 3 Stewart AM. Infant leukaemias and cot deaths. *Br Med J* 1975;ii:605-7.
- 4 Giulian GG, Gilbert EF, Moss RL. Elevated fetal haemoglobin levels in sudden infant death syndrome. *N Engl J Med* 1987;316:1122-6.
- 5 Weatherall DJ, Edwards JA, Donahue WTA. Haemoglobin and red cell enzyme changes in juvenile myeloid leukaemia. *Br Med J* 1968;ii:679-81.

Computerised advice on malaria prevention and immunisation

SIR,—Drs Angela J Williams and David J M Lewis (5 December, p 1449) highlight the problem of disseminating accurate antimalarial and pretravel health advice. We share their concern,¹⁻³ especially as surveys undertaken in Glasgow between 1977 and 1985 indicated that 36% of the 13 816 overseas travellers who were interviewed on their return reported illness—much of which might have been prevented. Moreover, in the past 10 years 703 patients with malaria have been recorded in Scotland.

These findings all emphasise the need for a source of continually updated information on suitable antimalarial drugs for travellers and appropriate immunisations. Since January 1987 this has been available on a database compiled by the Communicable Diseases (Scotland) Unit (CD(S)U) and installed by the Scottish Poisons Bureau Information Service on its mainframe computer. General practitioners who have access to a terminal or computer with a modem (such as that which links with Prestel) can register to use the service by contacting the CD(S)U (telephone 041 946 7120 ext 247).

This service covers recommended immunisations and malaria prophylaxis and is updated directly from the unit on the basis of advice from various sources, including the World Health Organisation, the Department of Health and Social Security, the Malaria Reference Laboratory, embassies, and drug companies. Information about the administration and availability of both common and more unusual vaccines is also provided. Direct feedback is encouraged so that the information can be improved. A modified version of the database will be made available soon to other interested parties such as airlines, travel agencies, and businesses.

It is hoped that this use of modern technology will help the medical profession to minimise preventable illness in travellers and reduce the costs associated with travel related illnesses.

JONATHAN H COSSAR
ERIC WALKER
DANIEL REID
ROBERT D DEWAR

Ruchill Hospital,
Glasgow G20 9NB

- 1 Cossar JH, Reid D, Grist NR, et al. Illness associated with international travel: a ten year review. *Travel Medicine International* 1985;3:13-8.

- 2 Reid D, Cossar JH, Ako TI, Dewar RD. Do travel brochures give adequate advice on avoiding illness? *Br Med J* 1986;293:1472.
- 3 Walker E, Quayyum A. Malaria in Scotland and its presentation in a Glasgow hospital. *Health Bull (Edinb)* 1981;39:281-6.

Rare tumour system

SIR,—We are indebted to you for bringing to our attention the Leeds Oncology Information Centre (28 November, p 1421). This organisation receives and collates information on the management of rare cancers and facilitates exchange of information between oncologists. For some time in this oncology unit we have considered floating the idea of a national rare tumour register to gather as much data as possible about the diagnosis, management, and treatment of such uncommon neoplasms.

The situation that obtains at the moment is unsatisfactory. A clinician encountering a rare tumour must review the published reports, which is time consuming, and the desired information is often incomplete. Many of these reports are case reports, often published as curiosities to expand a curriculum vitae for professional advancement. Because of the nature of case reports details tend to be scanty and their value may be lost in the mountain of papers produced each year.

We would propose the establishment of a central rare tumour register which would receive fully detailed accounts of any rare tumour covering all aspects of the disease. These accounts would amount to a detailed discharge summary. On receipt and acceptance the doctor submitting the report would be credited with a rare tumour register citation number. This number would be acceptable for inclusion on a curriculum vitae as a publication. This system would encourage those in training to submit such reports and would avoid the tedium of revising and redrafting less complete case reports for publication in journals. Such a system would require the active support of bodies such as the BMA, the royal colleges, and the Department of Health and Social Security so that the rare tumour register citation attained some status. Once the central registry had accumulated sufficient cases we would expect that authoritative monographs could be produced and would furnish a very useful source of information for the clinician confronted with the management of unusual tumours.

J J BOLGER
D A WILLIAMS
J GOEPFEL

Weston Park Hospital,
Sheffield S10 2SJ

Intervention by general practitioners to reduce smoking

SIR,—Dr M A H Russell and colleagues (14 November, p 1240) addressed an important issue in their study of the effects of interventions by general practitioners to help people stop smoking. In summarising their findings they state that "brief intervention given without the support and back up of the local smokers' clinic had no detectable advantage over general practitioners' usual care." They also conclude that the supported brief intervention procedure "significantly increased the success rate." We believe, however, that methodological flaws in the study design cast doubt on these conclusions.

Though the authors report that there was no significant difference in either self reported or "adjusted" abstinence rates between the unsupported brief intervention and usual care groups, several design issues render this finding difficult to interpret. Firstly, the usual care that was provided was neither described nor monitored. It is quite

possible that the "brief intervention" was quite similar to, if not identical with, "usual care." In addition, the doctors who provided brief intervention in both experimental groups also provided "usual care" to patients randomised to this condition, greatly increasing the chance of contamination.

The authors also do not provide any description of the procedure used to train doctors in the specific interventions or any description of the method used to determine their competence in performing the interventions. We cannot assume that the doctors actually provided the brief interventions the authors described. Because of these methodological problems, this study cannot be cited as an adequate test of the effectiveness of brief interventions.

Secondly, the authors' finding that the supported brief intervention procedure "significantly increased the success rate" also has to be considered with caution. As the authors note, the self report data which produced the significant differences among groups is tainted by the high failure rate on urinary cotinine testing in smokers who claimed to have stopped smoking. Validation of outcome is further tainted by the low compliance with urine cotinine testing in the supported brief intervention group. Additionally, since the authors did not ask about the use of nicotine gum at follow up, the meaning of a "failed" urine cotinine test is unclear. As the authors themselves conclude, without appropriate validation of self reported smoking status, smoking outcome results are subject to question.

Several previous studies have shown the effectiveness of doctors' counselling efforts,^{1,2} though carefully controlled studies that document the method of physician training and physician compliance with the counselling intervention are also needed. The United States National Cancer Institute is presently supporting a number of such studies. Their results are eagerly awaited.

MICHAEL G GOLDSTEIN
RAYMOND NIAURA

Miriam Hospital,
Department of Psychiatry,
Providence, RI 02906,
USA

- 1 Russell MAH, Wilson C, Taylor C, Baker CD. Effect of general practitioners' advice against smoking. *Br Med J* 1979;ii:231-5.
- 2 Schwartz JL. *Review and evaluation of smoking cessation methods: the United States and Canada, 1978-1985*. Bethesda, Md: National Cancer Institute, US Department of Health and Human Services, 1987:50-9.

Cocktail stick injuries

SIR,—Retained cocktail sticks are found not only in wounds of the extremities (19-26 December, p 1658).

A previously fit 19 year old male student presented with a six week history of abdominal pain relieved by antacids, two weeks of right sided backache with malaise, and loss of appetite and weight. On examination of the abdomen there was slight right upper quadrant discomfort and the tip of the spleen was palpable. He was feverish (40°C) and had three splinter haemorrhages and a soft ejection systolic murmur. The white cell count was $9.7 \times 10^9/l$ (81% neutrophils) and the erythrocyte sedimentation rate 65 mm in the first hour. Blood cultures one and two days after admission yielded *Streptococcus milleri* group C. Echocardiography showed no abnormality. Abdominal ultrasound examination showed no hepatic or renal abnormality but an enlarged spleen, and in the absence of any other obvious focus of infection treatment for endocarditis was started, with 14

days of penicillin and gentamicin. Symptoms and fever resolved although the peripheral white cell count was $13.2 \times 10^9/l$ in the second week of treatment.

He was discharged home well taking oral amoxycillin but within five days was readmitted with increasingly severe right sided abdominal pain; at laparotomy a cocktail stick was found protruding through a hole in the second part of the duodenum into omentum. *Serratia marcescens* was isolated from the small amount of surrounding pus. The patient subsequently admitted that the offending foreign body had been ingested during New Year's Eve revelries some three months earlier—serving to remind us of the importance, in the investigation of febrile gastrointestinal illness, of the question, "And what have you been eating?"

W R GRANSDEN
SUSANNAH EYKYN

Department of Microbiology,
United Medical and Dental Schools
of Guy's and St Thomas's Hospitals,
St Thomas's Campus,
London SE1 7EH

Embryo research

SIR,—Professor R M Winston is entitled to his opinion (12 December, p 1501) but must it masquerade as fact in a leading article? It is paradoxical that it should occur during a series in the same journal expounding the use of logic.

Firstly, we see yet again an unfounded, unscientific assumption that there is something different about the preimplantation human embryo—something not human, even undifferentiated. To support this argument he, for some reason, mentions the use of abortive, blatant contraceptives, the use of which was foisted on the public long before their real mode of action became known. He goes on to say that many patients would benefit from the detection of genetic defects. His examples—Lesch-Nyhan syndrome, Tay-Sachs disease in Ashkenazi Jews—actually leave one with the opposite impression—that is, that only a tiny minority of pregnant women would benefit and only those who had risk factors beforehand.

Next, Professor Winston justifies embryo experimentation in that it leads to the birth of healthy babies by in vitro fertilisation and, even more incredibly, might help to prevent miscarriages. Two thousand healthy babies by in vitro fertilisation is a drop in the lake of potentially healthy babies aborted, and a drop in the ocean if you add those prevented by contraceptives that prevent implantation. We are left with the idea that a decision banning embryo research imposed by a democratically elected government on unelected research workers, who are at best unrepresentative of the general population of Britain, is unjust. A strange meaning of the word "justice."

If the state banned embryo research it could at least claim to be representing a majority—not, as Professor Winston puts it, "the sincerely held views of a minority."

PAUL MALLETT

St Johns Wing,
Homerton Hospital,
London E9 6SR

AUTHOR'S REPLY—My leading article is, of course, perfectly appropriate as such because it reflects an important body of medical opinion, including that of the BMA.

Nor is it "unscientific." The contention that the preimplantation embryo is different from an implanted fetus is founded on biological facts. Before implantation it is indeed totally undiffer-

entiated with the potential only for becoming a baby. The evidence is that at least 40% (and possibly most) of human preimplantation embryos are lost by the time of the first menstrual period. The "human-ness" of the preimplantation embryo is something we can argue about, but are not sperm and eggs from our species (which are unprotected by law) equally human?

Mr Mallett, unlike most people, opposes blatant contraceptives; perhaps this explains his obviously deeply felt views. Apparently the low dosage pill and the intrauterine contraceptive device were "foisted on the public." Surveys in Hammersmith's family planning clinic confirm that few people worry about the mechanism of action of these contraceptives and that, after thorough explanation of how the device works, not one of our patients has discontinued this contraception for those reasons. Hardly "foisting," Mr Mallett.

There is now no doubt that embryo research will combat severe congenital disease. Of course, Tay-Sachs disease and Lesch-Nyhan syndrome are rare. Since when has affliction with a rare disease meant that that disease is unimportant to patients, their families, or their doctors? But Mr Mallett is typically selective in ignoring the article's discussion of common fatal diseases such as muscular dystrophy and cystic fibrosis—a defective gene carried by one in 20 of the population.

No, Mr Mallett, legislation banning this work would not represent the views of a majority. Most people in Britain support embryo research. In the past two years three independent national opinion polls—and four surveys by responsible television and radio programmes, too—have increasingly confirmed that the majority approve this research. But, more importantly, in a free democracy those suffering the appalling consequences of severe genetic disease and sad effects of infertility or repeated miscarriage should be allowed the benefit of research which they almost unanimously support and which is voluntarily accomplished solely as a result of their own treatment by in vitro fertilisation.

R M L WINSTON

Institute of Obstetrics and Gynaecology,
Hammersmith Hospital,
London W12 0HS

Light on seasonal affective disorders?

SIR,—Since Drs Melanie Abas and Declan Murphy (12 December, p 1504) admit that the mode of action of light treatment in seasonal affective disorder is unknown and that the extent (if any) of a placebo effect needs establishing one wonders what has been achieved in this subject.

They fail to comment on the possibility of deleterious side effects of intense photic radiations. For example, the work by Lewy *et al* which they quote involved daily two hourly exposures, for possibly 28 days,¹ to a luminance level of about half the threshold intensity producing photoreceptor damage in macaques in a single 12 hour session.² The traditional safety factor of 10 was not considered.

There may be important differences in safe exposures between phakic and aphakic patients because of the protection that the normal crystalline lens may provide for the retina.³ Since the density of the macular pigment appears to be correlated, if only on the basis of a cluster, with the presence of red hair, a further undocumented variable may have been included. Even the possible difference in pupillary diameter—for example, between blue and other eyes—ought to be considered.

As the difficulties of reliably determined retinal illumination (and hence the alleged therapeutic