- 14 Knoch M, Müller E-E, Höltermann W, Konder H, Lennartz H. Erfahrungen mit der extrakorporalen CO₂-Elimination. *Anaesthesist* 1987;36:210-6.
- 15 Gattinoni L, Agostini A, Damia G, et al. Hemodynamics and renal function during low frequency positive pressure ventilation with extracorporeal CO₂ removal. A comparison with continuous positive pressure ventilation. *Intensive Care Med* 1980;6:155-61.
- 16 Pesenti A, Kolobow T, Buckhold DK, Pierce JE, Huang H, Chen V. Prevention of hyaline membrane disease in premature lambs by apnoeic oxygenation and extracorporeal carbon dioxide removal. *Intensive Care Med* 1982;8:11-7.
- 17 Kolobow T, Moretti MP, Mascheroni D, et al. Experimental meconium aspiration syndrome in the preterm fetal lamb: successful treatment using the extracorporeal artificial lung. *Trans Am Soc Artif Intern Organs* 1983;29:221-5.
- 18 Dorrington KL, McRae K, Bellhouse BJ. Extracorporeal CO₂ removal (ECCO₂R) improves oxygenation in a lung lavage model of acute respiratory failure. *Trans Am Soc Artif Intern Organs* 1987;33 (in press).
- 19 Kolton M, Cattran CB, Kent G, Volgyesi G, Froese AB, Bryan AC. Oxygenation during high frequency ventilation compared with conventional mechanical ventilation in two models of lung injury. *Anesthesia and Analgesia* 1982;61:323-32.
- 20 Matamis D, Lemaire F, Harf A, Brun-Buisson C, Ansquer JC, Atlan G. Total respiratory pressure-volume curves in the adult respiratory distress syndrome. *Chest* 1984;86:58-66.
- 21 Hamilton PP, Onayemi A, Smyth JA, et al. Comparison of conventional and high-frequency ventilation: oxygenation and lung pathology. *J Appl Physiol* 1983;55:131-8.
- 22 Froese AB, Bryan AC. High frequency ventilation. *Am Rev Respir Dis* 1987;135:1363-74.
- 23 Gattinoni L, Pesenti A, Caspani ML, et al. The role of total static lung compliance in the management of severe ARDS unresponsive to conventional treatment. *Intensive Care Med* 1984;10:121-6.
- 24 Cornish JD, Gerstmann DR, Clark RH, Carter JM, Null D, deLemos RA. Extracorporeal membrane oxygenation and high frequency oscillatory ventilation: potential therapeutic relationships. *Crit Care Med* 1987;15:831-4.

The private hospital and the surgical trainee

Private practice is expanding rapidly in Britain. About one person in 10 belongs to a private insurance scheme, and new and increasingly luxurious private hospitals are appearing throughout the land. Some of these private hospitals will need resident staff. Young graduates may be attracted by these posts, which often provide considerable creature comforts, a good salary, and ample spare time in which to study towards higher examinations. But can such posts ever become a part of the training programme for young surgeons?

The requirements for recognising training posts in surgery (including all its specialties) are well defined by hospital recognition committees of the royal colleges and include close and personal interest by the one or two consultants on the "firm" in training the junior doctor; a responsible college tutor in the hospital; access to a wide variety of clinical experience, which should include dealing with emergency admissions; an active accident and emergency department; personal responsibility for patients; facilities for outpatient experience (including the clerking of new patients); and—most important of all—the chance to operate under the supervision of seniors. In addition, there should be excellent parasurgical facilities, including pathology services, the radiology department, and an intensive care unit; and these departments should actively teach junior staff. There should be a regular teaching programme, which should include seminars, tutorials, lectures, and, in particular, "death and disaster" meetings that regularly monitor the unit's activities. Also necessary are an efficient medical records department, a good library, and adequate provision for study leave.

Of course, many training posts in the National Health Service fall far short of these ideal requirements (and those in teaching hospitals are no exception), but the regional advisers, the college tutors, and the college inspectors struggle valiantly to achieve these ideals. Posts recognised as suitable for training for the examination to become a fellow of the Royal College of Surgeons are inspected every five years, and recognition for training can be withdrawn.

Some private hospitals may provide all these facilities, but

I doubt it. Does any post in the private sector include a one to one or one to two ratio between the trainee and consultant? Usually the one or two residents in the private hospital will be responsible to all the consultants using the hospital facilities. How many emergencies are juniors likely to see? And will they actually have hands on experience with many of them? Above all, will they get operative experience under supervision? This is surely the single most important requirement in a surgical training programme, yet are not many people choosing private care in order to ensure that their operations are done by the consultant surgeons of their choice?

These are difficulties that could be surmounted. Particularly in the specialties, a training programme could be planned in which juniors rotate between employment on specialised units in NHS hospitals and working for the same consultant teams in the private sector, where personal tuition could be continued. If a worthwhile scheme were to be put forward, including time in a private hospital, the colleges would need to inspect and consider it on its merits, but there seems no reason why this should not occasionally be acceptable.

Today resident posts in private hospitals provide a comfortable billet and pay while the young graduate raises funds, works for an examination, or simply waits for a job in the NHS. The future may bring some of these jobs into line with training posts in the NHS, and there then seems little practical (as compared with ideological) reason why they should not be recognised as counting towards surgical training.

HAROLD ELLIS

Professor of Surgery,
Charing Cross and Westminster Medical School,
London SW1P 2AP

Medical marriage: no bed of roses

Medical careers seem both absorbing and stressful to doctors themselves and to those who come into contact with them—from the patient who receives daytime care from the same junior doctor who admitted him in the small hours to the potential spouse attempting to coordinate time for courtship. How stressful is medicine, is the stress confined to the early postgraduate years, and what effects do the stress and unusual lifestyle have on the quality of marriage and family life in the longer term?

Many studies, mostly from the United States, highlight the stressful nature of medicine, the high suicide rate, particularly among women doctors, the use and abuse of drugs and alcohol, the frequency of psychiatric treatment, and the adverse effects on personal life.^{1,4} Many doctors are predisposed to emotional difficulties from childhood, but they often suffer severe emotional distress during the first postgraduate year: half of preregistration house officers showed emotional disturbance and 28% depression in a recent study.⁵ The stress diminishes with seniority, particularly in clinical rather than academic settings, and there is increasing job satisfaction.³

Doctors mostly seem to suffer unsatisfying but enduring marriages compared with other professions.^{1,2,6} In a recent study of the marriages of a group of middle aged American general practitioners and non-academic specialists, almost all

of whom were men, the greatest source of conflict perceived by both partners was "lack of time for self, family, and fun."⁷ The men blamed time spent on work, which averaged 56 hours a week, while their wives were primarily concerned about lack of intimacy. The doctors blamed their spouses for lack of sexual interest or sympathy for their job while their partners felt they did not talk to them enough or provide emotional support. Ironically, both complained that their partners did not listen, and satisfaction in marriage was clearly related to the time a couple spent talking to each other. Nearly half had sought marital counselling and a further fifth had considered it; these tended to be the couples who talked less to each other. In contrast to the findings of other studies, these couples, who had chosen to attend a re-education programme together, expressed high marital satisfaction overall.

Neither dissatisfaction in marriage nor emotional distress in house officers can be directly related to the numbers of hours worked,^{5,7} yet the houseman's main complaint is overwork and the primary source of marital conflict perceived by both partners is lack of time. How is it that traditional notions about the adverse effects of long hours are not supported by research? Firstly, intense preoccupation with work may be more harmful to people and their marriages. Secondly, doctors' hours are always long early in their careers, and time alone does not reflect the strain of disrupted sleep, study for postgraduate examinations, difficult staff relationships, or the pressure and responsibilities of certain specialties. Finally, there is self selection in the choice of specialties and on call rotas: women doctors tend to choose specialties they consider more compatible with family life rather than their preferred specialty, while men put their career choice first.⁸ Both men and women doctors marry early in their careers, and most women doctors, particularly in hospital medicine, now work full time. Within 10 years of qualification 85% of married female graduates had children.⁹

Little is known about the quality of recent medical marriages other than that doctors are increasingly marrying each other. These doctors experience the complications of hours on call, job moves, house moves, sharing housework, and childcare while trying to find time for one another and some social life. Couples and their marriages can survive these early years, even when both work, if they respect each other's careers and are both prepared to make some sacrifices, provide mutual emotional support, and devote much of their precious free time to talking. Otherwise, maladaptive patterns may set in with one or both partners withdrawing for protection, gratifying many needs at work until they discover their needs for intimacy are unmet; men tend to express a wish for greater physical intimacy and women greater verbal intimacy.^{2,7} Timing of starting a family is important, particularly in doctor-doctor marriages, and delay may allow for the more stressful years to be over; housework and childcare can then be afforded, and the couple can devote maximum amounts of free time for each other and their family. Little is known about the effects of medical marriages on the children other than that they tend to become doctors themselves. Certainly, if childcare is of high quality and consistent they need not suffer.

ALYSON HALL

Consultant Child Psychiatrist,
Emanuel Miller Centre for Families and Children,
London E14 8HQ

1 Vaillant GE, Sobowale NC, McArthur C. Some psychological vulnerabilities of physicians. *N Engl J Med* 1972;287:372-5.

2 Gerber A. *Married to their careers: career and family dilemmas in doctors' lives*. New York, Tavistock, 1983.

- 3 Mawardi BH. Satisfaction, dissatisfactions and causes of stress in medical practice. *JAMA* 1979;241:1483-6.
- 4 Ross M. Suicide among physicians. *Dis Nerv Syst* 1973;34:145-50.
- 5 Firth-Cozens J. Emotional distress in junior house officers. *Br Med J* 1987;295:533-6.
- 6 Garvey M, Tuason VB. Physician marriages. *J Clin Psychiatry* 1979;40:129-31.
- 7 Gabbard GB, Menninger MD, Coyne L. Sources of conflict in the medical marriage. *Am J Psychiatry* 1987;144:567-72.
- 8 Hutt R, Parsons D, Pearson R. The timing of and reasons for doctors' career decisions. *Health Trends* 1981;13:17-20.
- 9 Stephen PJ. Career patterns of women medical graduates 1974-84. *Med Educ* 1987;21:255-9.

Spout or pouch? Alternatives for patients with ileostomies

Advances in technology and ready access to a stoma therapy adviser have made life almost normal for many people with ileostomies.¹ None the less, there is a psychological and social cost to be borne,² and not surprisingly when a surgical alternative to the conventional spouted ileostomy becomes available patients are prepared to risk the dangers of a new procedure.

Until recently the best operation for patients with ulcerative colitis requiring surgery would have been a proctocolectomy since it removes all the diseased colon and any risk of cancer in one operation and is usually uncomplicated. A few patients with little disease in the rectum may be suitable for an ileorectal anastomosis. Though this avoids a stoma, about a third of patients eventually have to have an ileostomy.³ It does, however, allow suitable younger patients to mature through adolescence without a stoma. The disadvantage is a risk of malignant change in 3-5%, and since tumours arising in the retained rectum may be difficult to recognise follow up must be scrupulous. The stepping stone on the way to a procedure that does not necessitate an ileostomy was the Kock continent ileostomy, in which a reservoir was constructed from the ileum.⁴ Patients had no appliance but still had to intubate the reservoir intermittently for emptying. This operation, though ingenious, has now been superseded by the pelvic ileal reservoir or pouch procedure.

Joining the ileum to the anus was not a new idea, but putting a reservoir in the pelvis was. Ten years ago Parks *et al* and others recognised that the anal sphincter could be preserved in patients with ulcerative colitis, provided that the diseased mucosa in the uppermost anal canal and lower rectum was removed.^{5,6} The ileal reservoir, now of various designs (fig 1), is sewn on to the sphincter inside the anus, and the anastomosis protected temporarily by a loop ileostomy that is closed about eight weeks later (fig 2).⁶⁻⁸ The exact level of the anastomosis is controversial,^{9,10} but the theoretical danger of leaving behind islands of diseased rectal mucosa is unfounded.¹¹

Although there are no reports of postoperative deaths, patients have suffered formidable upsets during development of the procedure. Obstruction by small bowel adhesions is common, but the most serious complication is pelvic sepsis, usually around the anastomosis. It may be difficult to eradicate since it cannot drain freely, and healing with fibrosis may spoil the eventual functional result. None the less, over three quarters of patients will not have a complication. When the protective ileostomy is closed between 5% and 15% of patients will have minor discharge or soiling, especially at night, which is probably caused by an injury to the internal sphincter.¹² Frank incontinence is unusual, and function improves over the first 12 months. A stool frequency of three to six times a day without urgency can be expected,