therapy. On the other hand, treatment may be more effective for several reasons. Cryotherapy, for example, is most effective if it is given every three weeks,⁶ but owing to other demands on dermatologists' time this is not always possible. This causes the treatment course to be longer than necessary, thus compounding the original problem of insufficient available time. In a separate clinic run by a nurse it should be possible to provide more regular treatment.

Caution on the part of nurses may also mean that there would be fewer side effects from treatment. On the other hand, routine referral of all patients to a wart clinic may lead to more patients receiving cryotherapy because of its convenience to the operator, and this may result in more side effects. Medicolegal problems may occur for this reason.

In the end value judgments have to be made. We think that the potential advantages of delegating the routine treatment of warts to nurses outweigh the potential disadvantages, and we therefore propose a system for providing wart treatment.

Proposed system for providing wart treatment

All routine wart treatments should be provided by nurses or general practitioners.

Patients with atypical lesions should be referred to hospital for diagnosis, particularly when sites other than the hands, face, or feet are affected or the patient is over the age of 25. If the dermatologist agrees that the lesion is a wart then the patient can be referred back to the nurse or general practitioner for further treatment. The exception to this is that genital warts should be referred to departments of genitourinary medicine.

Patients whose warts fail to respond to six months of conventional treatment could be referred to the dermatologist so that other measures may be considered. The dermatologist may decide that

more prolonged treatment with conventional methods is appropriate and refer the patient back.

Conclusion

Despite our knowledge of the natural history of warts it seems unlikely that the patients will ever be convinced that no treatment is the best treatment except in very young children. There is no likelihood that the demand for treatment will go away, particularly in a society that is being encouraged to be conscious of personal appearance. The present system has been allowed to evolve to the detriment of the service that dermatologists provide to patients with skin disease. Dermatologists, general practitioners, nurses, and managers should take joint responsibility for implementing a policy that would lead to a far more sensible use of resources.

We thank Mrs N Dale, Mrs E Williams, and the medical records staff at Falkirk and District Royal Infirmary for their help. We also thank our colleagues for their cooperation and comments.

References

- 1 Lyell A. Management of warts. *Br Med J* 1966;iii:1576-9.
- 2 Rook A, Savin JA, Wilkinson DS. The prevalence, incidence and ecology of skin disease. In: Rook A, Wilkinson DS, Ebling FJG, Champion RH, Burton JL, eds. *Textbook of dermatology*. 4th ed. Vol 1. Oxford: Blackwell Scientific, 1986.
- 3 Benton EC, Hunter JAA. The dermatology outpatient service: a study of out-patient referrals in a Scottish population. *Br J Dermatol* 1984;110:195-201.
- 4 Royal College of General Practitioners. *Morbidity statistics from general practice. Third national study 1981-82*. London: HMSO, 1986.
- 5 Bunney MH, Nolan MW, Buxton PK, Going SM, Prescott R. The treatment of resistant warts with intralesional bleomycin: a controlled clinical trial. *Br J Dermatol* 1984;111:197-207.
- 6 Bunney MH, Nolan WN, Williams DA. An assessment of methods of treating viral warts by comparative treatment trials based on a standard design. *Br J Dermatol* 1976;94:667-79.

(Accepted 25 August 1987)

My Student Elective

Witchdoctors in Africa

A B KOZIELL, I F LAURENSEN

Baked insects, dried reptiles, and shrivelled white man's flesh are not accepted methods of treatment in most modern textbooks of medicine. In Zululand, however, many patients coming to hospital may already have been treated with similarly bizarre remedies. There is a deeprooted conviction among many tribespeople that a witchdoctor must be visited before a visit to the hospital is considered. It is not unusual to have to treat a sick Zulu first for the effects of a witchdoctor's potions and only subsequently for the original complaint. This sort of conviction emphasises the nature of

many problems we faced working as "modern medicine men" in an environment of strong tribal custom and mysterious beliefs.

We spent our electives at Charles Johnson Memorial Hospital in Nqutu, Kwazulu; it is a former mission hospital now under the jurisdiction of the Kwazulu government. The problems that we encountered are characteristic of those in other black rural homeland hospitals and contrast greatly with those in white hospitals.

The setting

Lying in the heart of Zululand, Charles Johnson Memorial Hospital serves a population of about 500 000 of South Africa's five million Zulu, the largest tribal group in the country. The Christian influence still prevails as witnessed by the centrally situated church and the prayers sung in harmony by the nurses at the change of each shift. The 600 or so beds and assorted floor space are divided among adults (anyone aged over 10), obstetrics, paediatrics, and psychiatry (nicknamed "the zoo" by the locals). The staff comprises Zulu

Department of Medicine, Queen Mary's Hospital, Sidcup, Kent
A B KOZIELL, BSC, MB, house physician

Roodlands General Hospital, Haddington, East Lothian EH4 3PF
I F LAURENSEN, MA, MB, house surgeon

Correspondence to: Dr Koziell.

Map of eastern South Africa, showing position of Nqutu and Zulu area (shaded).

nurses and fairly junior (mainly white) doctors aided by miscellaneous medical students. Nurses take on responsibilities not usually demanded in the United Kingdom, including interpreting for doctors and running the psychiatric unit and outlying clinics. Admissions are made from the outpatient/casualty department or the clinics. The clinics take the place of general practitioners, although general practitioners do exist in more urban areas. The clinics are visited fortnightly by doctors from the hospital.

The illness seen ranges from classic "Third World" diseases such as tuberculosis and those stemming from malnourishment to those found in developed countries: diabetes, congestive cardiomyopathy, and hypertension are particularly common. Trauma and burns contribute considerably to the workload, especially when weddings, funerals, or pay day are combined with home brewed Zulu beer. Thankfully, we saw only two cases of "necklace" burns, which are caused by a tyre filled with burning petrol placed around the necks of supposed traitors; in more troubled parts such cases are more common.

Interaction between Zulu and Western medicine

Many Zulu believe that some illnesses occur only among African peoples and so are amenable only to traditional treatment. Nevertheless, consultations with the isangoma, or witchdoctor, were expensive, costing £5.10 or more, whereas a hospital consultation would cost 50p and admission £1.00. An average family income in the area was roughly £66 a month. The isangoma is a powerful and important person in Zulu life; he is a diviner and doctor, acting as a link between spirits and their descendants, a protector of society. He "smells out" evil men affecting their neighbours and bringing disaster to the community. Interestingly, in such a male dominated society, many isangomas are women.

The isangoma is still regarded by many Zulu as superhuman and capable of striking them dead on the spot. Anything stolen from the hospital would reappear with miraculous rapidity if word was put

around that the isangoma would be asked to find the culprit. Although in this case belief in the isangoma could be put to good use, it was much more often the source of great frustration among the medical staff. Fear of incurring the isangoma's wrath led many Zulus to consult him before a hospital doctor. In addition, it was considered extremely unlucky to mix the two types of medicine, whether or not the isangoma had effected a cure. Against such a background patients often presented with a variety of bizarre complaints, not infrequently exacerbated by the ministrations of an isangoma. Differences in attitudes among the doctor, isangoma, and patient as to the cause and appropriate treatment of an illness meant that patients were sometimes terminally ill by the time they came to hospital.

The isangoma lives in a mud and grass hut similar to that of his Zulu patients, except that it is usually larger and has a flagpole outside. Each flag represents a miracle that the isangoma has performed, but we never spotted more than three together. The isangoma's knowledge of disease is limited, and his acquaintance with the structure of the human body is drawn mainly from analogy with the anatomy of animals. For an isangoma the symptoms of a disease are the disease, and so the great rule is: "As many symptoms, so many diseases." A person with acute cholecystitis may have pain radiating to the angle of the scapula. This pain would be regarded as a separate complaint and called "isibindi," while any symptoms in the region of the liver would be called "isibhobo." The strategy of the isangoma would be to treat each symptom vigorously but individually, without any thought as to what the underlying disease process may be.

The role of isangoma is assumed by inheritance and apprenticeship. The isangoma will ask questions of bone or stick patterns or his thumb while "possessed by spirits." These give answers, which allow the cause of the illness to be discovered and corrective measures prescribed. Illness is often thought to be caused by the black arts and the machinations of a wizard, which must therefore be counteracted by magic. The isangoma's medicines are classified according to use. "Black medicines" causing emesis are seen as working by washing the body, consuming badness in the stomach. "White medicines" may be rubbed into the body. Other medicines may be scattered around a hut or homestead. Enemas are the isangoma's favourite method of administering his medicines. These cause severe dehydration, especially in children, often complicated by hypoglycaemia. In addition, charms of various descriptions are worn to ward off evil spirits.

Marking the body

Health giving spells may be cast by marking the body with charcoal or razor blades. The blood from the incisions may be drained by suction into a cow's horn and ashes or boiled bark juice

Typical Zulu huts near the hospital.

rubbed into the wounds. Skin marking was commonly seen on patients with congestive cardiac failure. Small cuts were made over the medial and lateral aspects of the legs and by the big toes, presumably so that the oedema fluid could run out. Marking was also used for pneumonias and other chest complaints, a circular pattern of cuts being made over the chest wall with a razor blade. The fascinating aspect of these circles was that they always corresponded to the areas of maximal bronchial breathing. Apparently the isangoma listens to the chest wall and marks the areas of altered breath sounds as this is where the evil spirit is to be found.

Children did not escape marking; most seen at the hospital had been taken to an isangoma at least once. In most cases it was just to have small scratches made around the umbilicus to ward off evil spirits. If the child was ill and the marks were fresh, however, we could be fairly certain that the isangoma had administered Zulu medicines; this often complicated the clinical picture considerably.

The isangoma also plays an important part in questions of love, sex, and sexually transmitted disease. The cure for syphilis is to place a live frog against the affected part and then throw it into running water. The isangoma's advice may also be sought in cases of adultery; a husband who suspects his wife of having a lover eats special foods recommended by the isangoma. These are supposed to transmit disease or even death through him to his wife and thus to the illicit partner. If a husband commits adultery, however, it is regarded simply as high spirits. Male impotence is said to be best treated by an emetic in goats' milk.

Sky doctors

Lightning and its causes are important to the Zulu as evil spirits are thought to manifest themselves through its actions. This is perhaps not surprising as severe electrical storms are common in Zululand. Another type of doctor, the sky herd or heaven doctor, has a role in protecting people from thunder, lightning, and hail. Sky herds' power is derived from magic. During a storm they run out with rainshields and weapons shouting at the lightning, telling it to go away. For a fee they actually protect huts from lightning, while if a homestead is struck by lightning they must come to minister magic before eating and drinking may occur again. Such is the superstition attached to lightning that the occupants may decide to move to another place if this is recommended by the heaven doctor. After storms people regularly came to the hospital complaining of having been struck by lightning. Examination was generally unremarkable, and a disgruntled person left the hospital; "de-electrification" by therapeutic electrocardiography was often curative and satisfying in such cases. White man's magic?

Despite such apparent empiricism occasionally the isangoma's

Zulu women dancing to Christmas carols played on bagpipes.

cure was successful. This was usually when psychiatric, moral, or social problems were prominent. An interesting example was the case of a woman with severe puerperal psychosis that had proved totally refractory to treatment. Unusually, a female isangoma had decided to deliver her baby in hospital, and so as a last resort she was asked to talk to the woman. Within three hours the psychosis was cured: obviously, the isangoma was competent to deal with an aspect of Zulu psychology that we knew nothing about.

This article illustrates the types of experience we had at the Charles Johnson Memorial Hospital. At any such hospital the personalities of other staff are of paramount importance in influencing one's experiences and memories. Language and culture may readily obscure and frustrate in such a different environment, and attitudes, flexibility, and past experiences determine one's reactions. We would like to thank those whose advice and support made our stay worth while and possible.

IFL was financially supported for part of his elective by the Medical Research Council; Milupa Ltd; and Girton College, Cambridge.

Bibliography

- Bryant AT. *Zulu medicine and medicine men*. Cape Town: Centaur, 1983.
- Bryant AT. *The Zulu people*. Natal: Witness, 1949.
- Kriege EJ. *The social system of the Zulus*. Cape Town: Longmans Green, 1939.
- Ngubane H. *Body and mind in Zulu medicine*. London: Academic Press, 1977.

(Accepted 21 July 1987)

An 80 year old woman recovered from a cerebrovascular accident eight years ago, but now has distressing urge incontinence, presumably because of detrusor weakness. What treatment is advised?

The term "detrusor weakness" is somewhat misleading. The patient's symptoms could be due to detrusor underactivity or detrusor overactivity (instability). Either could cause the symptoms of urge incontinence—the first through inefficient bladder emptying, resulting in a considerable volume of residual urine and leakage due to overflow with an increased risk of urinary tract infection. The second problem, detrusor instability, is associated commonly with a cerebrovascular accident. Urinary control may have been lost after eight years because of the reduction in resistance offered by the urethral sphincter mechanism and periurethral supports associated with increasing age.

It is important to exclude a urinary tract infection, carcinoma of the bladder, vesical calculi, and interstitial cystitis. Postmenopausal oestrogen deficiency may be detected by atrophic changes in the vagina, which are mirrored in the urethra. Cystoscopy may be necessary to exclude major disease, but in the absence of haematuria this is unlikely. Simply passing a catheter immediately after voiding will assess the volume of residual urine.

Urodynamic studies are well tolerated by women of all ages and would be the only way to detect detrusor instability.

If the detrusor is underactive and the symptoms a result of incomplete bladder emptying clean intermittent catheterisation by the patient or carer should be considered.¹ This will reduce the frequency of voiding, episodes of incontinence, and the risk of infection. Atrophic changes should be treated by a short course of hormone replacement given orally or by vaginal applicator.² Detrusor instability responds well to bladder training and treatment with anticholinergics—for example, terodiline.³ Much helpful advice and support can be gained from the district continence adviser and her team of resource nurses in hospital and community. There are over 100 continence advisers in post throughout the United Kingdom, and all have a comprehensive knowledge of the methods of maintaining continence and managing incontinence.—J C GINGELL, consultant urologist, A M SHEPHERD, associate specialist in urogynaecology, Bristol.

- 1 Murray K, Lewis P, Blannin J, Shepherd A. Clean intermittent self-catheterisation in the management of adult lower urinary tract dysfunction. *Br J Urol* 1984;55:379-80.
- 2 Walter S, Wolff H, Barlebo H, Jensen HK. Urinary incontinence in post-menopausal women treated with oestrogens: a double blind clinical trial. *Urol Int* 1978;33:135-43.
- 3 Tapp A, Fall M, Norgaard J, et al. A dose-titrated, multi-centre study of terodiline in the treatment of detrusor instability. *Neurology and Urodynamics* 1987;6:254-5.