Doppler studies in diabetic pregnancy

SIR,—Dr R J Bradley and others (9 January, p 94) expertly demonstrated the combined approach of Doppler ultrasound and transabdominal cordocentesis in complicated pregnancy. I would caution them, however, in their conclusion that hypoxia with concomitant growth retardation marked by macrosomic effects of diabetes is the likely cause of the “unexplained” diabetic stillbirth, and that this sad event may be predicted by umbilical artery Doppler monitoring.

We have been monitoring our diabetic pregnancies for some time using Doppler ultrasound and have recorded one “unexplained” stillbirth in a poorly controlled diabetic. This fetus had entirely normal Doppler waveforms in the umbilical artery (A:B ratio 2.34 at 35 weeks) just three days before fetal death; the pregnancy was not complicated by hypertension. Bracero and others¹ have also reported one case of “unexplained” stillbirth in their series of 43 diabetic women, where an A:B ratio of 3.7 at 35 weeks (mildly abnormal) was found a few days before stillbirth. End diastolic flow was definitely present in both of these cases. Undoubtedly, poor placental perfusion, indicated by abnormal Doppler studies, will predispose to hypoxia and acidosis in the fetus, but this may be a result of hypertension, not poorly controlled diabetes. In the absence of other cases this report should remain interesting but not be grounds for complacency in the management of diabetic pregnancy characterised by normal Doppler studies.

S N TYRRELL

Department of Obstetrics and Gynaecology,
St James's University,
Leeds LS9 7TF

1 Bracero L, Schulman H, Fleischer A, *et al.* Umbilical artery velocimetry in diabetes and pregnancy. *Obstet Gynecol* 1986;68:654-8.

Obscure gastrointestinal bleeding

SIR,—The leading article by Mr A C Steger and Mr J Spencer (2 January, p 3) fails to distinguish between patients who suffer a single episode of acute gastrointestinal bleeding and those readmitted repeatedly with blood loss. In our experience the latter group can prove the more elusive because bleeding has often slowed or stopped before investigation can reveal its source. Arteriography is useful only when the rate of blood loss is greater than 1 ml/min during the investigation. When bleeding is slower we have found that radionuclide scanning with ^{99m}Tc labelled autogenous red cells is valuable for showing the general area of blood loss by accumulation of radioactivity.

Laparotomy alone may not show the source of obscure blood loss because the cause is often a mucosal lesion which is impalpable and invisible from outside the bowel. Within the past year we have diagnosed and treated three cases of acute recurrent gastrointestinal blood loss with intraoperative endoscopy. Upper gastrointestinal and colonic causes were excluded by endoscopy beforehand, and in each case the cause proved to be small bowel angiodysplasia. At laparotomy a sterilised colonoscope was passed into the small bowel via an enterostomy and then along its entire length. An “unscrubbed” operator handled the head of the instrument and carried out the inspection while the surgeon sleeved the bowel over the endoscope. The bowel wall was transilluminated (figure), and any angiodysplasias thus seen were underlaid with sutures from outside the bowel wall or else from within after opening

Transillumination of bowel wall at intraoperative endoscopy.

the bowel. One patient had three lesions in the distal duodenum, another had two lesions in the jejunum, and the third, a man of 79, had more than 30 lesions distributed throughout the small bowel.

We would recommend that intraoperative endoscopy should be available whenever a patient is undergoing laparotomy for obscure gastrointestinal blood loss.

C R G QUICK
R J DICKINSON

Hinchingsbrook Hospital,
Huntingdon, Cambridgeshire PE18 8NT

The NHS debate

SIR,—Sonia Alexander's letter (16 January, p 209) written in reply to Dr Tony Smith's new year message (2 January, p 1) causes me concern and dismay—words used by her to describe her reaction to Dr Smith's editorial.

To suggest that “the whole tenor of his article is founded on the premise that there is not a lot wrong with the NHS that a few thousand million pounds a year extra cash cannot put right” is a travesty of his well reasoned leading article. But Sonia Alexander is correct when she says that the root issue is philosophical and is concerned with the proper role of the NHS in Britain as we enter the 21st century: is the concept of health care for all, available at no cost to the patient at the time of use, a reasonable social and economic proposition? I prefer to use the phrase “available at no cost to the patient at the time of use” to her pejorative phrase “available freely on demand.”

While I accept that a service must be efficient, cost effective, and well managed and have the vested interests of all who work in the NHS under control, such measures are peripheral to the main philosophical debate. Whatever type of service is organised such principles are obviously important and must be dealt with. But what is central is to decide on which side of the debate one lies. Are we committed to the original aims and concepts of the NHS or not? Since 1948 there has been a consensus, held by all political parties, on the aims of the NHS.

But it would seem from Sonia Alexander's letter and from the attitude of the present government to the health service that they are closely allied to the views of the right wing of the Republican party in the United States of America, where health care is seen essentially as a marketplace commodity. Over

the past eight years the commitment to the original priorities of the NHS has decreased, and this must now be explained to the country openly and honestly. If Sonia Alexander represents the view of most of the chairmen of health authorities then the outlook for patients is indeed grim. Perhaps before she castigates people for intellectual bankruptcy and for jumping on a dissidents' bandwagon she should ask herself: Is she in favour of at least trying to provide an equitable comprehensive service to the poor and increasing percentage of the population with low wages, as well as to people like herself and myself? Perhaps she and her followers should also be reminded of the truism that taxes (which none of us like paying) are the price we pay for living in a civilised society.

W J STEPHEN

Wells,
Somerset

SIR,—On 9 January nearly 100 consultants met in Birmingham to discuss the funding crisis facing the National Health Service. It was a unanimous decision to form a group that would demonstrate the commitment of consultants to the concept of the NHS at a time when its survival is in doubt because of the absence of a national policy on health and because of measurable underfunding. Britain has a policy on economic growth and the creation of wealth, but we do not have a policy on health—the effective treatment of disease and the prevention of disease combined with the promotion of health.

Birmingham Consultants for Rescue of the NHS is planning to produce regular evidence about the lack of resources for services to patients in our five districts. We will show the efficiency savings that have been made and show that without more resources services inevitably will be further affected. We believe that doctors should put forward the facts because we appreciate the full implications, and, provided the data are presented responsibly, reasonable men and women will recognise the truth when they see it.

Birmingham is fighting its corner, but the whole country is similarly affected. We hope more cities and districts will follow our example. We think the general public needs to start its own campaign. The National Health Service faces a great threat to its future, and if people want it they must stand up and say so by every legitimate and democratic means. Strikes and working to rule solve nothing and damage the future of the NHS, although the frustration and despair which inspire that action are understandable.

We realise the importance of independence. We are not a trade union, and quality of service to patients is our prime objective. Most doctors take up the subject because it is a vocation, and consultants have been in training for as long as 15 years from entering medical school. Most wish to see their skills deployed to maximum benefit for their patients.

K G TAYLOR
Chairman

The Action Group,
Birmingham Consultants for Rescue of the NHS,
Birmingham B15 3AJ

Medicine and politics

SIR,—History is always fascinating, and even more so when the historian tells us of his views and actions at the time of events.

It is therefore disappointing that Sir Douglas Black in his Henry Cohen History of Medicine Lecture (2 January, p 53) does not describe his contemporaneous views and actions at the time of