CORRESPONDENCE

Testing for HIV: the non-medicolegal view Anonymous 497	Gas exocrine "failure" in the critically ill T J J Inglis, BM, and B L S Gallagher, FFARCS 499	Neonatal morbidity: an audit with medicolegal implications J S Samra, MRCOG, and others 501
Surgeons and others and HIV M B Smith, FFARCS 497	Cost of prophylaxis in joint replacement N I L Wilson, FRCS, and C R Dreghorn, FRCS 499	Merit awards R Gabriel, FRCP 501
Psychiatrists in the 1990s Adrianne Reveley, MRCPsych; K Checinski, MB 498	False negative colposcopic cervical biopsy M Jarmulowicz, MB, and others 499	Points Dehiscence of the infraorbital nerve as a cause of facial pain (G Flint); Low vision aids and the elderly (A J Elliott); Hearing aids (T L Pilkington; J W Stephenson) 501
Cocktail stick injury: a fatal outcome N Kasthuri, MRCPATH, and A Savage, MRCPATH 498	Providing donor organs for transplantation E Sherwood-Jones, FRCP 500	Collars and corsets (P Michaelson); The NHS debate (M Modell; W Norman-Taylor); Impact of new ice rinks on accident departments (D M Williamson and others; D M Evans); Motorcycle messengers (E T Lummis); Sexism and ageism (J Collin; Kay Dickensin) 502
The need to make rugby safer D A D Macleod, FRCS, and P H Pattison, FRCS; H C Burry, FRACP, and C J Calcinai, FRACS 498	Communication of drug treatment to general practitioners T R G Howard, MRCP 500	
	New year message A Paton, FRCP 500	
	Hyperbaric oxygen, carbon monoxide, and cerebral oedema P B James, MFOM 500	

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

Testing for HIV: the non-medicolegal view

SIR,—I note with more than casual interest the dilemma in which Dr A J Keidan and colleagues find themselves in the case of their patient positive for human immunodeficiency virus (23 January, p 288).

Firstly, their letter raises the problem of confidentiality, for when numbers of HIV positive women are so small there is a genuine danger of the patient's identity being discovered. Secondly, I am impressed by the authors' honesty in admitting their clumsiness in not only failing to provide adequate precounselling but also not informing the patient that the test was being performed. It is now two years since the incident, and I hope that we have all come a considerable distance in experience and sensitivity.

There is no doubt that the clinicians in this case were placed in a cleft stick, and whichever course of action they embarked on would have been hazardous. If the woman had received counselling consent for the test would probably never have been granted and she would probably have left the hospital as hostile as she did on receiving the positive result.

Outside London HIV positivity is still rare among women, and the odds are more than good that the result in any case of this nature will be negative, but clearly informed precounselling covering all eventualities is essential. Perhaps it is time that as a profession we were more honest about all tests that we perform on our patients; should we not explain why we want to test for syphilis and hepatitis B?

Unfortunately, this patient reacted dramatically to the seropositivity and sought help from three further medical experts, desperately clutching at straws in the hope that someone would give her the answer she wanted to hear—sadly to no avail. She has subsequently become embittered towards all doctors and feels no trust is possible; we represent a personal conspiracy that has to be beaten. To this

end she is resolved to show that the result is wrong and constantly displays her good health as witness.

Recently, a further complication has developed with the inevitable final evidence of her testimony; she is now pregnant and determined to proceed even after extensive counselling about the possible outcome.

There is no doubt that this case is not the first nor will it be the last case of an HIV positive pregnant woman but it does highlight numerous dilemmas which the medical profession has to recognise and debate honestly before our patients can receive genuine well informed advice. I thank Dr Keidan and colleagues for their letter even if it is only to serve as a lesson for us all.

THE PATIENT'S GP

Surgeons and others and HIV

SIR,—I would like to comment on the leading article by Mr A J W Sim and Professor H A F Dudley in the light of our experience (9 January, p 80).

In our operating rooms we provide anaesthesia for about 50 patients positive for the human immunodeficiency virus (HIV) each year, and for the past two months we have been implementing a set of precautions to prevent transmission of HIV.

The basic rule is that our health care workers must take precautions to prevent exposure to skin and mucous membranes when contact with blood or other body fluids of any patient may occur. Gloves must be worn for touching blood and body fluids, mucous membranes, or non-intact skin of all patients; for handling items or cleaning surfaces soiled with blood or body fluid; and for performing venepuncture and other vascular procedures. Gloves should be available in all rooms and be carried in the pockets of health care workers, and

they must be changed and hands washed after contact with each patient. Secondly, masks and protective eye wear or face shields must be worn during procedures likely to generate droplets of blood or other body fluids to prevent exposure of mucous membrane of the mouth, nose, and eyes. Gowns or aprons must be worn during procedures likely to generate splashes of blood and other body fluids. Thirdly, needles must not be recapped, bent, or broken by hand, removed from disposable syringes, or otherwise manipulated by hand. After use all needles, syringes, and sharps must be placed in puncture resistant containers for disposal. Fourthly, to minimise the need to use resuscitation mouthpieces resuscitation bags or other ventilation devices must be available wherever resuscitation is predictable. Finally, health care workers with exudative lesions or weeping dermatitis must refrain from all direct patient care and from handling patient care equipment.

In practice this means that we now wear gloves at all times. The type of glove does not appear to matter. There is no scientific evidence that intact latex and intact vinyl have any difference in barrier effectiveness. The thinner latex of condoms protects against transmission of HIV, and the Federal Drug Administration allows a failure rate of 4 in 1000 condoms. Sterile surgical gloves must conform to a Federal Drug Administration failure rate of 15 in 1000 (American Society for Testing and Materials), and non-sterile gloves must have a failure rate of 25 or less in 1000. We remove gloves and wash hands between patients and any time we leave the operating room.

The obvious disadvantages of this are, firstly, expense and inconvenience; secondly, the discomfort of wearing "plastic" gloves; and, thirdly, the loss of feel in performing delicate tactile procedures such as venepuncture. However, after an initial relearning period the success rate for procedures has returned to normal. To some of us

who wear prescription glasses protective eye wear has not presented a problem. Most of my colleagues have bought clear glass spectacles because the perspex goggles provided by the department distort vision. Gowns are now worn for all invasive procedures (except peripheral venepuncture).

We adhere to the precautions laid down in the policy document for all patients coming to our operating rooms. Although the occupational risk of HIV infection seems to be low, if transmission of HIV did occur during normal professional activity without reasonable precautions being taken employer compensation and insurance would be invalid. The moral of the modern era of health care is to take precautions against transmission of HIV.

MARK B SMITH

Department of Anaesthesiology,
University of Oklahoma,
Oklahoma City, Oklahoma 73152,
USA

Psychiatrists in the 1990s

SIR,—Dr Glin Bennet (23 January, p 274) suggests that psychiatrists can contribute most to community mental health teams by a return to their medical roots, and I agree with much that he says.

But Dr Bennet addresses his comments to "ordinary psychiatrists" suggesting that "we now have a new generation of neuropsychiatrists and psychogeriatricians . . . who have a clear role in aspects of psychiatry which make use of medical skills." Psychogeriatrics may indeed be a growing subspecialty, but neuropsychiatry is almost non-existent. Dr Bennet's idealised view may come from his proximity to the Burden Hospital, where three of the handful of designated neuropsychiatrists in Britain are to be found.

Neuropsychiatry is given a very low priority in training schemes, with only two of the 325 higher training posts in adult mental illness, and it is almost never considered in the creation of new consultant posts. Like biological psychiatry, neuropsychiatry does not have section status in the Royal College of Psychiatrists.

The 1990s may bring great changes in our understanding of the organic basis of some psychiatric conditions. Unless we claim our medical heritage and make use of our training and skills we may find ourselves treating a dwindling cohort of functional disorders, while striving to "maintain our usefulness" as the most expensive members of mental health teams.

ADRIANNE REVELEY

Maudsley Hospital,
London SE5 8AZ

SIR,—I disagree with Dr Glin Bennet's notion that the position of the doctor at the head of the mental health team is under serious threat (23 January, p 274).

A consultant psychiatrist remains the logical choice because of the amount of experience and skill gained during the 12 years or so it takes to reach that grade. It could be argued that the first half of this time is spent learning to be a doctor, while the second half is spent unlearning aspects of the role, but the training is not negligible. Furthermore, much continuity and consistency of philosophy and practice is ensured because this member of the multidisciplinary team will not be rewarded for success with promotion away from clinical work, unlike those in other disciplines, who can look forward to being managers or (undisciplined) team leaders, ensuring that their skills diffuse away from the direct care of mentally ill people.

The trend towards psychiatric treatment that is not purely medical could be taken as posing a threat to the doctor. Good psychiatric training, however, emphasises the triad of physical, psychological, and social factors in mental illness and its treatment, so that the doctor is in a good position to coordinate a multidisciplinary team. For major psychiatric illness—severe depression and schizophrenia, for instance—physical treatments are often a prerequisite for further psychosocial management. Few psychiatrists cannot have been touched by the advances in psychodynamic and behavioural therapies and in social psychiatry over the past 25 years, so a balanced approach is increasingly likely. I therefore found Dr Bennet's warning not to "retreat back into the medical model" at odds with his advice to "keep up with your general medicine."

Finally, at least some psychiatric training is geared to the needs of the exciting expansion in community psychiatry. I am presently in an experimental post acting to all intents and purposes as a general practitioner trainee. This is providing a fascinating perspective on the interface between primary and specialist psychiatric care. Psychiatrists are necessary but not sufficient on their own to provide good mental health care, and their role is guaranteed in the 1990s as long as this interface is kept in view.

KEN CHECINSKI

London SW19

Cocktail stick injury: a fatal outcome

SIR,—Three cases of injuries due to cocktail sticks have recently been reported by Mr C Rand (19-26 December, p 1658). We report a fourth case with a fatal outcome. There is no previous record of this occurrence.

A 38 year old man was admitted to the casualty department at Selly Oak Hospital with sudden onset of abdominal pain associated with profuse haematemesis while travelling in a car with his wife driving. On admission he was awake but confused and pale and had no recordable blood pressure. A rapid transfusion of blood was given and intubation attempted, but the larynx and cords could not be visualised owing to blood clots. In spite of all attempts at resuscitation he died 10 minutes later. His wife said that he had been complaining of pleuritic chest pain for a week and he had been given naproxen and erythromycin by his general practitioner. He had no history of a peptic ulcer.

At necropsy the most striking findings were in the chest. The oesophagus contained mucus

streaked with blood, and in its lower part, close to the cardio-oesophageal junction just above the diaphragm, half a cocktail stick about 4 cm long was impacted. The sharp point of the stick had penetrated through the oesophageal wall and into the adjacent aorta, into which it protruded by about 0.2 cm (figure). There was thrombus surrounding the stick at the site of penetration and a small quantity of thrombus between the aorta and the oesophagus. The stomach was markedly dilated and contained fresh blood clots weighing 1100 g. The small intestine was filled with blood, which was fresh in the upper part and altered in the lower part and extended to the ileocaecal valve. There was no blood in the colon. All other organs were normal. The cause of death was perforation of the oesophagus and aorta by a swallowed foreign body—half a cocktail stick. The coroner recorded a verdict of death by misadventure.

At the inquest the wife stated that on Christmas Day she and her husband were eating sausages wrapped in a piece of bacon secured by half a cocktail stick. The sticks were difficult to see and she herself found one in her mouth and removed it. Her husband jokingly said he must have swallowed his. This incident was not mentioned to the general practitioner, and at no time during the 10 days before death did the patient complain of dysphagia. A chest radiograph ordered by the general practitioner was normal but a cocktail stick would not be visualised on radiography. The patient's rapid deterioration would appear to have been due to torrential haemorrhage from the aorta at the point of penetration by the sharp end of the cocktail stick.

We thank Mr S K Banerjee for permission to report this case and Miss Robin Giles for the photograph.

NITHYA KASTHURI
ANN SAVAGE

Department of Histopathology,
Selly Oak Hospital,
Birmingham B29 6JD

The need to make rugby safer

SIR,—We write as honorary medical advisers to our respective rugby unions and members of the medical committee of the International Rugby Football Board.

We were disappointed in your leading article by Drs H C Burry and C J Calcinaï (16 January, p 149), the medical advisers to the New Zealand Rugby Football Union, although they appear to write from the Accident Compensation Cor-

The sharp point of the cocktail stick has penetrated into the wall of the aorta (arrow).