(oral and anal) inoculation of sperm and the development of antispermat antibodies in both men and women is needed. Widely read popular texts on sexuality advise that swallowing sperm carries no hazard.⁵ This advice may need to be modified for men and women who want children.

L A KAY

Royal Infirmary,
Sunderland SR2 7JE,
Tyne and Wear

- 1 Tomasi TB, Bienenstock J. Secretory immunoglobulins. *Adv Immunol* 1968;9:1-24.
- 2 Vaerman JP, Ferin J. Local immunological response in the vagina, cervix and endometrium. *Acta Endocrinol* 1974;194 (suppl):281-300.
- 3 McDermott M, Bienenstock J. Evidence for a common mucosal immunological system. I. Migration of B lymphocytes into intestinal, respiratory and genital tissues. *J Immunol* 1979; 122:1892-5.
- 4 Bronson R, Cooper C, Rosenfield D. Sperm antibodies: their role in infertility. *Fertil Steril* 1984;42:171-83.
- 5 Masters WH, Johnson VE, Kolodny RC. *Masters and Johnson on sex and human loving*. London: Macmillan, 1987:313.

Advice on contraceptive services for ethnic minorities

SIR,—In reply to criticisms of my article on contraceptive services for ethnic minorities (29 November, p 1365) by Drs Jill Tattersall and Jennifer Wordsworth (23 January, p 293) I accept that perhaps "advice" may have been the wrong word to use and would substitute "counsel" as the ideal that I aim for. I am sorry that what I was trying to say appears to have been misunderstood.

Of course, patients wish to make contact with a counsellor with whom they have confidence. What research evidence there is suggests that being understood is very important. This is confirmed by the paper by McAvoy and Raza to be published in *Health Trends*.¹ It sounds somewhat arrogant to say that "rudimentary English" is sufficient. This may be so for giving instructions, but surely not for "listening to patients' needs, likes, fears, and worries." It is correct that language is not the only consideration, but to say that some non-English speakers prefer an English doctor is as irrelevant as saying that some English speakers prefer a doctor whose first language is not English, as they do.

Advocates are not potentially disastrous. Advocates should come from the community they serve and should be selected and trained to represent the views of the patient, not of the doctors, or even of religious leaders. This may make the doctors feel very threatened. They work well in Hackney and are valued by the patients they serve and (most) local doctors. I suggest Drs Tattersall and Wordsworth read the article I referred to² in order to see how such a service may be set up successfully.

They do seem to have missed the two points I was trying to make, which were that patients should be offered a choice of counsellor which reflects the ethnic make up of a community, and that these counsellors should be trained for working across cultures.

JON FULLER

London E8 3JN

- 1 McAvoy BR, Raza R. Asian women: views on provision of contraceptive services and cervical cytology. *Health Trends* (in press).
- 2 Cornwell J, Gordon P. *An experiment in advocacy, the Hackney multiethnic women's project*. London: Kings Fund Centre, 1984.

Thrombocytopenia and interferon

SIR,—Dr M E Ellis and others and Dr Andrew M L Lever and others describe three cases in which alfa interferon was suggested as alternative treatment for thrombocytopenia associated with

human immunodeficiency virus infection (12 December, p 1519). As thrombocytopenia may be caused by interferon, however, this suggestion should be treated with caution.

We have studied the haematological, metabolic, and neurological effects of alfa interferon in the treatment of amyotrophic lateral sclerosis.¹ We used high intravenous doses of human leucocyte alfa interferon ranging from 25 MU/day up to 200 MU/day for five days.² On this schedule a dose dependent thrombocytopenia was a complication, which was reversed when interferon was stopped. After high dose interferon treatment thrombocytopenia developed more slowly but lasted longer than leucopenia (table). In spite of discontinuation of the treatment on day 5 the thrombocyte count

Leucocyte and thrombocyte counts before and after interferon infusions to 11 patients with amyotrophic lateral sclerosis

	Cell counts ($\times 10^9/l$; mean (and range))	
	Leucocytes	Thrombocytes
Before interferon	6.1 (4.8-7.9)	263 (262-289)
Day 2	3.4 (2.8-4.2)	176 (160-178)
Day 3	2.0 (1.0-2.9)	113 (96-140)
Day 7	3.6 (2.8-4.7)	71 (66-80)

dropped until day 7, returning to normal three to five days later. No bleeding complications were seen. Because of neurological side effects or severe leucopenia the interferon doses had to be reduced in some cases.

This thrombocytopenic effect was not due to dilution of blood because of infusion fluids, as all infusions ceased at day 5 and the thrombocyte count continued to decrease. The first decrease in thrombocyte count in the case of Dr Ellis and colleagues may well have been caused by interferon. We would like to emphasise the dose dependent thrombocytopenic effect of interferon and argue that it cannot be recommended for the treatment of thrombocytopenia.

MARKUS FÄRKKILÄ
MATTI IIVANAINEN

Department of Neurology,
University Hospital,
Helsinki, Finland

- 1 Färkkilä M, Iivanainen M, Roine R, *et al*. Neurotoxic and other side effects of high dose interferon in amyotrophic lateral sclerosis. *Acta Neurol Scand* 1984;70:42-6.
- 2 Cantell K, Hirvonen S, Koistinen V. Interferon. *Methods Enzymol* 1981;78:499-505.

Providing donor organs for transplantation

SIR,—Like Dr Eric Sherwood-Jones (13 February, p 500), I watched the BBC's *That's Life* programme on organ transplantation and noted the general feeling of the participants that the main cause of the shortage of donor organs was that intensive care units were not asking relatives of potential donors for the organs. Transplant surgeons are clearly frustrated by this and consider neurosurgeons, neurologists, intensivists, and anaesthetists as the villains of the piece, but I think they should consider their own role in organ procurement more closely.

In the early days of multiorgan transplantation the referring doctors had to coordinate several teams from various parts of Britain who all wanted to arrive at different times and who all felt that their organs had the greatest priority—leading to delays, often at the last minute, which caused distress to relatives and intensive care staff alike. Since the advent of the transplant coordinators this

has improved, but to reduce the stress on both relatives and intensive care unit resources whenever I refer a patient for organ harvest (and I ask all relatives) I specify a time by which the procedure must be completed, and if the transplant teams cannot fall in with that time then they will miss out on the organ. It is not tolerable for switching a ventilator off to be delayed while the transplant teams get their act together.

Secondly, I am sure it would help increase the awareness of the need for organ donation if the referring teams got better feedback. I have been asking relatives for organs for the past seven years and have never had a letter from a transplant surgeon either thanking me or my team for obtaining the organs or letting me know how the patients who had the transplants are getting on. It would be a simple matter to send a discharge summary of a patient after transplantation to the doctors who procured the organs so they could get some idea of the effectiveness of the exercise. Likewise it might help improve outcome if we were given some guidance on the best way to preserve organs for the 48 hours or so before their removal, and a better rapport might be built up if the transplant teams were to write articles on this subject in journals read widely by neurosurgeons, anaesthetists, and intensivists, or perhaps offered to give lectures.

I do not think that to appear on television and publicly berate colleagues for not providing enough organs is an approach that will improve the supply, and something more constructive ought to be offered by the transplant surgeons.

P RICHARDS

Department of Neurosurgery,
Charing Cross Hospital (Fulham),
London W6 8RF

Retreat from openness

SIR,—Dr Richard Smith's article (19-26 December, p 1633) confirms the fears which I have had for some time that the Department of Health and Social Security propaganda machine has replaced the daily experience and knowledge of hospital staff as the source of "official truth."

In addition to the manipulation of data, release of information at times which limit its availability, and the silencing of staff who speak out, we have the more recent experience of a nationwide "cleansing" of district health authorities. This process involves the unseating of "troublesome" trades union members of district health authorities when their terms expire and the refusal to appoint newly elected trades union nominees who are known to be actively concerned with health service affairs.

By tradition trades union nominees are automatically accepted for the single trades union seat unless the candidate is ineligible. The intention in providing a trades union seat was to ensure that district health authorities reflected a wide range of interests and opinions; but these seats are now being taken from the Trades Union Congress and given instead to supporters of the government or to inexperienced generalists.

There were many examples of this practice during 1987. In City and Hackney Lucy de Groot, an efficient and valuable trades union member, was unseated and replaced with a close relative of the prospective Conservative parliamentary candidate for south Hackney. In North Warwickshire Jack Sprung was unseated after five active years on the authority. In Tower Hamlets Tony Jewel was prevented from taking the general practitioner seat despite being the popular choice of general practitioners in the district. In the North Western region three candidates were rejected. In Dartford and