Royal College of Physicians of London (September 1984, revised November 1984) clearly state that such review is not required. The use of other androgenic steroids for intractable pruritus has been recommended for many years.¹ Our final sentence recommends formal controlled studies because we realise that our report is preliminary.

R P WALT

University Department of Medicine,
Queen Elizabeth Hospital,
Birmingham B15 2TH

1 Sherlock S. *Diseases of the liver and biliary system*. 6th ed. Oxford: Blackwell Scientific Publications, 1981:221.

Self referral to consultants

SIR,—The members of the Edgware Trainers' Workshop raise important points about the increasing tendency of patients to refer themselves to hospital. I suspect that the problem is much larger than is generally recognised, particularly in accident and emergency departments.

Many patients, dissatisfied with their general practitioner's appointments system or with the treatment already offered by their family doctor, will seek advice directly from the accident and emergency department for problems other than accidents and emergencies. This is even more common out of normal surgery hours, and in most cases the general practitioner never knows that the patient has attended, despite some departments having a policy of sending a letter.¹ Even when a letter does arrive I wonder how many doctors raise the matter with their patients.

The junior medical staff of the accident department are often unsure of their ground in redirecting patients with inappropriate conditions and tend to treat rather than educate. As a result of this easy alternative source of primary care the patient may never see a doctor who is experienced enough to plan long term follow up. My own advice to patients who attend for "second opinions" is that it would be unethical for me to change their general practitioner's own treatment, but those who come for help or advice about a genuine emergency have often taken the only course of action open to them and we should give any necessary treatment, communicating this to the general practitioner.

The training of senior accident and emergency nurses in triage has been suggested.² If this system were to gain acceptance patients could obtain advice at the time of registering in the department, saving those who do not need hospital care a long wait and allowing the staff more time to deal with serious cases.

ROBERT COCKS

North Western Injury Research Centre,
Hope Hospital,
Salford M6 8HD

1 Sherry M, Edmunds S, Touquet R. The reliability of patients in delivering their letter from the hospital A and E department to their general practitioner. *Arch Emerg Med* 1985;2:161-4.

2 Yates D. Nurse triage in the accident and emergency department. *Journal of the Royal Society of Health* 1987;4:153-4.

SIR,—The article by Dr Gerald Michael and colleagues (27 February, p 640) deploring self referrals to consultants provoked a torrent of correspondence (19 March, p 860). Mr Grant Williams said that self referral was "dangerous" and that patients might "receive treatment from physicians and surgeons who are not qualified to deal with their illness." In mental health, where self referral to non-medical practitioners in the community has flourished, Dr D M Bowker stated, "Any decision to supplant the primary screening role of the general practitioner by those

with less training... is theoretically reducing the diagnostic quality of the service."

These are noble ideals but not realistic. As Dr M M Lieberman pointed out, the general practitioner may not have received any vocational training in the specialty appropriate to the patient's problem and the appointments system and deputising services used by many general practitioners make it difficult for patients to see their own general practitioner in an emergency or out of hours and at weekends.

I conducted a retrospective survey of psychiatric problems presenting to a London casualty department over six months in 1986. Of 464 patients suffering from a psychiatric disorder, only 17 presented with a referral letter from their general practitioner. The patients were not just casual attenders: 168 were admitted to hospital, 63 were referred to the psychiatric emergency clinic, and 48 were offered outpatient treatment.

If general practitioners cannot provide an accessible and flexible service to their patients and if they do not possess the skills to manage particular illnesses they should not be surprised when their patients bypass them.

JOHN DUNN

Maudsley Hospital,
London SE5 8AZ

Postviral fatigue syndrome

SIR,—The paper by Dr Anthony David and others (5 March, p 696) has been overtaken perhaps by the recent publication of Yousef *et al* showing persisting enteroviral infection in many sufferers.¹ None the less, I am surprised the authors state that implication of a single virus would "greatly enhance the validity of the syndrome" given that well established syndromes such as hypertension, dementia, and hypothyroidism all have many possible causes.

The authors are correct to call for better definition of the syndrome. They recommend precision in the use of "postviral," but since the virus persists the term should now be abandoned. Their definition of the initial viral infection would include fever, but fever is often slight or absent^{2,3}—indeed, its absence in the presence of infection might be thought ominous. Choosing only those patients with fever and up to several months of illness risks throwing out the baby with the bathwater. The syndrome must be recognised promptly since advice to rest may make it less severe.² Furthermore, fluctuating neurological signs and symptoms often occur early on.^{2,3} Detailed observations of their pattern, not just at the bedside but also after limited physical activity, may increase understanding of the disease and help early recognition.

The baby may also be at risk from the use of the present state examination and the general health questionnaire.^{4,5} While both are highly reliable, their validity does not extend to a clear distinction between emotional lability of organic type (similar to that seen in certain brain stem disorders) and emotional upset which is a reaction to being ill. Both types of upset may occur in the postviral fatigue syndrome, in which brain infection is highly likely. The authors suggest that psychological changes may help cause the disease. In patients known to me they result from it; some (including children) whose symptoms were wrongly thought psychogenic were prescribed exercise, which made their illness worse.

The authors believe that understanding of this syndrome has been hindered by the work of doctors who are themselves sufferers. They claim that while medical training affords insight into subjective aspects of illness, it is "axiomatic" that

objectivity is not similarly enhanced. They would have to show, however, that research by sufferers was inferior to that by non-sufferers. Excellent reviews of the syndrome were written by Parish,^{6,7} himself a sufferer.⁸ Are the authors saying that Parish hindered understanding? Have they thoroughly checked that none of the better research among their references was done by doctors who suffered from the disease? If they did so they might be surprised. Percival Pott turned to writing about orthopaedics while recovering from a fractured tibia⁹; he does not seem to have hindered understanding.

A V MARK HUGHSON

Glasgow G12 0PU

- 1 Yousef GE, Bell EJ, Mann GF, *et al*. Chronic enterovirus infection in patients with postviral fatigue syndrome. *Lancet* 1988;i:146-50.
- 2 Ramsay AM. 'Epidemic neuromyasthenia' 1955-1978. *Postgrad Med J* 1978;54:718-21.
- 3 Compston ND. An outbreak of encephalomyelitis in the Royal Free Hospital Group, London, in 1955. *Postgrad Med J* 1978;54:722-4.
- 4 Wing JK, Cooper JE, Sartorius N. *Measurement and classification of psychiatric symptoms*. London: Cambridge University Press, 1974.
- 5 Goldberg DP. *Manual of the general health questionnaire*. Windsor: NFER Publishing Company, 1979.
- 6 Parish JG. Epidemic neuromyasthenia: a reappraisal. *J Int Res Communications* 1974;2:22-6.
- 7 Parish JG. Early outbreaks of 'epidemic neuromyasthenia.' *Postgrad Med J* 1978;54:711-7.
- 8 Ramsay AM. Accounts of outbreaks. Chairman's opening remarks. *Postgrad Med J* 1978;54:710.
- 9 Ellis H. *Bailey and Bishop's notable names in medicine and surgery*. 4th ed. London: H K Lewis, 1983:189-92.

SIR,—The postviral fatigue syndrome is increasingly being recognised as a distinct neurological entity with an organic background.¹ Dr Anthony S David and colleagues (5 March, p 696) referred to our study² in which we applied the technique of single fibre electromyography to measure jitter values in 40 patients with the syndrome in whom any concurrent neuromuscular problem had been carefully excluded. That study produced strong evidence to support the organic nature of this condition and provided some clues to its nature.²

Since then we have measured the fibre density in 10 of the original patients to seek more information about the pathophysiology of muscle disease in this syndrome. The patients, four of whom were women, were aged 18-61 (mean 40.9 (SD 15.4)) with a mean duration of illness of 4.1 (2.3) years. They were selected to include those who did not consume alcohol and who had shown positive results on virological, immunological, and muscle biopsy studies.¹ The patients were re-evaluated according to our previous protocol² to exclude any concurrent neuromuscular problem. Fibre density measurements were performed on the right extensor digitorum communis muscle according to standard procedures.³ The results showed normal fibre density but abnormally increased jitter values (table). Increased fibre density occurs in various neurogenic and myopathic disorders with muscle fibre degeneration in which there is a spatial rearrangement or a change in the number of muscle fibres in the motor unit.³ Normal fibre density in these patients with the postviral fatigue syndrome therefore suggests absence of these phenomena and further supports our argument that the jitter abnormalities probably arise from muscle membrane dysfunction.²

Our theory is also supported by several non-neurophysiological findings. Electron microscopy of muscle biopsy specimens showed definite though non-specific changes.¹ Examination of patients with the postviral fatigue syndrome by phosphorus-31 nuclear magnetic resonance showed evidence of disorder of muscular metabolic regulation,⁴ which might arise from muscle fibre