CORRESPONDENCE

The NHS versus private medicine P F Jones, FRCSed 1600	Training in social work Beulah R Bewley, FFCM, and A Elliott, FRCGP 1602	Funding and patient care in Scotland J A Ford, FRCPED 1604
Gas gangrene D J Coleman, FRCS, and A G Batchelor, FRCS; C Gollidge, MB, and others 1600	The Biopsy cut procedure for renal biopsies D P Worth, MRCP, and others 1603	Euthanasia P Hawker, MRCP; J Miller 1604
Proteinuria in malignancy P Gosling, MRCPATH, and others 1601	After the horror R Ramsay, MB, and S A Stansfeld, MRCPsych 1603	Drug points Toxic epidermal necrolysis and trimethoprim (G Das and others) 1604
Crisis in medical education G D Roberts, MRCP; Helen Vos, MB; C B Brown, FRCP; J A Davis, FRCP 1601	Nurses' pay Catherine A Gerrish, MSC 1603	Idiosyncratic reaction resembling toxic epidermal necrolysis caused by chloroquine and Maloprim (P A Phillips-Howard and J W Buckler); Acyclovir malabsorption (A Mindel and O Carney); Sterile abscess formation by continuous subcutaneous infusion of diamorphine (P J Hoskin and others) 1605
Non-mydratric photography and diabetic retinopathy R Taylor, MRCP; D Jones, FRCS, and others; R E J Ryder, MRCP 1602	Treatment for mild hyperparathyroidism S Minisola, MD, and others 1603	
	Deaths from injury E K Alpar, FRCS, and others 1604	
	BMA council election results D Gullick, MB 1604	

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

The NHS versus private medicine

SIR,—Sir Richard Bayliss (21 May, p 1457) documents the recent growth of independent hospitals and states his belief that "the electorate will accept, even welcome, a mixed medical economy, provided, and this is an important proviso, the NHS is improved, and 'free' medical care continues to be available at the time of need." The proviso is of fundamental importance because government ministers assume that encouragement of private hospital practice must benefit the acute hospitals of the NHS. Personal experience suggests that this should not be assumed.

When I was appointed as a consultant surgeon in 1958 the only contract available was for eight sessions so I began to see private patients and, for the reasons given by Sir Richard, found this enjoyable. As the volume of NHS and private work expanded, however, it became increasingly apparent that neither responsibility was being properly discharged. One cannot control workload by picking and choosing among referrals, so after eight years I applied for a whole time contract. This had several effects.

Firstly, instead of visiting four hospitals once or twice each day I could concentrate my efforts in one adult and one children's hospital. There was time to do a full ward round each day with our team and to organise outpatient clinics so that patients were usually seen by the same doctor and without undue haste—this can transform follow up clinics into useful occasions for both parties. We could set up a records system which showed our immediate and long term results. Secondly, there was time to teach. One of the most important tasks of the surgeon is to help the next generation to learn good operative technique, and there is no way to achieve this except for the surgeon to act as first assistant until the trainee is ready for opportunities to operate independently. Assistance will continue to be needed as more complex procedures are tackled. I consider the time given to assisting colleagues at all stages of training to have been

most enjoyable and rewarding. The increasing attention being paid to this aspect of surgical apprenticeship is most welcome.^{1,2} There is evidence that it still does not receive enough attention,³ and it is time that it was more widely accepted that some emergencies require technically advanced surgery for the best results and that this requires the presence of an experienced surgeon.

The commitment of the surgeon to any patient is comprehensive, but in private work the lack of junior colleagues means it can be heavy and unpredictable. It is unreasonable to call for more elective surgery to be performed in private hospitals and also to expect overextended surgeons to give as much time as they would wish to teaching and to out of hours emergency work in their NHS units.

What is needed urgently is for consultant

staffing to reach a level at which there is time for proper care of all patients, for teaching, and for investigation. There is a sad discrepancy between present surgical waiting lists and the fund of highly trained registrars and senior registrars longing to be given more responsibility. It would be a useful start, amidst all the advocacy of more private care, to give practical encouragement to the many who would be glad to pursue a full time career in academic or NHS hospital practice.

PETER F JONES

Aberdeen AB1 9HR

- 1 Matheson NA. Surgical technique. *Br J Surg* 1987;74:1190.
- 2 McKeown KC. Reoperative surgery for early complications following abdominal operations. *J R Soc Med* 1988;81:307.
- 3 Buck N, Devlin HB, Lunn JN. *The report of a confidential enquiry into perioperative deaths*. London: Nuffield Provincial Hospitals Trust/King's Fund, 1987:38.

Gas gangrene

SIR,—Dr C R Kirk and others (30 April, p 1236) and Professor Stephen T Holgate (30 April, p 1213) have clarified our position when considering treatment with penicillin in patients giving a history of penicillin allergy.

We would, however, strongly disagree with Dr Kirk and others that primary closure of soiled or penetrating wounds should be delayed. It is quite clear from the case report that the child's wound, although explored and cleaned extensively, was not excised. It was this failure to convert an untidy wound (which contained non-viable tissue) into a tidy one (in which the wound edges were viable) by the simple process of wound excision that led to the onset of gas gangrene. There is now ample evidence that untidy wounds such as deep dermal burns¹ and type III open tibial fractures² can safely undergo primary closure provided that adequate wound excision has been performed.

"Primary wound closure" may mean anything from direct suture, to skin grafting, local flaps, or free tissue transfer. It is important that wound excision should not be compromised by the surgeon's ability to close the resultant defect. Ideally the surgeon performing the excision should have access to whichever technique is most appropriate for its closure. Delayed primary closure is still appropriate when the surgeon cannot be certain, whether due to inexperience or the nature of the injury, that wound excision is complete. The real lesson of this case report is the long known importance of wound excision.³

DAVID J COLEMAN
ANDREW G BATCHELOR

St James's University Hospital,
Leeds LS9 7TF

- 1 Janzekovic Z. A new concept in the early excision and immediate grafting of burns. *J Trauma* 1970;10:1103-8.

- 2 Godina M. Early microsurgical reconstruction of complex trauma of the extremities. *Plast Reconstr Surg* 1986;78:285-92.
- 3 Trueta J. *Treatment of war wounds and fractures*. London: Hamish Hamilton, 1939.

SIR,—We do not agree with Dr C R Kirk and others (30 April, p 1236) that penicillin is so much the drug of choice in treating gas gangrene that a test dose should be given to confirm reported allergy.

Their comment that other agents are ineffective because of poor penetration of tissues cannot be substantiated. Metronidazole penetrates tissues well, achieving concentrations equivalent to those found in serum.¹ Dose is important in determining tissue concentrations, and we are not told the doses of metronidazole or amoxycillin trihydrate and clavulanic acid that were given in this case. In a murine model of gas gangrene responses to metronidazole were highly dose dependent,² and survival of mice treated with penicillin was not significantly better than that of untreated controls in spite of very high serum concentrations.

Antibiotics and hyperbaric oxygen have an adjunctive role in the management of clostridial gas gangrene, with surgical débridement offering the mainstay of treatment. It is not appropriate to give a drug, even as a test dose, to a patient with a possible allergy when the drug has not been shown to have clear benefit. Metronidazole may offer a good alternative to penicillin in such cases.

C GOLLEDGE
A KEIL

Department of Microbiology,
Royal Perth Hospital

T MCKENZIE

Department of Pharmacy,
Sir Charles Gairdner Hospital,
Nedlands,
Western Australia 6009

- 1 Ralph ED. Clinical pharmacokinetics of metronidazole. *Clin Pharm* 1983;8:43-62.
- 2 Stevens DL. Comparison of clindamycin, rifampin, tetracycline, metronidazole, and penicillin for efficacy in prevention of experimental gas gangrene. *J Infect Dis* 1987;155:220-8.

Proteinuria in malignancy

SIR,—Dr Nick Sawyer and colleagues (7 May, p 1295) describe an association between proteinuria and malignancy. Although proteinuria in patients with diabetes mellitus predicts diabetic nephropathy, it is also a risk factor for generalised vascular disease.¹ Diabetic patients with microalbuminuria, in contrast to those with normal albumin excretion, show increased vascular permeability to radio-labelled albumin.²

In patients with no clinical or biochemical evidence of renal disease we have found increases in renal permeability to plasma proteins after trauma,³ sepsis associated with burns and trauma,⁴ and acute pancreatitis⁵ and during elective surgery.⁶ We have proposed that increased renal permeability is part of a generalised increase in vascular permeability, rather than indicative of primary renal disease.⁵ This hypothesis is consistent with the finding that increased vascular permeability to radiolabelled albumin is found in patients after surgery and in septic shock.⁷ In that series nine out of 11 patients with malignant disease also showed increased vascular permeability. The mechanisms for these changes are not certain, but systemically active substances which cause massive increases in vascular permeability have been identified in several conditions, including advanced malignancy.⁸

We suggest that the proteinuria that often accompanies malignancy may not be due primarily

to subclinical nephritis but is a reflection of an increase in overall vascular permeability. We speculate that in advanced neoplastic disease mediators are present that produce systemic effects, which in the kidney are manifest as mild proteinuria. Rapid advances in the automated assay of low concentrations of specific proteins in urine mean that this relation could have implications for the early diagnosis and treatment of malignancy and many other conditions.

PETER GOSLING
CLIFFORD P SHEARMAN

Departments of Biochemistry and Surgery,
Selly Oak Hospital,
Birmingham B29 6JD

ANNE J SUTCLIFFE

Birmingham Accident Hospital,
Birmingham B15 1NA

- 1 Winocour PH, Durrington PN, Ishola M, Anderson DC, Cohen H. Influence of proteinuria on vascular disease, blood pressure, and lipoproteins in insulin dependent diabetics. *Br Med J* 1987;294:1648-51.
- 2 Feldt-Rasmussen B. Increased transcapillary escape rate of albumin in type 1 (insulin dependent) diabetic patients with microalbuminuria. *Diabetologia* 1986;29:282-6.
- 3 Gosling P, Sutcliffe AJ. Proteinuria following trauma. *Ann Clin Biochem* 1986;23:681-5.
- 4 Gosling P, Sutcliffe AJ, Cooper MACS, Jones AF. Burn and trauma associated proteinuria: the role of lipid peroxidation, renin, and myoglobin. *Ann Clin Biochem* 1988;25:53-9.
- 5 Gosling P, Shearman CP. Increased levels of urinary proteins: markers of vascular permeability? *Ann Clin Biochem* (in press).
- 6 Gosling P, Shearman CP, Gwynn BR, Simms MH, Bainbridge ET. Microproteinuria: response to operation. *Br Med J* 1988;296:338-9.
- 7 Fleck A, Raines G, Hawker F, et al. Increased vascular permeability: a major cause of hypoalbuminaemia in disease and injury. *Lancet* 1986;i:781-3.
- 8 Beutler B, Cerami A. Cachectin and tumour necrosis factor as two sides of the same biological coin. *Nature* 1986;320:584-8.

Crisis in medical education

SIR,—Professor Peter Richards (7 May, p 1278) has, unwittingly, made a strong case for moving a large part of undergraduate medical education away from the universities and into general practice. As he states, "As the emphasis moves to outpatient and community clinics, primary health care and prevention . . . the need is not for more educationalists but for more clinical teachers who understand the complexity of medicine." The existing network of general practitioner trainers, course organisers, and associate and regional advisers is well placed to meet this challenge along with the academic departments of general practice.

Perhaps some of the criticisms levelled at undergraduate education by Dr H J N Andreyev (7 May, p 1326) would then be overcome. Many of his criticisms are of the style of education and the lack of regard for patient autonomy. In postgraduate education in general practice we have had to learn the techniques needed to facilitate adult learning, and our work has taught us the importance of patient autonomy.

GEOFFREY D ROBERTS

Camberley,
Surrey GU15 2HJ

SIR,—Dr H J N Andreyev (7 May, p 1326) does not say where he trained, but his clinical course was obviously not at Addenbrooke's Hospital in Cambridge.

Much of what he says contains some truths, particularly with regard to the rigours of preclinical training, but surely his negative feelings relate to the hospitals where he trained. I thoroughly enjoyed the clinical course at Addenbrooke's, even though it was short (26 months). Individual teaching was promoted, and on peripheral hospital attachments there were often only one or two students on each firm. The teaching was

excellent, and much of this was because of mutual cooperation.

Learning of lists is tedious but much of the knowledge acquired is necessary. One often needs to think quickly and recall a likely differential diagnosis in order to act appropriately. Did Dr Andreyev do any student locums? I did three and found them invaluable. I was well supported during each of them.

Some attitudes that prevail in medical schools are unpleasant. Many doctors are downright rude, but there is no need for the enlightened to perpetuate this behaviour. I wonder what Dr Andreyev's career choice will be. He might find that some of his junior hospital training is also not what he had hoped for. He might be more fulfilled working in general practice if he does not become too bitter along the way.

HELEN VOS

Oxford OX4 1XD

SIR,—Dr Peter Richards (7 May, p 1278) discusses the crisis in undergraduate medical education. A feature of teaching hospitals has been unnecessary divisions between staff employed by the university and by the National Health Service. This has been exacerbated recently by problems in university funding and the loss of staff and support. Many people who work in teaching hospitals know the areas of division, friction, and lack of communication between these two groups of staff.

I suggest that teaching hospitals should be separately funded. All their staff should have equal status. If required, all consultants would be senior lecturers and, where appropriate, could be upgraded to reader or professor according to their abilities in clinical work or research.

New consultant staff should be appointed according to the overall requirements of the hospital. This would replace the current arrangement where appointments are limited by the nature of the post vacated. Consultants appointed to district general hospitals who have responsibilities for teaching medical students should also be part of the new system. These suggestions would remove the financial support of teaching hospitals away from the general university system because of the special requirements of providing clinical care to patients over and above research and teaching. They would put all consultants on an equal footing and remove confusion in the running and the development of the teaching hospital with respect to clinical service, teaching, clinical research, and basic research.

C B BROWN

Royal Hallamshire Hospital
Sheffield S10 2JF

SIR,—Professor Peter Richards discusses medical education (7 May, p 1278), but is there really a crisis? There is no shortage of first class recruits, and the product is excellent, although perhaps rather too stereotyped. We forget that the student-teacher ratio that we regard as optimal could be maintained by reducing the student intake as well as by shoring up the teaching establishment. Having spent nearly all my life in teaching hospitals—metropolitan and provincial, British and foreign, concerned with undergraduate and postgraduate education—I wish to make the following suggestions.

There is an overwhelming case for a reform of sixth form education, which, whether we like it or not, is determined by selection for university places. A regrouping of subjects is needed into the inorganic (mathematics, physics, and inorganic chemistry); the organic (biochemistry, botany,