CORRESPONDENCE

Accuracy of cervical cytology screening J R Hughes, MRCP 1534	Controlling symptoms in advanced cancer Jennifer Bulman, MB 1537	Improving outpatient clinics E Moonman, MSC 1539
Vocational training in general practice Dorothy M B Ward, MB; C Sayer, MB, and others; J Morris, MRCP, and others; J D Sinson, FRCP; D G Wilson, FRCP 1534	Prevalence of Gardnerella vaginalis K G Kerr, MB, and others; T C O'Dowd, MRCP, and R R West, PHD 1537	Iron and infection J J Bullen, PHD, and C G Ward, MD 1539
Rheumatic fever H McIntyre, MB, and others; P G Weaving, MRCP; J Pitts, MRCP 1535	Fetal intracardiac transfusions L Heyse-Moore, MRCP; M Westgren, MD 1538	Körner, nomenclature, and SNOMED W Dodd 1539
Acute appendicitis and social class N Anderson, MFCM, and Anne Cockcroft, MRCP 1536	The sickening of medical research A P Hemingway, FRCP, and D J Allison, FRCP 1538	Points Pre-employment chest radiographs (J M Grange; Freda Festenstein); Measles mortality (A O H Tellegen); Performance indicators (W A Wallace); Alcohol and road safety (B Sheerman); New drugs (K Summers); Insulin for the non-insulin dependent? (J S Garrow); Home visiting by consultants (R N Baldwin); Cervical biopsy (M B Gillett); Letter from Fiji (T G Hawley) 1540
Malaria prophylaxis L Rombo; P Fletcher, MRCP 1536	Penicillamine nephropathy P Emery, MRCP, and Gabriel Panayi, MRCP 1538	
Complications of nasogastric feeding C L M H Gibbons, MB, and others 1537	Psychiatric illness among British Afro-Caribbeans G Glover, MRCPsych 1538	
	Delays in treating bladder cancer M Jones, FRCS, and others 1539	

- All letters must be typed with double spacing and signed by all authors.
- No letter should be more than 400 words.
- For letters on scientific subjects we normally reserve our correspondence columns for those relating to issues discussed recently (within six weeks) in the *BMJ*.
- We do not routinely acknowledge letters. Please send a stamped addressed envelope if you would like an acknowledgment.
- Because we receive many more letters than we can publish we may shorten those we do print, particularly when we receive several on the same subject.

Accuracy of cervical cytology screening

SIR,—Mr John A Giles and others (16 April, p 1099) discuss the accuracy of cervical cytology screening. Screening competes with the traditional "demand led" services of a chronically underfunded health service. It is essential, therefore, that screening programmes are effective lest resources become diverted from areas of proved need to schemes with populist appeal but little scientific basis.

Since the 1960s screening for cervical cancer has been based on the assumption that the disease has a long preinvasive phase, rendering it amenable to testing at regular intervals. The work of Mr Giles's team and others suggests that the clinical course of at least some cervical cancers is changing, with an earlier age of onset and more rapidly invasive behaviour.^{1,3} The Intercollegiate Working Party on Cervical Cytology Screening has, therefore, recommended that screening should start at age 20 and continue at three yearly intervals.⁴ The behaviour of cervical dysplasia is unpredictable, and there is little evidence that this policy will reduce deaths from cervical cancer. The rapidly invasive variant may not be amenable to any practical screening interval.⁵ The only justification for screening is a proved reduction in mortality, not merely increased identification of abnormal cases.

Circumspection is required in interpreting cervical cancer statistics in Britain. Although deaths in women under 35 have increased from around 30 a year in the 1960s to about 110 a year in the 1980s, the mortality was as high as 70 a year in the 1950s (Office of Population Census and Surveys, personal communication). Comparing these small numbers with a total yearly mortality from all malignancies of about 150 000 puts the problem into perspective. The chances of screening

significantly reducing such a low mortality must be small, especially when the poor compliance of some populations is considered.⁶

National statistics relating to "positive" smears should be interpreted with equal caution. Returns of form SBH 140 from pathology laboratories to the Department of Health and Social Security show an overall increase in the proportion of positive cases. In some regions, however, notably Wessex, there has been virtually no change in incidence in the past decade whereas other regions have shown a large increase. It is important to know whether these discrepancies reflect different disease patterns or merely differing pathological or gynaecological management. It is also impossible to be sure whether there is a true increase in positive cases in younger women as positivity in any age group can be "weighted" by an increased number of smears performed in that group and by repetition of positive smears (it is positive smears that are counted, not women with positive smears).

In Britain we have the advantages of a large population base and a nationally organised health service. It seems absurd to miss the opportunity to organise properly controlled clinical trials to establish the behaviour of cervical neoplasia and assess the effectiveness of screening in younger women. The alternative is to plunge hysterically into a hotch potch of district based schemes which will be enormously expensive and can never be proved to have saved lives.

JOHN R HUGHES

Havant Health Centre,
Hampshire PO9 2AQ

1 Macgregor JE. Rapid onset cancer of the cervix. *Br Med J* 1982;284:441-2.

2 Wolfendale MR, King S, Usherwood MM. Abnormal smears: are we in for an epidemic? *Br Med J* 1983;287:526-8.

3 Campion MJ, McCance DJ, Cuzick J, Singer A. Progressive potential of mild cervical atypia: prospective cytological, colposcopic, and virological studies. *Lancet* 1986;ii:237-40.

4 Sharp F, Duncan I, Evans DMD, et al. *Report of the intercollegiate working party on cervical cancer screening*. London: Royal College of Obstetricians and Gynaecologists, 1987.

5 Hakama M, Rasamen-Virtanen U. Effect of a mass screening programme on the risk of cervical cancer. *Am J Epidemiol* 1976;103:512-7.

6 Nathoo V. Investigation of non-responders at a cervical screening clinic in Manchester. *Br Med J* 1988;296:1041-2.

Vocational training in general practice

SIR,—Dr Marie Campkin and colleagues (7 May, p 1331) imply that the Joint Committee on Post-graduate Training for General Practice has based its decision to withdraw recognition of vocational training in the North East Thames region solely on the report of its visitors to the Bloomsbury scheme in June 1987. This is far from so.

The committee has national responsibility for maintaining standards for vocational training. It arranges for its visitors to review the arrangements in each region every other year. These visits are undertaken in different parts of a region on each occasion. Reports of the visits to the North East Thames region have highlighted a series of problems extending back to 1983.

In 1985 the visitors reported that the standards of medical records in a large proportion of training practices in the region did not match the standards required by the committee's minimum criteria. Following receipt of that report the committee arranged for its officers to meet representatives of the North East Thames region to discuss the matter further. The region gave assurances that in future its criteria for selecting trainers would be in

accord with the committee's recommendations, and the committee felt able to reject the recommendations of its visitors in 1985 that recognition should be withdrawn at that stage.

In 1987 the visitors reported that, although the regional criteria were within the committee's recommendations, there were still problems in implementation and that two practices did not fulfil the basic criteria. This was discussed further by the officers of the committee and the representatives of the region. In February of this year the committee decided that, in the light of the reports of its visitors over six years, it could no longer recognise vocational training in the North East Thames region and that recognition should be withdrawn from 1 February 1989.

Dr Stella Lowry (7 May, p 1320) asked why action against individuals could not have sufficed. The committee has no direct relation with trainers or training practices. Its visits are to regions and its appraisal is of the regional organisation. Any action that its takes can only be in relation to a region. Responsibility for selecting trainers rests with regional general practice subcommittees, and they can take action against individuals.

The committee has always believed in the importance of the advisory and educational aspects of its visits. The issue on this occasion, however, was the failure of a regional organisation to achieve basic standards which were adopted by the committee three years before its most recent visit. We hope that those responsible for training in the North East Thames region can implement changes before the end of this year and satisfy the committee of progress so that the action proposed for February 1989 can be avoided.

DOROTHY M B WARD

Joint Committee on Postgraduate Training
for General Practice,
London SW7 1PU

SIR,—Dr Stella Lowry discusses the action taken against North East Thames (7 May, p 1320). We would like to invite Professor Pereira-Gray (chairman of the council of the Royal College of General Practitioners) to attend one of our meetings, which are held every Wednesday afternoon at the Whittington Hospital for the trainees on the University College Hospital, Middlesex, Whittington, and Royal Free vocational training schemes. As the people directly affected by the recent withdrawal by the Joint Committee on Postgraduate Training for General Practice of recognition of training in the North East Thames region and the subsequent decision by the Royal College of General Practitioners to prevent trainees in this region from sitting the MRCGP examination, except on the basis of individual assessment, we would welcome the opportunity to have the issue discussed and the situation clarified by him.

We are committed doctors practising in the inner city—supposedly the very people the government is trying to encourage to raise standards of practice in the inner city. The criticism of training in the region and the statements of the college may jeopardise our futures, whether in this region or outside it. Whatever the outcome, this controversy will have lasting and detrimental effects on the prospects of present and future trainees from this region. We have been placed undeservedly at a disadvantage in competition with our peers for jobs and will remain so for many years to come. We applaud the aims of the college to maintain and raise standards of postgraduate education, but we question whether the methods used in this episode are in the best interests of our training.

CAZ SAYER, and 20 others

Whittington Academic Centre,
London N19 5NF

SIR,—We consultants, course organisers, and trainees, having met to discuss the Royal Free vocational training scheme, would like to make public our anger at the blanket manner in which general practitioner training in the North East Thames region has been branded as inadequate.

Visitors to our scheme from the Royal College of General Practitioners in 1987 claimed that the training was satisfactory. In 1988, however, we read in the medical press that all trainees in the North East Thames region are now stigmatised as a result of a visit to five inner London practices, associated with a previous visit in 1985 to five practices in Essex, none of these practices having anything to do with our scheme.

It is difficult to think of any other professional body concerned with standards criticising a whole region (over 100 trainees) on such limited evidence. The result is that the trainees coming off our scheme and applying outside the North East Thames Region now feel they have been disadvantaged, while those responding to our present advertisement to join the scheme may consider withdrawing their application.

We protest at this unjust criticism of our training on the grounds that the problems identified have nothing to do with the Royal Free scheme. We ask that those in authority in the Joint Committee for Postgraduate Training in General Practice to rectify immediately the difficulty they have created for our trainees.

J MORRIS

J GODLEE

C DONOVAN, and 13 others

Royal Free Hospital,
London NW3 2QG

SIR,—Dr Stella Lowry argues that many answers are required as a result of the decision to withdraw recognition of training from North East Thames region (7 May, p 1320). As a former regional adviser in general practice, I support this submission and argue that the matter should be placed before the council of the BMA. There are three issues at stake which impinge on many professional interests.

Firstly, the annual report of the Joint Committee on Postgraduate Training for General Practice for 1987 gives no hint of an impending crisis in training anywhere. The section on the visitors' meeting and study day contains bland remarks about trainee assessments and the objectives of visiting. As a result of some visits, however, a working party chaired by Dr Peter Sprackling was set up to consider the procedures which may be followed after an unsatisfactory report from the visitors. No report from this working party is evident yet. It seems, therefore, that the committee's action in North East Thames region was not backed by research from the working party. One can conclude only that the committee, in describing its work, did not see the need to lay any foundation for action against a region.

Secondly, recognition or approval of training for general practice is split between two separate bodies. The approval of trainers is a statutory function of the regional general practice subcommittees and cannot be subsumed by any other authority. The approval of hospital posts is mediated through the royal colleges, and the regional postgraduate committees select those considered suitable for general practice training. The colleges do not seem to have been consulted by the committee about the changed status of their posts in North East Thames region or the manpower blight that could result.

Thirdly, the committee's power to influence training standards resides solely in its responsibilities to issue or withhold certificates on application.

These powers are exercised in relation to individual applications. There are no provisions to make prospective judgments, either for a single applicant or a group of applicants from a region. I have, therefore, to disagree with Dr Marie Campkin and others (7 May, p 1331) when they suggest that the committee could take disciplinary action against the regional and associate advisers. That this could even be suggested underlines the utter confusion that has followed the extension of powers taken by the committee so that it is now seen as virtually taking over the function of the general practice subcommittee.

The consequences of this action will be severe however the matter is resolved. The whole profession has an interest in seeing the matter settled. The evidence relied on by the committee should be submitted to the council of the BMA for rigorous review and ultimate report.

JOHN D SINSON

Leeds 17

SIR,—Dr Stella Lowry (7 May, p 1320), Dr Marie Campkin and others (7 May, p 1331), and Dr H J N Andreyev (7 May, p 1326) draw together a common theme of standards of teaching and training. I comment from the perspective of a past regional adviser in general practice who made many visits of inspection to vocational training schemes in other regions.

In general practice a statutory framework has allowed (or compelled?) the definition of criteria for those wishing to have trainees in their practices. In hospital those trainees are an indispensable component of the clinical teams, and their service contribution means that if the standards of training are inadequate withdrawal of recognition has dire consequences. The lack of formal assessment of teaching ability and commitment when consultants are appointed can, of course, lead to the situations described so graphically by Dr Andreyev. On the other hand, when training standards in general practice are perceived, rightly or wrongly, to be inadequate by the visitors the options set out by Dr Campkin and colleagues are all available. The trick is to use the right one. On one of my visits our team criticised the lack of central monitoring and control of standards in what was, admittedly, a large region. We thought that we had expressed our opinion rather wittily and were surprised and alarmed to hear that our report was considered libellous by some of the recipients.

If, as in hospitals, professional common sense and conscience dictate training standards a marvellous job will be done with occasional horrible lacunae. If, as in general practice, more formal systems are adopted the results will still be a marvellous job, with occasional horrible rows and, one hopes, fewer lacunae. I favour open accountability, which must mean defining ways of selecting acceptable trainers and adequate assessment of the training. There must also be enough sanctions to make the system work. We all perform best when we know that our efforts will be suitably judged.

D G WILSON

Cambridge CB4 1HX

Rheumatic fever

SIR,—Dr J G Bissenden (30 April, p 1215) advises of the need to prepare for a return of rheumatic fever, and our recent experience suggests that this warning is apposite.

A 37 year old woman developed a sore throat, fever, and cough with green sputum. Four days later she developed a flitting annular red rash on her legs and arms, which lasted for 48 hours. The