sense. We also urge them to correct the other anomaly and make needles alone and insulin pen injector needles available on prescription.

WILLIAM D ALEXANDER
Honorary Secretary, Medical Advisory Committee

ROBERT TATTERSALL
Chairman, Medical Advisory Committee

British Diabetic Association,
London W1M 0BD

- 1 Strathclyde Diabetes Group. Disposable or non-disposable syringes and needles for diabetics? *Br Med J* 1983;286:369-70.

- 2 Lester E, Woodroffe FJ, Grant AJ. Experience with routine re-use of plastic insulin syringes. *Br Med J* 1984;289:1498-9.
3 Allen AP, Tymms DJ, Leatherdale BA, Lloyd RS. Clinical and financial implications of a district scheme to provide plastic insulin syringes to diabetics. *Br Med J* 1986;292:1710-1.
4 Greenhough A, Cockcroft PM, Bloom A. Disposable syringes for insulin injection. *Br Med J* 1979;i:1467-8.
5 Hodge RH, Krongaard L, Sande MA, Kaiser DL. Multiple use of disposable insulin and syringe-needle units. *JAMA* 1980;244:266-7.
6 Oli JM, Gugani HC, Ojiegbe GC. Multiple use of ordinary disposable syringes for insulin injections. *Br Med J* 1982;284:236.
7 Aziz S. Recurrent use of disposable syringe-needle units in diabetic children. *Diabetes Care* 1984;7:118-20.
8 Bosquet F, Grimaldi A, Thervet F. Insulin syringe re-use. *Diabetes Care* 1986;9:310.
9 Collins BJ, Spence BK, Richardson SG, Hunter J, Nelson JK. Safety of re-using disposable plastic insulin syringes. *Lancet* 1983;i:559-61.
10 Koivisto VA, Feliq P. Is skin preparation necessary before insulin injection? *Lancet* 1978;i:1072-3.
11 Border LM, Bingham PR, Riddle MC. Traditional insulin-use practices and the incidence of bacterial contamination and infection. *Diabetes Care* 1984;7:121-7.

Polycystic ovaries in childhood

Polycystic ovaries are thought to cause 25% of cases of amenorrhoea and 85% of cases of oligomenorrhoea and can be detected in 95% of women with hirsutism.¹ Furthermore, the condition is probably the commonest cause of delayed puberty or menarche or heavy irregular periods in teenage girls.

During childhood two appearances of multiple cystic changes in the ovary may be recognised by ultrasonography. Multicystic ovaries, which contain more than six cysts with a diameter greater than 4 mm distributed throughout the ovary with no increase in stromal tissue, occur in normal girls before and during puberty.² This is the ovarian response to the nocturnal pulsatile secretion of gonadotrophin. Polycystic ovaries, which are larger than normal and contain many small cysts often arranged circumferentially with an increase in stromal tissue, are typically seen in women with the Stein-Leventhal syndrome. (It is now recognised that this classic syndrome applies only to a few obese and hirsute women who also have polycystic ovaries and menstrual irregularity.) The initial description of cystic ovaries in childhood as polycystic has caused confusion because this description probably applied to what we now call multicystic ovaries.³

Not all girls or women with polycystic ovaries have what is called the polycystic ovary disease or syndrome, although women with either polycystic ovaries or polycystic ovary syndrome often have daughters with the same condition. In addition, we have observed the daughters of several such patients whose ovarian morphology evolved from a normal multicystic pattern to a polycystic pattern during puberty.⁴ Women with polycystic ovary syndrome often report delayed menarche,⁵ and the condition probably thus begins before or during puberty. The disorder is clearly primarily an ovarian problem since polycystic ovaries have been described in girls with gonadotrophin deficiency since the age of 6 years.⁶ Only detailed longitudinal ultrasound studies will determine the prevalence and natural course of polycystic ovaries in children. But adolescent girls⁷ as well as adult women⁸ with polycystic ovaries have characteristic abnormalities of endocrine function that provide additional markers of the condition.

Since polycystic ovaries are associated with delayed puberty and irregular heavy periods the question arises of what should be done about girls with polycystic ovaries detected on ultrasonography. Our approach is pragmatic. We provide a controlled induction of puberty for a girl who is suffering

from the adverse effects of pubertal delay and for patients in the later stages of puberty with an irregular cycle we give, if necessary, the lowest dose of cyclical oestrogen and progestogen treatment compatible with good cycle control.

What is more difficult is to answer questions about the long term importance of the findings. Not all women with polycystic ovaries or indeed polycystic ovary syndrome have problems with fertility; indeed, a condition that occurs in almost a quarter of normal volunteers probably cannot be described as pathological.⁹ Yet we have evidence that polycystic ovaries associated with obesity reduce fertility.¹⁰

The ability of ultrasonographers to recognise the increase in ovarian volume and stroma and detect the typical distribution of follicles will put this syndrome increasingly before doctors treating prepubertal and adolescent girls. Children with polycystic ovaries may broadly be reassured and treated if necessary for their symptoms, but because of the effects of obesity on fertility we should warn them against excessive weight gain.

C G D BROOK

reader in paediatric endocrinology

H S JACOBS

professor of reproductive endocrinology

Endocrine Unit,
Middlesex Hospital,
London W1N 8AA

R STANHOPE

Lecturer in Paediatric Endocrinology,
Institute of Child Health,
London WC1N 1EH

- 1 Adams J, Polson DW, Franks S. Prevalence of polycystic ovaries in women with anovulation and idiopathic hirsutism. *Br Med J* 1986;293:355-9.
2 Stanhope R, Adams J, Jacobs HS, Brook CGD. Ovarian ultrasound assessment in normal children, idiopathic precocious puberty and during low dose pulsatile GnRH therapy of hypogonadotrophic hypogonadism. *Arch Dis Child* 1985;60:116-9.
3 Polhemus DW. Ovarian maturation and cyst formation in children. *Pediatrics* 1953;11:588-94.
4 Stanhope R, Adams J, Brook CGD. The evolution of polycystic ovaries in a girl with delayed menarche. *J Reprod Med* 1988 (in press).
5 Yen SSC. The polycystic ovary syndrome. *Clin Endocrinol* 1980;12:177-208.
6 Stanhope R, Adams J, Pringle JP, Jacobs HS, Brook CGD. The evolution of polycystic ovaries in a girl with hypogonadotrophic hypogonadism before puberty and during puberty induced with pulsatile gonadotrophin-releasing hormone. *Fertil Steril* 1987;47:872-5.
7 Zumoff B, Freeman R, Coupey S, Saenger P, Markowitz M, Kream J. A chronobiologic abnormality in luteinizing hormone secretion in teenage girls with the polycystic ovary syndrome. *N Engl J Med* 1983;309:1206-9.
8 Rebar R, Judd HL, Yen SSC, Rakoff J, Vandenberg G, Naftolin F. Characterisation of the inappropriate gonadotrophin secretion in polycystic ovary syndrome. *J Clin Invest* 1976;57:1320-9.
9 Adams J, Polson DW, Wadsworth J, Franks S. Polycystic ovaries—a common finding in normal women. *Lancet* 1988 (in press).
10 Eshel A, Abdulwahid NA, Armar NA, Adams J, Jacobs HS. Pulsatile luteinizing hormone releasing hormone therapy in women with polycystic ovary syndrome. *Fertil Steril* 1988 (in press).