Characterization of Infectious Hematopoietic Necrosis Virus mRNA Species Reveals a Nonvirion Rhabdovirus Protein†

GAEL KURATH AND JOANN C. LEONG*

Department of Microbiology, Oregon State University, Corvallis, Oregon 97331-3804

Received 20 July 1984/Accepted 5 October 1984

The genome RNA and six mRNA species of infectious hematopoietic necrosis virus were analyzed by denaturing gel electrophoresis. The following molecular weights were determined: genome RNA, 3.7×10^6 ; mRNA 1, 2.26×10^6 ; mRNA 2, 5.63×10^5 ; mRNA 3, 4.84×10^5 ; mRNA 4 (containing two different mRNA species), 3.00×10^5 ; and mRNA 5, 1.95×10^5 . Densitometer analyses of gels were used to calculate the molar ratios of the intracellular mRNA species: mRNA 1, 0.02; mRNA 2, 0.49; mRNA 3, 1.0; mRNA 4, 2.52; and mRNA 5, 0.41. Hybrid selection studies determined the mRNA coding assignments as follows: mRNA 1 encodes the viral polymerase, L; mRNA 2 encodes the glycoprotein, G; mRNA 3 encodes the nucleocapsid protein, N; mRNA 4 is composed of two comigrating mRNA species which encode the matrix proteins, M1 and M2; and mRNA 5 encodes a previously unrecognized viral protein which is induced in infected cells but is not present in mature virions. This nonvirion protein has been designated the NV protein.

Infectious hematopoietic necrosis virus (IHNV) is a piscine rhabdovirus of economic importance as a pathogen of salmon and trout (2, 30). This virus is responsible for increasingly devastating losses of young fish at hatcheries in western North America (12). Survivors of a hatchery epizootic of IHNV appear to be virus free until they return from the ocean and begin to spawn (2). At that time, the virus appears in many tissues of the fish and is presumably transmitted vertically via the spawning fluids. There also is substantial evidence for horizontal waterborne transmission of IHNV (28, 30). The life cycle of IHNV in relation to its carrier host is not known (23, 28, 30).

IHNV is similar to the rhabdovirus prototypes, vesicular stomatitis virus (VSV), and rabies, in that it is a bullet-shaped, enveloped virus with glycoprotein surface projections and a genome of single-stranded, negative sense RNA (3, 13, 25). The protein structure of IHNV has been reported to resemble that of rabies virus and consists of a viral polymerase (L), a surface glycoprotein (G), two matrix proteins (M1 and M2), and a nucleocapsid protein (N) (13, 19, 23). Although the proteins of IHNV have been well characterized, the viral nucleic acid species have received little attention. A system for *in vitro* transcription of IHNV has been described previously (24), but individual transcripts were not resolved or identified. Therefore, to further our understanding of the molecular biology of IHNV both in the carrier state and during lytic infections, it was first necessary to characterize the viral genome and mRNA species.

We report here the isolation of genomic RNA from purified IHNV virions and six viral mRNA species from IHNV-infected salmon cells. These molecules were resolved by electrophoresis on both glyoxal and methyl mercury gels, and the molecular weights and molar ratios present in infected cells were determined. In addition, the mRNA species were characterized by hybrid selection and *in vitro* translation studies which resulted in the identification of the viral protein coded for by each message. Our results identify

mRNA species for all the known viral proteins and, in addition, identify a message for a previously unrecognized viral protein which is present only in the infected cells.

MATERIALS AND METHODS

Cells and virus. The IHNV used in this study was isolated in 1975 from an adult steelhead trout at the Round Butte Hatchery in central Oregon. The virus was propagated in a chinook salmon embryo cell line (CHSE-214) obtained from J. L. Fryer, Department of Microbiology, Oregon State University, Corvallis. Cell monolayers were infected with IHNV at a multiplicity of infection of 0.001 and incubated at 16°C for 7 days. When radioactive labeling of viral genome RNA was desired, [5,6-³H]uridine (30 Ci/mmol; New England Nuclear Corp.) was added to 10 μ Ci/ml 24 h postinfection. The supernatant fluid was harvested and clarified of cellular debris by centrifugation at $2,400 \times g$ (4,000 rpm) in an HS-4 rotor in a Sorvall RC5 centrifuge for 10 min at 4°C. Clarified supernatant fluids were stored at -20°C, or the virions were purified on discontinuous and continuous sucrose gradients as described previously (20). The clarified supernatant contained 10^7 to 10^8 50% tissue culture infective dose units of IHNV per ml.

Preparation of viral genome RNA. Purified virions were suspended in STE (0.1 M NaCl, 20 mM Tris-hydrochloride [pH 7.4], 1 mM EDTA). Self-digested pronase and sodium dodecyl sulfate (SDS) were added to 1 mg/ml and 1% (vol/vol), respectively, and the mixture was incubated for 30 min at 37°C. RNA was isolated from disrupted virions by two extractions with STE-saturated phenol, followed by one extraction with STE-saturated chloroform-isoamyl alcohol (24:1). The final aqueous phase was made 0.3 M with potassium acetate and precipitated overnight with 2.5 volumes of ethanol. RNA was pelleted and resuspended in water, and the quantity of RNA was measured by UV absorbance at 260 nm. One liter of tissue culture fluid yielded 12 to 15 μ g of IHNV genome RNA. All solutions (excluding Tris buffers) and glassware used for RNA extraction were treated with diethyl pyrocarbonate to inactivate contaminating RNases.

Preparation of intracellular RNA. Monolayers of CHSE-214 cells were infected with IHNV at a multiplicity of

* Corresponding author.

† Oregon Agricultural Experiment Station technical paper no. 7323.

infection of 10 to establish a synchronous infection, and actinomycin D was added to 0.5 µg/ml to inhibit cellular transcription. At 6 h postinfection, [5,6-³H]uridine (30 Ci/mmol; New England Nuclear Corp.) was added to 10 µCi/ml. The infection was allowed to proceed until ca. 25% of the cell monolayer exhibited a cytopathic effect (24 to 28 h postinfection), at which time the monolayers were rinsed three times with ice-cold Tris-buffered saline. A lysing solution consisting of 0.5% SDS, 250 µg of proteinase K per ml, 0.1 M NaCl, 5 mM EDTA, and 30 mM Tris-hydrochloride (pH 7.4) was added, using 4 ml/150 cm² flask (40). The flasks then were incubated for 1 h at 37°C. Cell lysates were then pooled, and total nucleic acid was isolated by two extractions with STE-saturated phenol-chloroform-isoamyl alcohol (25:24:1). The aqueous phase was extracted with chloroform-isoamyl alcohol (24:1), brought to 0.3 M potassium acetate, and precipitated with 2.5 volumes of ethanol.

The total nucleic acid preparation was treated with DNase by incubation for 1 h at 37°C in the presence of 0.1 mg of proteinase K-treated DNase (41) per ml, 10 mM CaCl₂, 10 mM MgCl₂, 20 mM Tris-hydrochloride (pH 7.8), and 2 mM vanadyl ribonucleoside complexes (Bethesda Research Laboratories). Reactions were stopped by the addition of EDTA to 50 mM. RNA was extracted with phenol-chloroform-isoamyl alcohol (25:24:1) containing 0.1% 8-hydroxyquinoline until the dye remained yellow, indicating the complete removal of the vanadyl ribonucleoside complexes. The aqueous phase then was brought to 0.3 M potassium acetate and precipitated with 2.5 volumes of ethanol. The RNA precipitates were resuspended in water, measured spectrophotometrically for concentration and purity, and stored at -70°C.

Polyadenylated RNA was selected from the total RNA preparation by column chromatography with oligodeoxythymidylic acid-cellulose by a modified procedure of Aviv and Leder (4). RNA was suspended in 1 mM EDTA and heat denatured at 65°C for 10 min. After cooling the RNA on ice, Tris-hydrochloride (pH 7.5) and NaCl were added to 10 mM and 0.5 M, respectively. Up to 10 mg of RNA was loaded onto an 8-ml column of oligodeoxythymidylic acid-cellulose and washed extensively with binding buffer (10 mM Tris-hydrochloride [pH 7.5], 0.5 M NaCl). Bound RNA was eluted by washing with 10 mM Tris-hydrochloride (pH 7.5), and fractions containing radioactivity were pooled, brought to 0.3 M potassium acetate, and precipitated with 2.5 volumes of ethanol. Precipitated RNA was resuspended in diethyl procarbonate-treated water, quantitated, and stored at -70°C. Diethyl procarbonate was used as above to inactivate RNases in solutions and glassware.

Electrophoresis of RNA. Glyoxal treatment of RNA and subsequent electrophoresis of RNA on 1% agarose gels in 10 mM sodium-phosphate buffer (pH 7.0) was carried out as described previously (26), with the exception that the glyoxal (Fluka AG, Buchs, Switzerland) required more extensive deionization. Glyoxal was passed twice over each of five 10-ml columns of Bio-Rad AG501-X8 mixed-bed resin and stored in small portions at -20°C for single use.

Electrophoresis of RNA samples in 5 mM methyl mercuric hydroxide (Alfa Products), on 1% agarose gels containing 5 mM methyl mercuric hydroxide, was carried out as described previously (5).

Both types of agarose gels were prepared for fluorography by soaking them for 2 h with agitation in liquid autoradiography enhancer (En³Hance; New England Nuclear Corp.), followed by 2 h with agitation in water. Gels were dried between filter paper and cellophane for 2 h on a gel dryer

(Bio-Rad) and exposed to X-ray film (Kodak X-Omat AR) for 20 to 48 h at -70°C.

Electrophoresis of proteins. Proteins were analyzed by polyacrylamide gel electrophoresis on 10% polyacrylamide gels with a 4.75% stacking gel (18). Samples were boiled for 2 min in SDS denaturing buffer (62.5 mM Tris-hydrochloride [pH 6.8], 10% glycerol [wt/vol], 5% 2-mercaptoethanol, 2.3% SDS) before electrophoresis with a constant current of 20 mA. The Laemmli gel buffer system was used for gel and reservoir buffers (18).

After electrophoresis, gels were fixed for at least 1 h in 10% acetic acid-10% trichloroacetic acid-30% methanol. Gels were prepared for fluorography by treatment with liquid autoradiographic enhancer (En³Hance, New England Nuclear Corp.), dried between two pieces of cellophane, and exposed to X-ray film as above.

Densitometer analysis of autoradiograms. Autoradiographs of electrophoretic gel profiles were scanned on a Zeineh soft laser scanning densitometer (model SL-504-XL; BioMed Instruments, Inc.). Scans were printed on an Apple personal computer, using an electrophoresis reporting integrator program. Scans were photocopied, and individual peaks were cut out and weighed. Relative peak weights and mRNA molecular weights were used to calculate the molar ratio of each mRNA band with respect to mRNA band 3, which was given the arbitrary value of 1.0. The equation used was as follows: molar ratio of mRNA X = (peak weight mRNA X/peak weight mRNA 3) × (molecular weight of mRNA 3/molecular weight of mRNA X), in which mRNA X was any one of the five mRNA bands.

Cell-free translation. Polyadenylated RNA or RNA selected by hybridization to plasmid DNA was translated in vitro in a nuclease-treated rabbit reticulocyte lysate system (Bethesda Research Laboratories) as specified by the manufacturer. Reactions were carried out in 15-µl volumes for 1 h at 30°C in the presence of [³⁵S]methionine (NEG-009A; 1166.5 Ci/mmol; New England Nuclear Corp.) Newly synthesized proteins were visualized by polyacrylamide gel electrophoresis and fluorography as described above.

Hybrid selection. Hybrid selection of individual RNA species was carried out by hybridization of the polyadenylated RNA from IHNV-infected cells to plasmid DNA immobilized on nitrocellulose filters essentially as described previously (22, 33). Plasmids carrying cDNA sequences to IHNV mRNA species are listed in Table 1. The construction and characterization of these plasmids are described in an accompanying manuscript.

Plasmids, determined by RNA blot hybridization to carry sequences from each of the five translatable mRNA species, were cleaved with the restriction enzyme *Bam*HI or *Hind*III. Each linearized plasmid (10 µg) was heat denatured and spotted onto separate 3-mm squares of nitrocellulose filter paper (BA 85; Schleicher & Schuell, Inc.). The filters were dried overnight and baked at 80°C in vacuo for 2 h. Each filter was hybridized with 15 µg of polyadenylated RNA from IHNV-infected CHSE-214 cells for 6 h at 52°C under the conditions specified by Maniatis et al. (22). After hybridization the filters were washed, and the RNA was released, extracted, and precipitated as described above. Precipitated RNA was rinsed twice with 70% ethanol and translated in vitro.

RESULTS

Genome molecular weight. The genome of IHNV has been estimated by sucrose gradient sedimentation to be 38S to 40S (13, 25). To obtain a more exact size estimate, the

TABLE 1. Summary of hybrid selection results

Plasmid	mRNA band ^a selected	Protein encoded
pG480	2	G ₀
pN421	3	N
pN419	3	N
pM1163	4	M1
pM219	4	M2
pM2132	4	M2
pM2173	4	M2
pNV58	5	NV
pNV137	5	NV
pNV711	5	NV

^a Determined by mRNA blot hybridizations (17).

genome RNA was extracted from purified virions and analyzed by denaturing gel electrophoresis. This RNA migrated as a single high-molecular-weight band on both glyoxal and methyl mercury gels (Fig. 1), indicating that it was intact, genome-length RNA. Although both glyoxal and methyl mercury gels are reported to fully denature RNA (5, 26), the IHNV genome RNA ran differently in the two systems when compared with VSV genome standard RNA (the VSV genome has a molecular weight of $3.82 \times 10^6 \pm 0.14 \times 10^6$) (32). Glyoxal-treated IHNV genome RNA migrated slightly faster than glyoxal-treated VSV genome RNA, indicating a molecular weight of $3.7 \times 10^6 \pm 0.05 \times 10^6$. On methyl mercury gels the same IHNV genome RNA migrated significantly faster than the VSV genome, indicating a smaller molecular weight of ca. 3.5×10^6 to 3.6×10^6 . Although this discrepancy should be kept in mind, we have chosen to use the glyoxal gel size estimate.

mRNA molecular weights. Isolation of the viral mRNA species was carried out by infecting CHSE-214 cells with IHNV in the presence of actinomycin D and tritiated uridine. Polyadenylated RNA from IHNV-infected cells was resolved by both glyoxal and methyl mercury gel electrophoresis into six species of RNA (Fig. 1). The largest band of RNA comigrated with purified IHNV genome RNA and was presumed to be genome-length, positive- or negative-sense molecules involved in the replication process. When purified IHNV genome RNA was fractionated with oligodeoxythymidylic acid-cellulose, less than 2% of the RNA bound to the column. Therefore, the presence of this genome-length RNA in the polyadenylated RNA preparations does not indicate a true polyadenylic acid [poly(A)] tail, but perhaps some internal short poly(A) sequences, or carry-over of genome template molecules, hybridized to polyadenylated mRNA molecules during oligodeoxythymidylic acid-cellulose chromatography.

The five subgenomic polyadenylated RNA bands were designated IHNV mRNA 1 through 5, from largest to smallest (Fig. 1). In all preparations, mRNA 2, 3, 4, and 5 were relatively abundant, whereas mRNA 1 was present in minor, more variable quantities. The marker RNA species (VSV genome, 3.82×10^6 ; 28S ribosomal RNA, 1.7×10^6 ; 18S ribosomal RNA, 6.5×10^5) (27, 32) established a reproducibly linear plot of log molecular weight versus mobility for each glyoxal gel. This was used to determine the following values for the molecular weights of the IHNV mRNA species: mRNA 1, $2.26 \times 10^6 \pm 0.035 \times 10^6$; mRNA 2, $5.63 \times 10^5 \pm 0.08 \times 10^5$; mRNA 3, $4.84 \times 10^5 \pm 0.07 \times 10^5$; mRNA 4, $3.0 \times 10^5 \pm 0.05 \times 10^5$; and mRNA 5, $1.95 \times 10^5 \pm 0.03 \times 10^5$. These values were calculated as the average of 8 to 10 separate determinations including four different preparations of polyadenylated RNA.

Although the methyl mercury gels provided excellent resolution of the RNA species, in our hands the plots of log molecular weight versus mobility were not linear over the size range of the marker RNA species and therefore were not used to determine the molecular weights of unknown RNA species.

mRNA molar ratios. The molar ratios of each mRNA species produced intracellularly during IHNV infection were determined by densitometer analysis of the autoradiograms of several glyoxal gel profiles (Fig. 2). The areas under each peak and the molecular weights of the mRNA species were used to compute the following molar ratios, normalized to 1 mol of mRNA 3: mRNA 1, 0.02 ± 0.01 ; mRNA 2, 0.49 ± 0.03 ; mRNA 3, 1.00; mRNA 4, 2.52 ± 0.40 ; and mRNA 5, 0.41 ± 0.14 .

In vitro translation of mRNA. To confirm the viral mRNA character of the polyadenylated RNA preparations, the preparations were tested for the ability to direct viral protein synthesis in vitro. Figure 3 shows a polyacrylamide gel of the proteins synthesized in an mRNA-dependent cell-free translation reaction primed with 1 μ g of the polyadenylated RNA from IHNV-infected cells. Proteins were synthesized which comigrated with IHNV standard proteins N, M1, and

FIG. 1. Denaturing gel analyses of IHNV mRNA from infected cells. Polyadenylated RNA from IHNV-infected cells was analyzed by electrophoresis under denaturing conditions. (A) Glyoxal gel electrophoresis: a, ribosomal RNA markers (18S and 28S); b, IHNV genomic RNA extracted from purified virions; c, VSV genome marker RNA; d, polyadenylated RNA from IHNV-infected cells. (B) Methyl mercury gel electrophoresis: a, VSV genome marker RNA; b, IHNV genomic RNA from purified virions; c, polyadenylated RNA from IHNV infected cells; d, ribosomal marker RNA (18S and 28S). The number designations for the major species of mRNA are shown on each gel.

FIG. 2. Densitometer scan of the glyoxal gel profile of IHNV mRNA from infected cells. Gel profiles such as those shown in Fig. 1 were scanned with a soft laser scanning densitometer and printed with a personal computer, using an electrophoresis integration program. The mRNA band number corresponding to each peak is indicated at the top. The letter G indicates IHNV genomic RNA.

M2. A fourth protein was synthesized which migrated faster than the G protein and had a molecular weight of 56,000. This corresponds with the molecular weight of the unglycosylated G protein, G_0 (the molecular weight of G_0 is $55,000 \pm 3,000$; Y. Hsu, personal communication). The viral L protein was never observed in the *in vitro* translation products. A fifth protein with a molecular weight of 12,000 also was synthesized *in vitro*. Although this protein does not correspond to any of the proteins in purified virions, there is a protein of this size induced in relatively high quantities in IHNV-infected cells (Fig. 3). We have designated this the NV protein due to its nonvirion nature.

mRNA coding assignments. It was necessary to correlate which viral mRNA encoded each of the viral proteins. This was done by hybrid selection of individual mRNA species from the mixture with cloned plasmids carrying partial cDNA sequences from each viral mRNA. The construction and characterization of these clones is described elsewhere (17). *In vitro* translation of the mRNA selected by hybridization with each plasmid then identified the protein encoded by that mRNA.

In this way the protein coding assignments were made for all the mRNA species which could be translated *in vitro*. Examples for each protein are shown (Fig. 4), and a summary of the hybrid selections carried out is presented (Table 1). The plasmid which carried cDNA to mRNA band 2 hybrid selected mRNA which translated into the G_0 protein. Two plasmids which carried cDNA to band 3 selected the mRNA which coded for the N protein. Of four plasmids which carried cDNA to mRNA band 4, three selected an mRNA which coded for the M2 protein, and one selected the mRNA for the M1 protein. All three plasmids which carried cDNA to mRNA band 5 hybrid selected mRNA which translated into the small-molecular-weight NV protein.

DISCUSSION

The mRNA species of IHNV have been isolated and characterized as to molecular weights, molar ratios, and coding assignments (Table 2).

The molecular weights of the six IHNV mRNA species were used to estimate their coding capacities, after subtracting the molecular weight of a 100-residue poly(A) tail (Table

2). Our estimate of 100 residues was based on data for the poly(A) tails of VSV mRNA species (64 to 125 residues) (37) and of rabies virus mRNA species (100 to 250 residues) (14). Table 2 shows the estimated coding capacity of each IHNV mRNA, along with the actual size of the protein encoded. In each case the size of the mRNA species is more than sufficient to code for its assigned protein product. The molecular weight of 150,000 for the IHNV L protein is likely to be an underestimate by analogy with the L protein of VSV. The complete sequence of the VSV L mRNA indicates that the true molecular weight of the VSV L protein is 241,000 (38). This is significantly higher than values determined by SDS-polyacrylamide gel electrophoresis, which ranged from 160,000 to 230,000 (38). In like manner, the true molecular weight of the IHNV L protein may be higher than the value determined by gel migration and may be closer to the estimated coding capacity of mRNA 1 (Table 2). The sum of the molecular weights of the six mRNA coding sequences [excluding 100 residue poly(A) tails] is 3.9×10^6 . Thus, these six genes account for 105% of the IHNV genome length, which is 3.7×10^6 (this paper).

The molar ratios of the viral mRNA species produced intracellularly during infection are shown in Table 2. The

FIG. 3. Electrophoretic analyses of proteins translated *in vitro* from IHNV mRNA produced in infected cells. Polyadenylated RNA from IHNV-infected cells was used to prime an *in vitro* translation system, and the proteins produced were analyzed by SDS-polyacrylamide gel electrophoresis along with IHNV protein standards. Lanes: a, proteins from uninfected CHSE-214 cells; b, proteins from IHNV-infected cells; c, proteins from purified IHNV virions; d, proteins translated *in vitro* from IHNV mRNA from infected cells. The nomenclature of the IHNV proteins is indicated on the right.

FIG. 4. Electrophoretic analysis of proteins translated in vitro from individual IHNv mRNA species after hybrid selection. Protein was translated from individual mRNA species selected from polyadenylated RNA from IHNv-infected cells by hybridization with the following cDNA plasmid: a, pM1163; b, pM219; c, pN144; d, pG480; e, pNV58. The mRNA band selected by each plasmid is indicated at the top of each lane. Lane f is a marker lane of IHNv proteins translated in vitro from the total mRNA preparation.

ratios for mRNA species 2, 3, and 4 were very consistent among different preparations of polyadenylated RNA, whereas those for mRNA species 1 and 5 were more variable. The molar ratio calculated for mRNA 1 is most likely lower than the actual value due to the increased probability of degradation for such a large mRNA (17). Since the M1 and M2 mRNA species comigrate in the gel, their individual ratios cannot be distinguished. However, the combined ratio of 2.52 indicates that one or both of these mRNA species are present in molar quantities greater than the N mRNA, which was given the arbitrary molar ratio of 1.0.

The gene order on the IHNv genome has been determined by R-loop mapping to be (3')N-M1-M2-G-NV-L(5') (17). With the exception of the NV gene, this is identical to the genetic maps of VSV (3')N-NS-M-G-L(5') (1, 6) and rabies virus (3')N-M1-M2-G-L(5') (10). In VSV, transcription has been shown to proceed sequentially from the 3' to the 5' end of the genome, with no gene transcribed until the transcription of its 3' neighboring gene is complete (1, 6). In addition, studies with VSV have shown an attenuation of transcription at each gene junction, resulting in a decrease in the molar quantities of mRNA transcribed from each gene from the 3' to the 5' end of the genome (15, 42). If IHNv transcription also proceeds from the 3' to the 5' end along the genome, then the increase in the molar ratio of transcription from the

N to the M1 or M2 genes, or both, may suggest a lack of the attenuation phenomenon. However, 5' to the M2 gene sequential attenuation could occur, since the G, NV, and L mRNA species, respectively, are found in decreasing molar quantities (Table 2). Alternatively, the individual mRNA species may have unequal rates of turnover, in which case the observed molar ratios could not be used to assess sequential transcription or attenuation phenomena.

Although five bands of mRNA were resolved by electrophoresis (Fig. 1), there was not a one-to-one correspondence among these mRNA bands and the five known viral proteins. Instead, it was found that the mRNA species for two viral proteins comigrated in mRNA band 4, and mRNA band 5 encoded a previously undescribed viral protein.

The viral polymerase protein, L, was never observed as an in vitro translation product, presumably due to its large size (molecular weight, 150,000) (19) and the difficulty of translating large mRNA species in in vitro systems. However, there is evidence to indicate that mRNA 1 encodes the L protein, even though this could not be proven directly by hybrid selection and in vitro translation. First, cloning data confirms that mRNA 1 is of viral origin by the use of plasmids carrying cDNA to mRNA 1 (17). These cDNA inserts were shown to be viral sequences by hybridization with a cDNA probe to the viral genome and by heteroduplex formation with the viral genome. Second, the large molecular weight of the L protein (150,000) means that its mRNA must have a minimum molecular weight of 1.39×10^6 . Thus, mRNA 1 is the only viral mRNA of sufficient size to encode the L protein (Table 2), in addition to being the only mRNA not accounted for by other viral proteins (Table 1).

Coding assignments for the four smaller mRNA bands were made by hybrid selection of individual mRNA species, followed by in vitro translation (Fig. 4, Table 1). IHNv mRNA bands 2 and 3 encoded the viral G₀ and N proteins, respectively. Hybrid selection with different plasmids carrying cDNA to mRNA band 4 showed the synthesis of the M1 or M2 protein, but not both. Thus, it was proven that mRNA band 4 contained two comigrating mRNA species which code for the virion M1 and M2 proteins. This is analogous to the situation in VSV, in which the fourth mRNA band visible on denaturing polyacrylamide gels includes both the NS and M mRNA species (16, 37).

Thus, all five of the known virion proteins were accounted for by mRNA species 1 through 4, and the identity of mRNA 5 remained to be determined. Proof that mRNA 5 was of viral origin came again from the cloning data (17). Three plasmids were isolated which carried cDNA to mRNA 5. The cDNA inserts of these plasmids hybridized with a cDNA probe to the viral genome and also formed heteroduplexes with the viral genome. Thus, mRNA 5 is coded for by the IHNv genome and is not a cellular mRNA.

TABLE 2. Characteristics of IHNv mRNA species

IHNv mRNA	Mol wt	Molar ratio	Estimated coding capacity ^a	Protein encoded	Protein ^b mol wt
1	2.26×10^6	0.02	240,000	L	150,000
2	5.63×10^5	0.49	57,000	G ₀	55,000
3	4.48×10^5	1.0	48,500	N	40,500
4	3.00×10^5	2.52	28,700	M1 M2	22,500 17,000
5	1.95×10^5	0.41	17,400	NV	12,000

^a Calculated, assuming a poly(A) tail of 100 residues (14, 37).

^b Determined by SDS-polyacrylamide gel electrophoresis (19).

Hybrid selection in which the plasmids with cDNA to mRNA 5 were used resulted in the synthesis of a protein with a molecular weight of 12,000. Electrophoretic profiles of purified IHN virions labeled with [³⁵S]methionine confirmed that this protein is not present in mature virions (Fig. 3). However, IHNV-infected cell proteins labeled with [³⁵S]methionine did show the induction of substantial amounts of a 12,000-molecular-weight protein. Since the function of this protein remains to be determined, it was designated the NV protein, due to its nonvirion nature. The NV protein translated from hybrid-selected mRNA 5 appeared on gels as a more diffuse band compared with the intact band produced during *in vitro* translation of the whole polyadenylated RNA preparation (Fig. 3 and 4). This may be due to some alteration or preferential degradation of this small mRNA during the manipulations of hybrid selection.

Hybrid-selected preparations of individual mRNA species were sometimes contaminated with trace amounts of mRNA for other proteins, as shown by the faint N and M2 protein bands in lanes a and d (Fig. 4). This may be due to some regions of homology between these mRNA species or simply to the high relative abundance of the N and M2 mRNA species. Nevertheless, in each case a single viral protein was reproducibly the major species translated from hybrid-selected mRNA, enabling us to make conclusive coding assignments which are summarized in Table 1.

To our knowledge, the NV protein is the first nonvirion viral protein discovered in a rhabdovirus. Viral proteins synthesized in infected cells but not assembled into mature virions have been reported for positive-strand RNA viruses such as picornaviruses and togaviruses, in which the function(s) proposed for the nonvirion proteins is generally related to an RNA replicase activity (21). However, IHNV, like all rhabdoviruses, carries a viral polymerase in the virion. This polymerase is capable of carrying out the transcription of viral mRNA species in *in vitro* reactions containing lysed purified virions (24; G. Kurath and J. C. Leong, manuscript in preparation). Since these reactions would not contain any NV protein, it appears that NV protein is not necessary for viral transcription. It may be involved with viral replication or influence host cell processes, both of which occur in the infected cells.

With the exception of the NV protein, IHNV resembles VSV and rabies virus very closely with respect to its proteins, mRNA species, genome size, and genome organization. Thus, we are led to question whether there might be a nonvirion protein analogous to the IHNV NV protein associated with VSV, rabies, or both. The small molecular weight of the NV protein (12,000) and its mRNA (1.95×10^5 ; ca. 570 bases) mean that the coding sequence on the genome would be only 370 to 470 bases, depending on the length of the NV mRNA poly(A) tail (14, 37). Classical genetics with VSV mutants indicate the possible existence of six complementation groups in both the Indians and New Jersey serotypes (9, 31). This suggests that there are six viral genes rather than five. However, cloning and sequencing studies with VSV now have yielded the nucleotide sequence of the entire VSV genome, and there is no indication of a sixth gene (11, 34–36, 38). In addition, the original characterizations of the mRNA species of VSV and rabies virus did not show any mRNA species which were not accounted for by the virion proteins (7, 8, 16, 29, 37, 39).

If there is a nonvirion protein analogous to the IHNV NV protein in VSV and other rhabdoviruses, then this protein may have a general role in the rhabdovirus replication cycle. Alternatively, if VSV and rabies virus have no analog to the

NV protein, it may be that the role of this protein is somehow specific to replication in the fish cells of its host or to replication at colder temperatures. The optimum temperature for IHNV replication is 16°C, and the virus is inactivated at 20°C (30). It would be of interest to determine whether a protein analogous to the NV protein is coded for by other piscine rhabdoviruses such as spring viremia of carp virus, pike fry rhabdovirus, or viral hemorrhagic septicemia virus (30).

ACKNOWLEDGMENTS

We thank Mark Engelking for technical assistance and Carlene Pelroy for secretarial assistance.

G.K. was the recipient of an N. L. Tarter graduate fellowship award and a Mark H. Middlekauf graduate fellowship award. This work was supported by U.S. Department of Agriculture SEA grant 901-15-139 and Bonneville Power Administration grant DEA 17984BP16479.

LITERATURE CITED

1. Abraham, G., and A. K. Banerjee. 1976. Sequential transcription of the genes of vesicular stomatitis virus. *Proc. Natl. Acad. Sci. U.S.A.* 73:1504–1508.
2. Amend, D. F. 1975. Detection and transmission of infectious hematopoietic necrosis virus in rainbow trout. *J. Wildl. Dis.* 11:471–480.
3. Amend, D. F., and V. C. Chambers. 1970. Morphology of certain viruses of salmonid fishes. I. *In vitro* studies of some viruses causing hematopoietic necrosis. *J. Fish. Res. Board Can.* 27:1285–1293.
4. Aviv, H., and P. Leder. 1972. Purification of biologically active globin messenger RNA by chromatography on oligothymidylic acid-cellulose. *Proc. Natl. Acad. Sci. U.S.A.* 69:1408–1412.
5. Bailey, J. M., and N. Davidson. 1976. Methyl mercury as a reversible denaturing agent for agarose gel electrophoresis. *Anal. Biochem.* 70:75–85.
6. Ball, L. A., and C. N. White. 1976. Order of transcription of genes of vesicular stomatitis virus. *Proc. Natl. Acad. Sci. U.S.A.* 73:442–446.
7. Both, G. W., S. A. Moyer, and A. K. Banerjee. 1975. Translation and identification of the viral mRNA species isolated from subcellular fractions of vesicular stomatitis virus-infected cells. *J. Virol.* 15:1012–1019.
8. Coslett, G. D., B. P. Holloway, and J. F. Obijeski. 1980. The structural proteins of rabies virus and evidence for their synthesis from separate monocistronic RNA species. *J. Gen. Virol.* 49:161–180.
9. Flamand, A. 1980. Rhabdovirus genetics, p. 115–139. *In* D. H. L. Bishop (ed.), *Rhabdoviruses*, vol. III. CRC Press, Inc., Boca Raton, Fla.
10. Flamand, A., and J. F. Delagneau. 1978. Transcriptional mapping of rabies virus *in vivo*. *J. Virol.* 28:518–523.
11. Gallione, C. J., J. R. Greene, L. E. Iverson, and J. K. Rose. 1981. Nucleotide sequences of the mRNA's encoding vesicular stomatitis virus N and NS proteins. *J. Virol.* 39:529–535.
12. Groberg, W. J. 1983. The status of viral fish disease in the Columbia River basin, p. 1–22. *In* J. C. Leong and T. Y. Barila (ed.), *Proceedings of a workshop on viral diseases of salmonid fishes in the Columbia River basin*. Bonneville Power Administration, Portland, Oreg.
13. Hill, N. J. 1975. Physico-chemical and serological characterization of five rhabdoviruses infecting fish. *J. Gen. Virol.* 27:369–378.
14. Holloway, B. P., and J. F. Obijeski. 1980. Rabies virus-induced RNA synthesis in BHK21 cells. *J. Gen. Virol.* 49:181–195.
15. Iverson, L. E., and J. K. Rose. 1981. Localized attenuation and discontinuous synthesis during vesicular stomatitis virus transcription. *Cell* 23:477–484.
16. Knipe, D., J. K. Rose, and H. F. Lodish. 1975. Translation of individual species of vesicular stomatitis viral mRNA. *J. Virol.* 15:1004–1011.

17. Kurath, G., K. G. Ahern, G. D. Pearson, and J. C. Leong. 1985. Molecular cloning of the six mRNA species of infectious hematopoietic necrosis virus, a fish rhabdovirus, and gene order determination by R-loop mapping. *J. Virol.* **53**:469-476.
18. Laemlli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680-685.
19. Leong, J. C., Y. L. Hsu, and H. M. Engelking. 1983. Synthesis of the structural proteins of infectious hematopoietic necrosis virus, p. 61-71. *In* J. H. Crosa (ed.), *Bacterial and viral diseases of fish*. Washington Sea Grant Program, University of Washington, Seattle.
20. Leong, J. C., Y. L. Hsu, H. M. Engelking, and D. Mulcahy. 1981. Strains of infectious hematopoietic necrosis (IHN) virus may be identified by structural protein differences. *Dev. Biol. Stand.* **49**:43-55.
21. Luria, S. E., J. E. Darnell, Jr., D. Baltimore, and A. Campbell. 1978. Animal virus multiplication: the RNA viruses, p. 317-327. *In* S. E. Luria et al. (ed.), *General virology*, 3rd ed. John Wiley & Sons, Inc., New York.
22. Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. Molecular cloning, p. 329-333. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
23. McAllister, P. E., and R. R. Wagner. 1975. Structural proteins of two salmonid rhabdoviruses. *J. Virol.* **15**:733-738.
24. McAllister, P. E., and R. R. Wagner. 1977. Virion RNA polymerases of two salmonid rhabdoviruses. *J. Virol.* **22**:839-843.
25. McCain, B. B., J. L. Fryer, and K. S. Pilcher. 1974. Physicochemical properties of RNA of salmonid hematopoietic necrosis virus (Oregon strain). *Proc. Soc. Exp. Biol. Med.* **146**:630-634.
26. McMaster, G. K., and G. G. Carmichael. 1977. Analysis of single- and double-stranded nucleic acids on polyacrylamide and agarose gels by using glyoxal and acridine orange. *Proc. Natl. Acad. Sci. U.S.A.* **74**:4835-4838.
27. Metzler, D. E. 1977. *Biochemistry: the chemical reactions of living cells*, p. 927. Academic Press, Inc., New York.
28. Mulcahy, D., R. J. Pascho, and C. K. Jené. 1983. Detection of infectious hematopoietic necrosis virus in river water and demonstration of waterborne transmission. *J. Fish Dis.* **6**:321-330.
29. Pennica, D., B. P. Holloway, J. T. Heyward, and J. F. Obijeski. 1980. *In vitro* translation of rabies virus mRNAs. *Virology* **103**:517-521.
30. Pilcher, K. S., and J. L. Fryer. 1978. Viral diseases of fish: a review through 1978, part 2. *Crit. Rev. Microbiol.* **7**:287-364.
31. Pringle, C. R., and J. F. Szilagy. 1980. Gene assignment and complementation group, p. 141-161. *In* D. H. L. Bishop (ed.), *Rhabdoviruses*, vol. II. CRC Press, Inc., Boca Raton, Fla.
32. Repik, P., and D. H. L. Bishop. 1973. Determination of the molecular weight of animal RNA virus genomes by nuclease digestions. I. Vesicular stomatitis virus and its defective T particle. *Virology* **12**:969-983.
33. Ricciardi, R. P., J. S. Miller, and B. E. Roberts. 1979. Purification and mapping of specific mRNAs by hybrid-selection and cell-free translation. *Proc. Natl. Acad. Sci. U.S.A.* **76**:4927-4931.
34. Rose, J. K. 1980. Complete intergenic and flanking gene sequences from the genome of vesicular stomatitis virus. *Cell* **19**:415-421.
35. Rose, J. K., and C. K. Gallione. 1981. Nucleotide sequences of the mRNA's encoding the vesicular stomatitis virus G and M proteins determined from cDNA clones containing the complete coding regions. *J. Virol.* **39**:519-528.
36. Rose, J. K., and K. Iverson. 1979. Nucleotide sequences from the 3'-ends of vesicular stomatitis virus mRNA's as determined from cloned DNA. *J. Virol.* **32**:404-411.
37. Rose, J. K., and D. Knipe. 1975. Nucleotide sequence complexities, molecular weights, and poly(A) content of the vesicular stomatitis virus mRNA species. *J. Virol.* **15**:994-1003.
38. Shubert, M., G. G. Harmison, and E. Meier. 1984. Primary structure of the vesicular stomatitis virus polymerase (L) gene: evidence for a high frequency of mutations. *J. Virol.* **51**:505-514.
39. Sokol, F., D. Stanček, and H. Koprowski. 1971. Structural proteins of rabies virus. *J. Virol.* **7**:241-249.
40. Thomas P. 1983. Hybridization of denatured RNA transferred or dotted to nitrocellulose paper. *Methods Enzymol.* **100**:255-266.
41. Tullis, R. H., and H. Rubin. 1980. Calcium protects DNase I from proteinase K: a new method for the removal of contaminating RNase from DNase I. *Anal. Biochem.* **107**:260-264.
42. Villareal, L. P., M. Breindl, J. J. Holland. 1976. Determination of molar ratios of vesicular stomatitis virus induced RNA species in BHK21 cells. *Biochemistry* **15**:1663-1667.