Published in final edited form as:

Brain Res. 2008 July 7; 1218: 77–86. doi:10.1016/j.brainres.2008.04.007.

Effects of glucose and glutamine concentration in the formulation of the artificial cerebrospinal fluid (ACSF)

Je Hi An, Yuzhuo Su, Thomas Radman, and Marom Bikson¹

Department of Biomedical Engineering, City College of New York, New York, NY 10031 USA

Abstract

The composition of the ACSF is fundamental in controlling the extracellular environment of the brain slice preparation. ‘Typical’ formulations lack amino acids and contain a higher concentration of glucose (10 mM) than in the cerebrospinal fluid (0.47 – 4.4 mM). We examined the effects of different concentrations of glutamine, the most abundant amino acid in the CSF, and glucose on rat hippocampal slice physiology. Bipolar paired-pulse stimulation was applied to the Schaffer collaterals and population spikes were monitored in the CA1 pyramidal layer. Addition of glutamine (0.5 mM) to slices superfused with 10 mM of glucose did not enhance population spike amplitude. Higher concentration of glutamine (2–5 mM) resulted in spreading-depression. Decreasing glucose concentration from 10 mM to 5 mM, in the absence of glutamine, attenuated population spikes. Decreasing glucose to 2 mM, in the absence of glutamine, suppressed evoked population spikes. Superfusing brain slices with ACSF containing ‘physiological’ concentrations of both glucose (2 mM) and glutamine (0.5 mM) similarly suppressed population spikes. In separate experiments, during high-K⁺ induced epileptiform activity, glutamine (0.5 mM) did not affect the burst duration, frequency or waveform. These results suggest that the concentration of glucose in ACSF should conservatively be 10 mM in order to maximize paired-pulse population responses while the presence of physiological concentration of glutamine (0.5 mM) has minimal effects on paired-pulse responses and high-K⁺ induced epileptiform activity. These results are discussed in the context of fundamental differences between *in vitro* brain slice superfusion and *in vivo* brain perfusion.

Section

Neurophysiology; neuropharmacology

Keywords

glutamine; glucose; ACSF; superfusion; blood-brain barrier

1. Introduction

The composition of artificial cerebrospinal fluid (ACSF) is fundamental in controlling the extracellular environment of the *in vitro* brain slice preparation and will affect the ‘baseline’

¹To whom all correspondence and/or reprint requests should be addressed. T-403B, Steinmann Hall, Department of Biomedical Engineering, City College of New York, 140th St. and Convent Avenue, New York, NY 10031 USA, E-mail: bikson@ccny.cuny.edu, Tel: 212-650-6791, Fax: 212-650-6727.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

levels of neuronal activity (e.g. resting membrane potential, action potential rate, transporter activity, intracellular ion content), metabolism (e.g. energy-related parameters, oxygen and glucose consumption), and protein synthesis (Jefferys 1995). Despite the ubiquity and established utility of the brain slice preparation, there remain fundamental questions about ‘appropriate’ ACSF formulation and significant variations in ACSF composition across discrete slice studies; moreover, typical ACSF formulation deviate fundamentally from *in vivo* brain organic concentrations.

In the brain, glucose is a key energy substrate due to its high plasma concentration and abundant glucose transporters across the blood-brain-barrier (Dienel and Hertz 2001). In ‘typical’ ACSF formulations, glucose concentration is 10–11 mM (in some studies up to 25 mM; Christie and Jahr, 2006). These values are significantly higher than the glucose range reported *in vivo*, 0.47–4.4 mM (Silver and Erecinska, 1994; McNay and Gold, 1999; Kirchner et al., 2006). *In vitro*, the minimum glucose concentration required to maximize *specific* electrophysiological indicators, including population spikes, has been reported to vary in the range of 1 to 10 mM (Cox and Bachelard, 1982; Schurr et al., 1989; Rosen and Andrew 1991; Yuan et al., 2004; Kirchner et al. 2006). Increasing glucose concentration (> 5 mM) exerts a neuroprotective role *in vitro* (Schurr et al., 1989; Schurr 1999a; Schurr et al., 1999b; Cater et al., 2003). The ‘appropriate’ glucose concentration in ACSF thus remains controversial.

In addition to glucose, the cerebrospinal fluid (CSF) contains numerous amino acids of which glutamine is the most abundant, in the range of 0.4–0.8 mM (Ames and Nesbett, 1981; Schiff et al., 1985; Szerb and O’Regan, 1985; Schurr et al., 1987; Kapetanovic et al., 1993; Nishimura et al., 1995; Zheng et al., 2000; Tani et al., 2007). Glutamine serves as a major precursor of neurotransmitters including glutamate and GABA (Szerb and O’Regan 1985; Battaglioli and Martin 1991; Berg-Johnsen et al., 1993; Bacci et al., 2002; Bak et al., 2006; Kam and Nicoll 2007; Tani et al., 2007), has been found to induce enzyme activity (Baudry et al., 1988) and may serve as energy substrate (Bradford et al., 1978; Bacci et al., 2002; Cater et al., 2003). However, glutamine is omitted in ‘typical’ ACSF formulations. Superfusion with glutamine-free ACSF leads to depletion of brain slice glutamine content (and related neurotransmitter content), which can be (partially) reversed by addition of glutamine to the ACSF (Kapetanovic et al., 1993; Battaglioli and Martin 2005; Tani et al., 2007). Experiments with fluorocitrate suggest that inhibition of glutamine synthesis disrupts brain slice function (Berg-Johnsen et al. 1993). The effects of ACSF glutamine on slice physiology appear system and experimental protocol specific (Schiff et al., 1985; Kan and Nicoll 2007; Tani et al., 2007).

To better understand the role of glucose and glutamine in ACSF formulation, we examined the effects of glutamine and different concentrations of glucose on evoked paired population responses in rat hippocampal slices and spontaneous epileptiform activity. The goal of this study was to determine what concentrations of ACSF glucose and glutamine are required to support slice function; paired-population responses and epileptiform activity are sensitive to a broad range of excitability and excitatory/inhibitory synaptic changes. In addition to extending previous work examining the effects of ACSF glutamine (in the presence of physiologically ‘high’ glucose concentration) or only reducing ACSF glucose concentration (in the absence of glutamine), we specially investigated the concurrent restoration of *both* ACSF glucose and glutamine to *in vivo* levels.

As a result of physical slicing, inherent connectivity/morphological, metabolic, and neurophysiological differences exist between *in vivo* and *in vitro* brain tissue (Elliot 1969; Dingledine et al., 1980; Langmoen and Andersen 1981; Lipton and Whittingham 1984; Schwartzkroin et al., 1998; Schurr et al., 1999b; Zheng et al., 2000; Novotny et al., 2001). There are fundamental differences between brain slice superfusion and brain perfusion, including blood, CSF, and interstitial flow (Fig 1). Because of these inherent differences, ACSF

formulations closer to CSF concentrations may not necessarily result in more 'relevant' *in vitro* model systems (Kirchner et al 2006). In general, the relevance of data collected using brain slices to the *in vivo* situation is dependent on how well 'important' *in vivo* characteristics are preserved *in vitro*; which *in vivo* characteristics are considered important depends on the questions asked (Alger et al., 1984). The conditions of brain slice preparation and maintenance, including the formulation of the slice superfusate (ACSF), are thus critical to the valid interpretation of *in vitro* data. A necessary first step toward addressing this complex topic is to quantify what effects compounds normally present in CSF, including glucose and glutamine, have on brain slice function.

2. Results

2.1 Effect of ACSF glutamine and glucose concentration of evoked population spikes

Paired-pulse evoked population spikes provide a gross indicator of overall slice excitability, including synaptic function. The role of glutamine in ACSF was tested by monitoring its effect on population spike amplitude. Under both control (10 mM glucose, 0 mM glutamine) and glutamine (10 mM glucose, 0.5 mM glutamine) conditions, population spike amplitude increased gradually over the course of most experiments. In this and other tests, the role of drugs over time was therefore compared to control drift in population spike amplitude. There was no difference in normalized population spike amplitude at 50 minutes, between control ($n = 7$) and glutamine-treated ($n = 18$; 0.5 mM) slices (control pulse 1 (P1), 192 ± 34 %; glutamine-treated P1, 155 ± 14 %; control pulse 2 (P2), 127 ± 11 %; glutamine-treated P2, 160 ± 83 %; Fig 2B). There was no significant difference in paired-pulse ratio for either facilitation or depression cases (control paired-pulse facilitation (PPF), 85 ± 20 %; glutamine-treated PPF, 99 ± 8 %; control paired-pulse depression (PPD), 59 ± 11 %; glutamine-treated PPD, 120 ± 33 %; Fig 2C). Lower concentration of glutamine (0.2 mM) similarly had no effect on population spike amplitude ($n = 3$; not shown). Higher concentration of glutamine (2–5 mM) induced spreading-depression in 5 of 6 slices tested (Fig 3), in the remaining slice the population spike amplitude attenuated over time.

The concentration of glucose in ACSF was varied while monitoring evoked population spike amplitude. Reduction of glucose from 10 mM to 5 mM ($n = 6$) resulted in a significant decrease in the pulse 1, but not pulse 2, population spike amplitude with respect to control cases (control P1, 192 ± 34 %; 5 mM glucose-treated P1, 92 ± 21 %; control P2, 127 ± 11 %; 5 mM glucose-treated P2, 113 ± 28 %; Fig 4B). There was no significant difference in paired-pulse ratio for either facilitation or depression cases (control PPF, 85 ± 20 %; 5 mM glucose-treated PPF, 113 ± 72 %; control PPD, 59 ± 11 %; 5 mM glucose-treated PPD, 126 ± 25 %; Fig 4C). Reduction of glucose to 2 mM significantly attenuated P1 population spike amplitude such that within 30 minutes of low-glucose expose, P1 spikes were reduced to 41 ± 21 % of baseline, while at 30 minutes P2 population amplitude was not reduced significantly (78 ± 28 %). At 50 minutes, both pulses were significantly attenuated relative to control conditions (2 mM glucose-treated P1, 34 ± 23 %; 2 mM glucose-treated P2, 62 ± 26 %; Fig 5B). These results are consistent with previous studies (Fan et al. 1988; Schurr 1989) using single and paired-pulses where 'synaptic silence' was reported when glucose concentration was reduced below 2 mM.

After confirming that 2 mM glucose, in the absence of glutamine, is insufficient to maintain slice function, we investigated if the combination of 2 mM glucose and 0.5 mM glutamine could maintain or restore slice function. In this case, *both* glucose and glutamine were concurrently resorted to representative *in vivo* levels. We found that in the presence of glutamine ($n = 5$, 0.5 mM), reduction of glucose from 10 to 2 mM still caused disruption in the population response after 40 min exposure (P1, 134 ± 23 % to 39 ± 16 %, $p < 0.05$; P2, 158 ± 27 % to 69 ± 28 %, $p < 0.05$; Fig 5C). These results indicate that glutamine (0.5 mM) cannot reverse the disruption in gross slice function resulting from glucose reduction to 2 mM.

We further tested in 'moderate' reductions in ACSF glucose (to 5 mM) would induce a (metabolic) dependence on ACSF glutamine. In the presence of baseline 5 mM glucose, addition of 1 mM glutamine had no significant effect of paired-pulse evoked orthodromic responses (not shown); spreading-depression was not observed in any slices after addition of 1 mM glutamine.

2.2 Effect of ACSF glutamine on high-K⁺ induced epileptiform activity

We next investigated the effect of glutamine in high-K⁺ ACSF, on slice function including cell excitability or synaptic function. Effects of ACSF glutamine on excitability of synaptic function would be expected to induce a change in high-K⁺ epileptiform waveform (Jensen and Yaari 1988; Leschinger et al. 1993; Bikson et al. 2002). In addition, we hypothesized that the increased neuronal activity/metabolic demand associated with epileptiform activity may induce a dependence on ACSF glutamine. However, addition of glutamine (0.5 mM; n = 4) did not affect high-K⁺ induced epileptiform activity compared to baseline burst duration ($116 \pm 9\%$), burst amplitude ($114 \pm 11\%$), or burst frequency ($89 \pm 8\%$; Fig 6).

3. Discussion

Despite the acceptance of the brain slice as a valuable neuroscience research tool, there are significant differences in the ACSF formulation used across laboratories, and little consensus on 'appropriate' formulations, particularly concerning organics. The implications of the results of the present study, to this complex problem, are discussed.

3.1. Role of ACSF glutamine

Our experiments indicate that addition of glutamine to ACSF formulation does not significantly affect neuronal excitability or synaptic function (reflected in evoked population spike response) during low-frequency stimulation of hippocampal brain slices over 1–2 hours of recording time. Considering that *in vivo* clearance of glutamine from brain to endothelium or from CSF to choroids plexus, is more rapid than the influx from blood (Xiang et al., 2003), our results corroborate with a *net* production of glutamine by the brain (tissue) and efflux to the blood/ACSF. Under normal conditions, sufficient glutamine may be produced by brain slices (Berg-Johnsen et al. 1993), and excess glutamine diffused by the ACSF. Kapetanovic et al. 1993 reported time-dependent loss of glutamine in unstimulated slices. Thus, even in the absence of ACSF glutamine, brain slice neuronal glutamine pools may be adequately replenished, at least under the conditions tested in the present study. Considering that 0.5 mM ACSF glutamine results in overshoot of basal glutamine level (Kapetanovic et al., 1993), 'pathological' increases in ACSF glutamine concentration (2–5 mM) may *reverse* the direction of glutamine diffusion resulting in excessive interstitial glutamine accumulation, interference with glutamine-glutamate cycling, and/or saturation of amino acid transporters, ultimately leading to the spreading-depression observed in the present study and previously in cortical slices (Tani et al. 2007). Spreading-depression was not noted during 4 mM glutamine superfusion at room temperature (Kam and Nicoll 2007).

The glutamate-glutamine cycle is considered integral to replenishing the neurotransmitter pool (Pfrieger and Barres 1996; Tsacopoulos and Magistretti 1996; Rothman et al., 1999; Bacci et al., 1999; Bacci et al., 2002; Kam and Nicoll, 2007); recent reports have highlighted remaining questions regarding the kinetics of glutamine regulation (Winkler et al., 2000; Blin et al., 2002; Hasegawa et al., 2006; Masson et al., 2006; Conti and Melone, 2006; Kam and Nicoll 2007). Brain slice studies examining glutamine regulation have typically used relatively 'high' baseline glucose levels (compared to *in vivo* levels). Moreover, because of fundamental differences between slice and brain perfusion (Fig 1), and related differences in metabolic kinetics, insights from *in vitro* studies must be cautiously applied to the *in vivo* situation. Rather,

the focus of the present report was characterizing under what conditions ACSF glutamine would modulate the activity of brain slices and hence play a role in *in vitro* experimental design and interpretation.

The results of the present study indicate that omission of glutamine from ACSF formulation will not *necessarily* confound interpretation of results from brain slice studies. However, under conditions of prolonged (>2 hours) incubation or during intense neuronal activity, glutamine pools in the slice may be depleted resulting in functional uptake of glutamine, if it is present, from the ACSF (Schiff et al., 1985; Szerb 1985). The high-K⁺ activity induced in the present report, however, did not lead to such dependence, at least as manifest in epileptiform waveform over 40 minutes. Glutamine (0.1 to 5 mM) has been shown to protect hippocampal slices against hypoxia while 10 mM glutamine aggravated loss of synaptic function (Schurr et al., 1987). Kam and Nicoll (2007) reported an effect on hippocampal slice synaptic function only at relatively high (4 mM) glutamine levels and after >4 hours; they verified this was not limited by slice diffusion kinetics (Fig 1) by also testing dissociated neurons. In cortical slices, glutamine (<1 mM) increased glutamine content and evoked potentials in both normal and hyper-excitable tissue (Tani et al., 2007). Consistent with our findings in hippocampal slices, Tani et al. (2007) did not observe spreading-depression in normal cortical slices after addition of physiological (<1 mM) glutamine to ACSF; however, ≥ 0.5 mM glutamine promoted spreading-depression in hyper-excitable ‘under-cut’ cortical slices. In the retinal slice preparation, glutamine (0.5–5 mM) modulates protein synthesis and electrophysiological responses after ~2 hours (Parks et al., 1976; Ames and Nesbett 1981; Ames and Masland 1990). Glutamine-containing ACSF (0.6 mM) increased the spontaneous firing rate of most neurons in the ventromedial nucleus of hypothalamic slices (Nishimura et al., 1995). Thus the importance of glutamine in ACSF formulation may depend on the specific experimental paradigm used (‘what question is asked’).

3.2 Role of ACSF glucose

The attenuation of excitability following glucose reduction is not attributed to osmotic challenge because of the scale of changes and because decreases in glucose/osmolarity would be expected to increase excitability (see Experimental Procedure).

We report that reduction of glucose from 10 mM to 5 mM attenuated pulse 1 after >40 minutes, but there was no significant change in pulse 2 amplitude or paired-pulse ratio even at 50 minutes. Characterizing the dentate region of Guinea-Pig slices, Cox and Bachelard (1982) observed significant decrease in *unpaired* population responses after glucose reduction from 10 to 5 mM. Paired-pulse *ratio*, in the dentate region of rat hippocampal slices, was not compromised by reduction in glucose to from 5 to 1 mM or 10 to 2 mM (Kirchner et al. 2006). We found that reduction in glucose to 2 mM significantly reduced pulse 1, but did not significantly change pulse ratio. Fan et al. (1988) and Schurr et al. (1989) reported a decrease in spike size with <2 mM ACSF glucose using single and paired-pulses, but did not examine paired-pulse ratios. Our results are thus consistent with, and reconcile, these previous studies. Across slice neurophysiological studies, 10 mM ACSF glucose maximizes responses to ‘physiological’ stimuli (and no *attenuation* was reported after *increasing* ACSF glucose from 5 to 10 mM); thus 10 mM ACSF glucose may be a “conservative” baseline concentration to maximize typical slice neurophysiological responses. This result serves as a rational guideline for ACSF formulation, but does not necessarily inform *in vivo* glucose regulation mechanisms (see below).

Less than 2 mM ACSF glucose fails to support specific brain slice function indicators. However, *in vivo* extracellular glucose concentration is in the range of 0.47–4.4 mM (Silver and Erecinska, 1994; McNay and Gold, 1999; Kirchner et al., 2006), thus 2 mM glucose is within the physiological CSF range. One plausible explanation for this considerable

discrepancy is that the distance for metabolite diffusion in brain slice (set by slice thickness; $>300\ \mu\text{M}$) is generally higher than the distance *in vivo* (set by intercapillary distance, $\sim 50\ \mu\text{m}$, LaManna et al., 2004). The ‘U’ shaped concentration profile of metabolites in a slice (Fujii 1982; Schiff and Somjen 1985) may require that glucose concentration at the *surface* of the slice (set by ACSF formulation) be maintained above physiological concentration to ensure adequate glucose delivery at the slice core. Important from the perspective of ACSF formulation, the relatively high glucose concentration near the slice surface may not adversely affect neuronal function in this region; increasing glucose from 5 to 10 mM does not *decrease* overall slice excitability.

3.3 ACSF formulation and the brain slice preparation: experimental design

As illustrated in the cases of hyperglycemic aggravation of neuronal damage (Schurr et al., 1999b; Mehta 2003; Payne et al., 2003) and hypoglycemic associated seizures (Kirchner et al. 2006), the (systemic) effects of compounds *in vivo* may differ categorically from their effects on brain slices. Particularly in the case of metabolites, such differences may be attributed to fundamental differences in neuronal microenvironment regulation (Fig 1). The results of the present study directly inform brain slice experiment design and interpretation, rather than *in vivo* metabolic dynamics.

The above discussion reinforces the need to consider both fundamental differences between *in vivo* and *in vitro* conditions and specific brain slice experimental paradigms (‘what question is being asked’) when determining ACSF formulation. Considering the significant differences between *in vivo* and *in vitro* conditions (Fig 1), disparities in metabolic activity and chemical microenvironment are expected; moreover, the role of glucose and glutamine concentrations in ACSF will presumably vary depending on the experimental conditions including high frequency stimulation, bath temperature, protocol time ($> 2\ \text{hrs}$), slice thickness, animal species (Cox and Bachelard., 1982; Schiff et al., 1985; Newman et al., 1991; Bacci et al., 2002). Our results support ACSF formulation approaches that methodically deviate from CSF levels to adequately maintain brain slice function. ACSF glucose levels may ‘conservatively’ be maintained at 10 mM (presumably reflecting limited slice diffusion), while varied benefits of supplementing (0.5 to 4 mM) glutamine may be balanced against increased risk of spreading-depression (presumably reflecting reversing diffusion gradients).

4. Experimental procedure

Transverse hippocampal slices ($350\ \mu\text{m}$) were prepared from male Sprague-Dawley rats (125–150 g); anaesthetized with intraperitoneal ketamine ($7.4\ \text{mg kg}^{-1}$) and xylazine ($0.7\ \text{mg kg}^{-1}$) and killed by cervical dislocation (Radman et al., 2007). A total of 30 animals have been used and no more than 3 slices from the same animal were used for any experimental condition. The slices were stored in a holding chamber submerged in “normal” ACSF consisting of (mM): 125 NaCl, 26 NaHCO_3 , 3 KCl, 1.6 CaCl_2 , 1.5 MgSO_4 , 1.25 NaH_2PO_4 , and 10 glucose, bubbled with a mixture of 95% O_2 –5% CO_2 . After $>60\ \text{min}$, slices were transferred to an interface recording chamber at 34°C .

Extracellular recordings of population responses in the CA1 pyramidal cell region were made using glass micropipettes ($2\text{--}8\ \text{M}\Omega$) filled with 150 mM NaCl. Bipolar platinum-iridium stimulating electrodes (Merrill et al. 2005) were placed in Schaffer collaterals (orthodromic stimulation) and paired-pulses (30 ms interpulse delay, 0.2 ms pulse duration) were evoked once every 2 minutes. Slices were stimulated with intensity $< 500\ \mu\text{A}$ (2200 and 2300; A–M Systems, Carlsborg, WA, USA) to produce first population spike amplitude between 1–15 mV and adjusted to 25–75 % of the maximum response. Unless otherwise stated, superfusate changes were applied (at $t = 0\ \text{min}$) for 40 minutes after a baseline was established. Baseline was established as $<\pm 10\%$ change in first population spike amplitude over 12 minutes. For

each experimental trial, the first (P1) and second (P2) population spike amplitudes were normalized to the respective averages during the baseline period. PPF was determined as P2/P1 greater than 1, while PPD was determined as P2/P1 less than 1. Paired-pulse ratios (P2/P1) were normalized to the averages during the baseline period of each trial. Results are reported as normalized percentage $\pm$ standard error; n=number of slices. Student's t-test was performed compared to the normalized control values at t = 50 min unless otherwise indicated and the null-hypothesis was rejected if $p < 0.05$. We identified outliers using boxplots (Gurcan et al., 1997; Gruetter et al., 2001).

Extracellular recording of epileptiform activity in the CA1 pyramidal cell region were made using glass micropipettes (2–8 M Ω) filled with 150 mM NaCl. After 30 minutes of perfusion with 'high K⁺' ACSF (125 NaCl, 26 NaHCO₃, 8 KCl, 1.0 CaCl₂, 1.5 MgSO₄, 1.25 NaH₂PO₄, and 10 glucose), complete mechanical lesions were made, in the CA2 region, across the Schaffer collateral pathway (Jensen and Yaari 1988).

The osmolality of ACSF was measured using a Vapro-5520 vapor pressure osmometer (Wescor, Logan, UT, USA). The osmotic difference between high-K⁺ glutamine absent (308 ± 1 mOsm/L) and high-K⁺ plus 0.5 mM glutamine (310 ± 1 mOsm/L) is significantly less than that previously shown to modulate epileptiform waveform (Andrew et al., 1989; Traynelis and Dingledine 1989; Rosen and Andrew 1991; Saly and Andrew 1993). The osmotic differences between "normal" ACSF (295 ± 2 mOsm/L), ACSF plus 0.5 mM glutamine (295 ± 3 mOsm/L), and ACSF with 2 mM Glucose (291 ± 2 mOsm/L) are not expected to significantly modulate excitability (Rosen and Andrew 1991; Saly and Andrew 1993; Azouz et al., 1997).

All signals were amplified, and low-pass filtered (1–10 kHz) with an Axoclamp-2B (Axon Instruments, Union City, CA, USA) and FLA-01 amplifiers (Cygnus Technology, Delaware Water Gap, PA, USA); then digitized and processed using a Power 1401 and Signal software (CED Cambridge Electronic Design, Cambridge, UK).

It is important to note that neither population spike amplitude nor epileptiform waveform is not necessarily indicative of a "healthier" slice; however, these extracellular field waveforms are established and quantitative indicators of gross brain slice function. Changes in neuronal excitability, metabolism, or excitatory or inhibitory synaptic function (including pre- or post-synaptic effects) would be manifest in these signals (Schwechter et al., 2003); specifically using a 30 ms interpulse delay and the high-K⁺ model (Korn et al., 1987; Traynelis and Dingledine 1988; Patrylo et al., 1994; Leschinger et al., 1993). These metrics are therefore used in the context of this study to access any effects of ACSF organics on gross slice function. In addition, monitoring of evoked (paired) population spikes and epileptiform activity is ubiquitous across slice studies on normal and pathological brain function, such that the results of this paper using these metrics directly impact the appropriate design of such studies.

All chemicals were obtained from Sigma (St. Louis, MO, USA).

Acknowledgments

We thank Phillip Hahn for discussions on general ACSF formulation.

References

- Alger, BE.; Dhanjal, SS.; Dingledine, R.; Garthwaite, J.; Henderson, G.; King, GL.; Lipton, P.; North, A.; Schwatzkroin, PA.; Sears, TA.; Segal, M.; Whittingham, TS.; Williams, J. Appendix: Brain Slice Methods. In: Dingledine, R., editor. *Brain Slices*. New York: Plenum Press; 1984. p. 381-437.
- Ames A 3rd, Nesbett FB. In vitro retina as an experimental model of the central nervous system. *J Neurochem* 1981;37:867–877. [PubMed: 7320727]

- Ames, A.; Masland, RH. The rabbit retina *in vitro*. In: Jahnsen, H., editor. Preparations of Vertebrate Central Nervous System In Vitro. Chichester: John Wiley; 1990. p. 183-202.
- Andrew RD, Fagan M, Ballyk BA, Rosen AS. Seizure susceptibility and the osmotic state. *Brain Res* 1989;498:175–180. [PubMed: 2790471]
- Azouz R, Alroy G, Yaari Y. Modulation of endogenous firing patterns by osmolarity in rat hippocampal neurones. *J Physiol* 1991;502:175–187. [PubMed: 9234205]
- Bacci A, Verderio C, Pravettoni E, Matteoli M. The role of glial cells in synaptic function. *Phil Trans R Soc Lond B* 1999;354:403–409. [PubMed: 10212490]
- Bacci A, Sancini G, Verderio C, Armano S, Pravettoni E, Fesce R, Franceschetti S, Matteoli M. Block of glutamate-glutamine cycle between astrocytes and neurons inhibits epileptiform activity in hippocampus. *J Neurophysiol* 2002;88:2302–2310. [PubMed: 12424271]
- Bak LK, Schousboe A, Waagepetersen HS. The glutamate/GABA-glutamine cycle: aspects of transport, neurotransmitter homeostasis and ammonia transfer. *J Neurochem* 2006;98:641–653. [PubMed: 16787421]
- Battaglioli G, Martin DL. GABA synthesis in brain slices is dependent on glutamine produced in astrocytes. *Neurochem. Res* 1991;16:151–156. [PubMed: 1881516]
- Baudry M, Shahi K, Gall C. Induction of ornithine decarboxylase in adult rat hippocampal slices. *Mol Brain Res* 1988;4:313–318.
- Berg-Johnsen J, Paulsen RE, Fonnum F, Langmoen IA. Changes in evoked potentials and amino acid contents during fluorocitrate action studied in rat hippocampal cortex. *Exp Brain Res* 1993;96:241–246. [PubMed: 7903642]
- Bikson M, Id Bihi R, Vreugdenhil M, Kohling R, Fox JE, Jefferys JG. Quinine suppresses extracellular potassium transients and ictal epileptiform activity without decreasing neuronal excitability in vitro. *Neuroscience* 2002;115:251–261. [PubMed: 12401338]
- Blin M, Crusio WE, Hevor T, Cloix JF. Chronic inhibition of glutamine synthetase is not associated with impairment of learning and memory in mice. *Brain Res Bull* 2002;57:11–15. [PubMed: 11827732]
- Bradford HF, Ward HK, Thomas AJ. Glutamine—a major substrate for nerve endings. *J Neurochem* 1978;30:1453–1459. [PubMed: 670985]
- Cater HL, Chandratheva A, Benham CD, Morrison B III, Sundstrom LE. Lactate and glucose as energy substrates during, and after, oxygen deprivation in rat hippocampal acute and cultured slices. *J Neurochem* 2003;37:1381–1390. [PubMed: 14713294]
- Christie JM, Jahr CE. Multivesicular release at Schaffer collateral-CA1 hippocampal synapses. *J Neurosci* 2006;26:210–216. [PubMed: 16399689]
- Conti F, Melone M. The glutamine commute: lost in the tube? *Neurochem Int* 2006;48:459–464. [PubMed: 16517023]
- Cox DW, Bachelard HS. Attenuation of evoked field potentials from dentate granule cells by low glucose, pyruvate + malate, and sodium fluoride. *Brain Res* 1982;239:527–534. [PubMed: 6284306]
- Dienel GA, Hertz L. Glucose and lactate metabolism during brain activation. *J Neurosci Res* 2001;66:824–838. [PubMed: 11746408]
- Dingledine R, Dodd J, Kelly JS. The in vitro brain slice as a useful neurophysiological preparation for intracellular recording. *J Neurosci Methods* 1980;4:323–362. [PubMed: 6106092]
- Elliot, KA. The Use of Brain Slices. In: Lajtha, A., editor. *Handbook of Neurochemistry*. Vol II. New York: Plenum Press; 1969. p. 103-114.
- Gruetter R, Seaquist ER, Ugurbil K. A mathematical model of compartmentalized neurotransmitter metabolism in the human brain. *Am J Physiol Endocrinol Metab* 2001;281:E100–E112. [PubMed: 11404227]
- Gurcan MN, Yardimci Y, Cetin AE, Ansari R. Automated detection and enhancement of microcalcifications in mammograms using nonlinear subband decomposition. *IEEE Conference on acoustics, speech, and signal processing* 1997;4:3069–3072.
- Fan P, O'Reagan PA, Szerb JC. Effect of low glucose concentration on synaptic transmission in the rat hippocampal slice. *Brain Res Bull* 1988;21:741–747. [PubMed: 2851374]
- Frigge M, Hoaglin DC, Iglewicz B. Some implementations of the boxplot. *Amer Statistician* 1989;43:50–54.

- Fujii T, Baumgartl H, Lubbers DW. Limiting section thickness of guinea pig olfactory cortical slices studied from tissue pO₂ values and electrical activities. *Pflugers Arch* 1982;393:83–87. [PubMed: 7088686]
- Jefferys JG. Nonsynaptic modulation of neuronal activity in the brain: electric currents and extracellular ions. *Physiol Rev* 1995;75:689–723. [PubMed: 7480159]
- Jensen MS, Yaari Y. The relationship between interictal and ictal paroxysms in and in vitro model of focal hippocampal epilepsy. *Ann Neurol* 1988;24:591–598. [PubMed: 2849367]
- Kam K, Nicoll RA. Excitatory synaptic transmission persists independently of the glutamate-glutamine cycle. *J Neurosci* 2007;27:9192–9200. [PubMed: 17715355]
- Kirchner A, Veliskova J, Velisek L. Differential effects of low glucose concentrations on seizures and epileptiform activity in *in vivo* and *in vitro*. *Eur J Neurosci* 2006;23:1512–1522. [PubMed: 16553614]
- Korn SJ, Giacchino JL, Chamberlin NL, Dingledine R. Epileptiform burst activity induced by potassium in the hippocampus and its regulation by GABA-mediated inhibition. *J Neurophysiol* 1987;57:325–340. [PubMed: 3559679]
- LaManna JC, Chavez JC, Pichiule P. Structural and functional adaptation to hypoxia in the rat brain. *J Exp Biol* 2004;207:3163–3169. [PubMed: 15299038]
- Langmoen, IA.; Andersen, P. The Hippocampal Slice *In Vitro*. A description of the technique and some examples of the opportunities it offers. In: Kerkut, GA.; Wheal, HV., editors. *Electrophysiology of Isolated Mammalian CNS Preparations*. London: Academic Press; 1981.
- Leschinger A, Stabel J, Igelmund P, Heinemann U. Pharmacological and electrographic properties of epileptiform activity induced by elevated K⁺ and lowered Ca²⁺ and Mg²⁺ concentration in rat hippocampal slices. *Exp Brain Res* 1993;96:230–240. [PubMed: 7903641]
- Lipton, P.; Whittingham, TS. Energy metabolism and brain slice function. In: Dingledine, R., editor. *Brain Slices*. New York: Plenum Press; 1984.
- Masson J, Darmon M, Conjard A, Chuhma N, Ropert N, Thoby-Brisson M, Foutz AS, Parrot S, Miller GM, Jorisch R, Polan J, Hamon M, Hen R, Rayport S. Mice lacking brain/kidney phosphate-activated glutaminase have impaired glutamatergic synaptic transmission, altered breathing, disorganized goal-directed behavior and die shortly after birth. *J Neurosci* 2006;26:4660–4671. [PubMed: 16641247]
- McNay EC, Gold PE. Extracellular glucose concentrations in the rat hippocampus measured by zero-net-flux: microdialysis flow rate, strain, and age. *J Neurochem* 1999;72:785–790. [PubMed: 9930754]
- Mehta S. The glucose paradox of cerebral ischaemia. *J Postgrad Med* 2003;2003:299–301. [PubMed: 14699225]
- Merrill DR, Bikson M, Jefferys JGR. Electrical stimulation of excitable tissue: design of efficacious and safe protocols. *J Neurosci Methods* 2005;141:171–198. [PubMed: 15661300]
- Newman GC, Hospod FE, Schissel SL. Ischemic brain slice glucose utilization: effects of slice thickness, acidosis, and K⁺. *J Cereb Blood Flow Metab* 1991;11:398–406. [PubMed: 2016347]
- Nishimura F, Nishihara M, Mori M, Torii K, Takahashi M. Excitability of neurons in the ventromedial nucleus in rat hypothalamic slices: modulation by amino acids at cerebrospinal fluid levels. *Brain Res* 1995;691:217–222. [PubMed: 8590056]
- Novotny EJ Jr, Ariyan C, Mason GF, O'Reilly J, Haddad GG, Behar KL. Differential increase in cerebral cortical glucose oxidative metabolism during rat postnatal development is greater in vivo than in vitro. *Brain Res* 2001;888:193–202. [PubMed: 11150475]
- Parks JM, Ames A III, Nesbitt FB. Protein synthesis in central nervous tissue: studies on retina in vitro. *J neurochem* 1976;27:987–997. [PubMed: 12170628]
- Payne RS, Tseng MT, Schurr A. The glucose paradox of cerebral ischemia: evidence for corticosterone involvement. *Brain Res* 2003;971:9–17. [PubMed: 12691832]
- Patrylo PR, Schweitzer JS, Dudek FE. Potassium-dependent prolonged field bursts in the dentate gyrus: effects of extracellular calcium and amino acid receptor antagonists. *Neuroscience* 1994;61:13–19. [PubMed: 7969887]
- Pfriegeer FW, Barres BA. New views on synapse-glia interactions. *Curr Opin Neurobiol* 1996;6:615–621. [PubMed: 8937825]
- Radman T, Su Y, An JH, Parra LC, Bikson M. Spike timing amplifies the effect of electric fields on neurons: implications for endogenous field effects. *J Neurosci* 2007;27:3030–3036.

- Rosen AS, Andrew RD. Glucose concentration inversely alters neocortical slice excitability through an osmotic effect. *Brain Res* 1991;555:58–64. [PubMed: 1933330]
- Rothman DL, Sibson NR, Hyder F, Shen J, Behar KL, Shulman RG. In vivo nuclear magnetic resonance spectroscopy studies of the relationship between the glutamate-glutamine neurotransmitter cycle and functional neuroenergetics. *Philos Trans R Soc Lond B Biol Sci* 1999;354:1165–1177. [PubMed: 10466144]
- Rothman DL, Behar KL, Hyder F, Shulman RG. In vivo NMR studies of the glutamate neurotransmitter flux and neuroenergetics: implications for brain function. *Annu Rev Physiol* 2003;65:401–427. [PubMed: 12524459]
- Saly V, Andrew RD. CA3 neuron excitation and epileptiform discharge are sensitive to osmolality. *J Neurophysiol* 1993;69:2200–2208. [PubMed: 8350139]
- Schiff SJ, Szerb JC, Somjen GG. Glutamine can enhance synaptic transmission in hippocampal slices. *Brain Res* 1985;343:366–369. [PubMed: 2996699]
- Schiff SJ, Somjen GG. Hyperexcitability following moderate hypoxia in hippocampal tissue slices. *Brain Res* 1985;337:337–340. [PubMed: 2992681]
- Schurr A, Changaris DG, Rigor BM. Glutamine protects neuronal function against cerebral hypoxia: a study using the in vitro hippocampal slice preparation. *Brain Res* 1987;412:179–181. [PubMed: 3607453]
- Schurr A, West CA, Rigor BM. Electrophysiology of energy metabolism and neuronal function in the hippocampal slice preparation. *J Neurosci Methods* 1989;28:7–13. [PubMed: 2725015]
- Schurr A, Miller JJ, Payne RS, Rigor BM. An increase in lactate output by brain tissue serves to meet the energy needs of glutamate-activated neurons. *J Neurosci* 1999a;19:34–39. [PubMed: 9870935]
- Schurr A, Payne RS, Miller JJ, Rigor BM. Study of cerebral energy metabolism using the rat hippocampal slice preparation. *Methods* 1999b;18:117–126. [PubMed: 10356342]
- Schwartzkroin PA, Baraban SC, Hochman DW. Osmolarity, ionic flux, and changes in brain excitability. *Epilepsy Res* 1998;32:275–285. [PubMed: 9761327]
- Schwechter EM, Veliskova J, Velisek L. Correlation between extracellular glucose and seizure susceptibility in adult rats. *Ann Neurol* 2003;53:91–101. [PubMed: 12509852]
- Silver IA, Erecinska M. Extracellular glucose concentration in mammalian brain: continuous monitoring of changes during increased neuronal activity and upon limitation in oxygen supply in normo-, hypo-, and hyperglycemic animals. *J Neurosci* 1994;14:5068–5076. [PubMed: 8046468]
- Szerb JC, O'Regan PA. Effect of glutamine on glutamate release from hippocampal slices induced by high K^+ or by electrical stimulation: interaction with different Ca^{2+} concentrations. *J Neurochem* 1985;40:1724–1731. [PubMed: 2859354]
- Tani H, Bandrowski AE, Parada I, Wynn M, Huguenard JR, Prince DA, Reimer RJ. Modulation of epileptiform activity by glutamine and system A transport in a model of post-traumatic epilepsy. *Neurobiol Dis* 2007;25:230–238. [PubMed: 17070687]
- Traynelis SF, Dingledine R. Potassium-induced spontaneous electrographic seizures in the rat hippocampal slice. *J Neurophysiol* 1988;59:259–276. [PubMed: 3343603]
- Traynelis SF, Dingledine R. Role of extracellular space in hyperosmotic suppression of potassium-induced electrographic seizures. *J Neurophysiol* 1989;61:927–938. [PubMed: 2723735]
- Tsacopoulos M, Magistretti PJ. Metabolic coupling between glia and neurons. *J Neurosci* 1996;16:877–885. [PubMed: 8558256]
- Winkler BS, Kapousta-Bruneau N, Arnold MJ, Green DG. Effects of inhibiting glutamine synthetase and blocking glutamate uptake on b-wave generation in the isolated rat retina. *Vis Neurosci* 1999;16:345–353. [PubMed: 10367968]
- Xiang J, Ennis SR, Abdelkarim GE, Fujisawa Mutsuo, Kawai N, Keep RF. Glutamine transport at the blood-brain and blood-cerebrospinal fluid barrier. *Neurochem Int* 2003;43:279–288. [PubMed: 12742070]
- Yuan H, Yamada K, Inagaki N. Glucose sensitivity in mouse substantia nigra pars reticulata neurons in vitro. *Neurosci Lett* 2004;355:173–176. [PubMed: 14732459]
- Zheng L, Godfrey DA, Waller HJ, Godfrey TG, Chen K, Wong W. Metabolism of the dorsal cochlear nucleus in rat brain slices. *Hearing Res* 2000;143:115–129.

Fig. 1. Schematic of key

A) *in vivo* brain perfusion mechanisms and B) *in vitro* superfusion mechanisms relating to metabolite/reactant delivery/clearance. Mechanical slicing of tissue results in ‘fundamental’ traumas regardless of the details of the preparation method. During slice preparation and handling some mechanical damage occurs at the boundary of the slice. The vasculature of the slice is rendered inoperative. In the absence of circulation, blood vessels no longer move substances such as oxygen and glucose into the tissue and CO_2 out of the tissue. In addition, CSF flow and bulk flow of interstitial fluid no longer operate as a transport mechanism for substances into and through the brain. Surface ACSF flow is the only convective mechanism for transport remaining in the slice. In brain slices, the delivery and clearance kinetics of metabolites is limited by diffusion with the surface.

Fig. 2. Effect of 0.5 mM glutamine on orthodromic population spikes evoked in the CA1 region of hippocampal brain slices

A) Sample traces of control and glutamine- treated population spikes. B) Paired-pulses were delivered to the Schaeffer-collaterals every 2 minutes and supra-threshold responses monitored in the CA1 pyramidal layer. Extracellular glucose was maintained at 10 mM. There was no significant difference between control (○; n = 7) and 0.5 mM glutamine (◆; n = 18) treated slices. C) No significant difference between control and glutamine-treated ratios for either facilitation or depression cases.

Fig. 3. Effect of 2–5 mM glutamine on hippocampal brain slice neurophysiology. Baseline ACSF glucose was 5 mM. Asterisks (*) indicates initiation of spreading-depression

A) Continuous recording of CA1 field potential during application of 5 mM glutamine (bar). Paired-pulse orthodromic responses were monitored every 2 minutes. B) Average responses across slices; for each slice data is presented as time before of spreading-depression (at $t = 0$).

Fig. 4. Effect of 5 mM glucose on orthodromic population spikes evoked in the CA1 region of hippocampal brain slices

A) Sample traces of 5 mM glucose- treated population spikes. B) Extracellular glucose was reduced from 10 to 5 mM and paired-pulse responses monitored every 2 minutes. There was a significant difference between control ($\circ$; $n = 7$) and 5 mM glucose ($\blacklozenge$; $n = 6$) treated slices only in pulse 1. C) There was no significant difference in paired-pulse ratio for either facilitation or depression cases.

Fig. 5. Effect of 2 mM glucose in combination with glutamine on paired-pulse orthodromic population spikes evoked in the CA1 region of hippocampal brain slices

A) Sample traces of 2 mM glucose- treated population spikes. B) Change of glucose concentration from 10 to 2 mM. Population spike amplitude of both pulses is significantly reduced at t = 50 min (n = 6). C) Change of glucose concentration from 10 to 2 mM and the presence of 0.5 mM glutamine. Glutamine does not prevent the population spike attenuation (n = 5).

Fig. 6. Effect of 0.5 mM glutamine on high- K^+ induced epileptiform activity

A) No significant changes were observed between baseline (top trace) and glutamine application (bottom trace, $t = 52$ min). B) Glutamine does not affect burst duration, amplitude, or frequency compared to baseline ($n = 5$).