Key messages

- Major differences exist among European countries in the timing and degree of adoption of measures to control a bloodborne HIV epidemic
- Despite many weaknesses, official AIDS notification data help in understanding the size of the problem in Europe and the United States
- Annual age adjusted incidence rates of bloodborne AIDS for 1985-93 were over 3 per million in Romania, the United States, and France
- Rates were lower than 1 per million in Italy, Germany, Sweden, and the Netherlands
- Declines in incidence of bloodborne AIDS in Europe and the United States started in the late 1980s (children) and early 1990s (adults)

Data from the European non-aggregate AIDS dataset (ENAADS), version AIDS9403.DAT, prepared by the European Centre for the Epidemiological Monitoring of AIDS in Paris, was used in this study. Compilation of this data file was made possible by the continuing participation of clinical doctors in mandatory and voluntary national reporting schemes for AIDS cases. We wish to thank the National Center for Infectious Diseases, Centers for Disease Control and Prevention (CDC), Atlanta, for AIDS public information dataset updated to June 1993; Dr Eva Negri for useful comments; Mrs Luigina Mei for editorial assistance; and Dr Angelo Lo Re and Mr Roberto Ricci for technical assistance.

Funding: This work was supported by two grants from the Ministero della Sanità-Istituto Superiore di Sanità, VIII Progetto AIDS contracts 9303-12 and 9303-31; it was conducted within the framework of a European Union project on sexually transmitted diseases and AIDS.

Conflict of interest: None.

1 Centers for Disease Control. Possible transfusion-associated acquired immune deficiency syndrome (AIDS) —California. *MMWR Morb Mortal Wkly Rep* 1982;31:652-4.

- 2 Ness PM, Nass CC. Blood donor testing for HIV-I/II and HTLV-I/II. *Arch Pathol Lab Med* 1994;118:337-41.
- 3 Anderson C. AIDS scandal indicts French government. *Nature* 1991;353:197.
- 4 Bader J-M. France: doctors face charges over HIV-infected blood. *Lancet* 1992;340:360.
- 5 Hersh BS, Popovici F, Apetrei RC, Zolotuska L, Beldescu N, Calomfirescu A, et al. Acquired immunodeficiency syndrome in Romania. *Lancet* 1991;338:645-9.
- 6 Patrascu IV, Dumitrescu O. The epidemic of human immunodeficiency virus infection in Romanian children. *AIDS Res Hum Retroviruses* 1993;9:99-104.
- 7 Dal Maso L, Franceschi S, Negri E, Serraino D, La Vecchia C, Ancelle-Park RA. Trends of AIDS incidence in Europe and the United States. *Soz Präventivmed* 1995;40:239-65.
- 8 Selik RM, Ward JW, Buehler JW. Trends in transfusion-associated acquired immune deficiency syndrome in the United States, 1982 through 1991. *Transfusion* 1993;33:890-3.
- 9 Selik RM, Ward JW, Buehler JW. Demographic differences in cumulative incidence rates of transfusion-associated acquired immunodeficiency syndrome. *Am J Epidemiol* 1994;140:105-12.
- 10 Franceschi S, Dal Maso L, La Vecchia C, Negri E, Serraino D. AIDS incidence rates in Europe and the United States. *AIDS* 1994;8:1173-7.
- 11 Centers for Disease Control. Revision of the case definition for acquired immunodeficiency syndrome. *MMWR Morb Mortal Wkly Rep* 1985;34:373-5.
- 12 Centers for Disease Control. Revision of the CDC surveillance case definition for acquired immunodeficiency syndrome. *MMWR Morb Mortal Wkly Rep* 1987;36:3-15S.
- 13 World Health Organisation. Acquired immunodeficiency syndrome (AIDS)—WHO/CDC case definition for AIDS. *Wkly Epidemiol Rec* 1986;61:69-76.
- 14 World Health Organisation. Acquired immunodeficiency syndrome (AIDS)—1987 revision of CDC/WHO case definition for AIDS. *Wkly Epidemiol Rec* 1988;63:1-8.
- 15 European Centre for the Epidemiological Monitoring of AIDS. *AIDS surveillance in Europe*. No 37. Paris: Hospital Saint-Maurice, 1993.
- 16 Doll R, Smith PG. Comparison between registries: age-standardized rates. In: Waterhouse J, Muir C, Shanmugaratnam K, Powell J, eds. *Cancer incidence in five continents*. Vol IV. Lyons: International Association for Research on Cancer, 1982:58-88. (Scientific publication No 42.)
- 17 Rosenberg PS. A simple correction of AIDS surveillance data for reporting delays. *J Acquir Immune Defic Syndr* 1990;3:49-54.
- 18 Goedert JJ, Kessler CM, Aledort LM, Biggar RJ, Andes WA, White GC II, et al. A prospective study of human immunodeficiency virus type 1 infection and the development of AIDS in subjects with hemophilia. *N Engl J Med* 1989;321:1141-8.
- 19 Ward JW, Bush TJ, Perkins HA, Lieb LE, Allen JR, Goldfinger D, et al. The natural history of transfusion-associated infection with human immunodeficiency virus. Factors influencing the rate of progression to disease. *N Engl J Med* 1989;321:947-52.
- 20 Medley GF, Anderson RM, Cox DR, Billard L. Incubation period of AIDS in patients infected via blood transfusion. *Nature* 1987;328:719-21.
- 21 Blaxhult A, Granath F, Lidman K, Giesecke J. The influence of age on the latency period to AIDS in people infected by HIV through blood transfusion. *AIDS* 1990;4:125-9.

(Accepted 11 October 1995)

Prognostic trees to aid prognosis in patients with cutaneous malignant melanoma

Tom C Aitchison, Jane M Sirel, Douglas C Watt, Rona M MacKie, for the Scottish Melanoma Group

Abstract

Objectives—To design user friendly guides to prognosis for patients who have had invasive primary cutaneous malignant melanomas surgically excised.

Design—Adaptation of the classification tree method was used to derive prognostic trees for four different subgroups of malignant melanoma patients in whom known and possible prognostic variables interacted in different ways.

Setting—Scotland.

Subjects—Statistical modelling for prognostic trees was based on 1978 patients whose primary malignant melanoma was first diagnosed in 1979-86 for whom five year follow up and all relevant clinical pathological data were available. The resultant model was validated with 300 patients first diagnosed in 1987 for whom the same information was available.

Main outcome measures—Actual and predicted rate of survival after diagnosis of primary cutaneous malignant melanoma.

Results—The four subgroups of patients were men and women with ulcerated and non-ulcerated cutaneous primary melanomas. Validation of the model showed excellent agreement between actual

status of patients in the relevant subgroups and their status as predicted by the model.

Conclusions—The prognostic trees are simple to use and give more accurate prognosis for individual patients than is currently available from tumour thickness alone.

Introduction

Studies have shown that several factors affect the prognosis and survival of patients with primary invasive cutaneous malignant melanoma.¹⁻⁴ Most reports agree that tumour thickness is the most important prognostic factor, but several other factors seem to interact, including sex, presence or absence of ulceration, level of invasion, age at diagnosis, site of tumour, and type of tumour.

We previously reported the results of proportional hazards models developed by the Scottish Melanoma Group to predict survival of patients with primary cutaneous malignant melanoma.⁵ Four distinct subgroups of patients were identified from this analysis—men and women with ulcerated or non-ulcerated lesions. Within each of these subgroups different sets of prognostic factors were identified, although tumour thickness was always a key factor. For example, among

Department of Statistics,
Glasgow University,
Glasgow G12 8QW
Tom C Aitchison, senior
lecturer
Jane M Sirel, research
assistant
Douglas C Watt, lecturer

Department of
Dermatology, Glasgow
University, Glasgow
G12 8QQ
Rona M MacKie, professor

Correspondence to:
Professor MacKie.

BMJ 1995;311:1536-9

women with ulcerated lesions the site of the primary lesion and the mitotic count were also of prognostic value, whereas tumour thickness was the only relevant prognostic factor for women with non-ulcerated lesions. The details of this analysis are available from us on request in the form of a technical report and would be the preferred basis for predicting prognosis for an individual patient; however, predicting individual patients' survival profiles, and hence likely prognoses, by means of such a proportional hazards model is reasonably complicated, and, as the calculations are computer based, the model is not really suitable for general and widespread use.

We therefore adapted the proportional hazards models to create simplified, user friendly guides to prognosis that could be used for counselling individual patients as well as for identifying different risk groups for inclusion and stratification in clinical trials of any new treatment.

Methods

PROPORTIONAL HAZARDS MODEL

The Scottish Melanoma Group carried out a full modelling exercise with proportional hazards models that included the variables sex, level of invasion, age at

Fig 3—Prognostic tree for survival of patients with primary cutaneous malignant melanoma: men with ulcerated lesions

diagnosis, site of tumour, type of tumour, numbers of mitoses (low or high), pre-existing naevus, regression, and presence or absence of obvious ulceration on histological examination. The analysis was based on all 1978 patients registered with the Scottish Melanoma Group who had a stage 1 primary cutaneous malignant melanoma of Clark level ≥ 2 during 1979–86, for whom there was full information on all the possible prognostic factors, and who had a minimum of four years follow up. The endpoint in this and all subsequent analyses was death due to malignant melanoma.

SIMPLE PROGNOSTIC TREES

We reanalysed the results for the 1978 patients by means of an adaptation (see below) of classification trees' for each of the four subgroups identified in the full analysis (that is, based on the proportional hazards models). For each subgroup, the classification tree analysis provided a tree structure, with two branches at each step of the tree and new branches being formed until the data had no further important prognostic information (that is, the data could not be further subdivided by any of the possible prognostic factors to be of any benefit in predicting survival).

These versions of classification trees were based on log rank tests' at each step in the tree and were therefore appropriate for analysing censored data by a tree based approach. An empirical rule based on a cross validation approach' was used to determine the pruned trees (figures 1–4), with the extra condition that, whenever possible, at least 50 patients were available at each of the final nodes of the trees. This was achieved in 14 of the 16 nodes among the four trees.

After each node had been arrived at by means of the above approach a Kaplan-Meier estimate of the survival function for the particular subgroup was calculated and used to provide the estimates of two year and five year survival.

ESTIMATING PERFORMANCE OF THE PROGNOSTIC TREES

As a first step in assessing the usefulness and accuracy of these trees, they were applied to all 300 patients registered with the Scottish Melanoma Group whose disease was diagnosed in 1987 and who had a minimum of four years of follow up. We applied the appropriate Kaplan-Meier estimate based on the 1979–86 database to each of the 300 test patients and evaluated, for each, the probability of surviving at least one, two, three, and four years. These individual estimates were then summed over all patients in each of the subgroups and compared with the actual number of patients who survived past each of one, two, three, and four years.

Results

THE PROGNOSTIC TREES

Figures 1–4 show the results of the classification tree analyses for the four subgroups. At the end of each final branch is a node (the boxes at the foot of each tree diagram). These nodes contain the number of patients in that particular subdivision of the data and, for

Fig 1—Prognostic tree for survival of patients with primary cutaneous malignant melanoma: women with ulcerated lesions

Fig 2—Prognostic tree for survival of patients with primary cutaneous malignant melanoma: women with non-ulcerated lesions

illustration, the predicted two year and five year survival for these patients.

For example, if a woman presented with a 4.5 mm thick ulcerated lesion, figure 1 indicates that she would have a 66% chance of surviving at least two years and a 43% chance of surviving at least five years (based on a sample of 151 women with ulcerated tumours thicker than 3.9 mm). However, if another woman presented with an ulcerated lesion of only 2 mm then one would take the left hand branch of the tree in figure 1 and ask her age; if she was, say, 50 years old one would carry on through the next step to estimate her prospects of surviving two years as 92% and of surviving five years as 77%.

PERFORMANCE OF THE PROGNOSTIC TREES

Table 1 shows that the general performance of the prognostic trees was very good when the observed survival of the 300 patients diagnosed in 1987 was compared with their predicted survival based on the trees. For example, of the 151 women with non-ulcerated lesions diagnosed in 1987, 129 actually survived at least four years, while the predicted number based on the tree in figure 2 was 127. In general the predicted numbers of surviving patients

Table 1—Survival of 300 patients with primary cutaneous malignant melanoma diagnosed in 1987: comparison of observed results and survival predicted from prognostic trees. Values are numbers of patients

	No of years after diagnosis			
	1	2	3	4
Women with ulcerated lesions:				
Patients at risk*	46	44	44	33
Survival (observed†:predicted)	42:42-4	36:33-9	33:30-3	21:19-8
Women with non-ulcerated lesions:				
Patients at risk*	151	149	143	132
Survival (observed†:predicted)	150:149-9	148:146-5	141:139-5	129:126-8
Men with ulcerated lesions:				
Patients at risk*	32	30	30	28
Survival (observed†:predicted)	28:29-1	20:22-1	18:18-5	15:13-7
Men with non-ulcerated lesions:				
Patients at risk*	71	68	66	61
Survival (observed†:predicted)	71:69-2	65:63-3	61:59-8	54:51-8

*Number of patients at risk from diagnosis up to and including the specified number of years.

†Number of patients whose status (that is, whether dead due to melanoma) was certain at the appropriate time after prognosis.

Table 2—Estimated disease free survival of patients at five years after diagnosis of primary cutaneous malignant melanoma: comparison of estimates from proportional hazards model and prognostic trees

Case information		Estimated five year survival (%)	
Tumour thickness (mm)	Other relevant risk factors	Proportional hazards model	Prognostic trees
Women with ulcerated lesions:			
4.5	Mitosis medium to high*; site axial	35	43
3.0	Mitosis medium to high*; site extremity; age 70	63	67
2.0	Mitosis medium to high*; site extremity; age 70	65	67
Women with non-ulcerated lesions:			
2.5	Age 50	92	89
1.5	None	94	93
0.5	Level of invasion 3; site axial	96	92
Men with ulcerated lesions:			
5.5	None	45	34
4.5	None	46	52
3.5	None	48	52
Men with non-ulcerated lesions:			
3.5	Level of invasion 4, 5	69	52
1.5	Level of invasion 4, 5	79	76
0.5	Level of invasion 2, 3	94	99

*Mitosis levels were low or medium to high. While they were significant in the proportional hazards model, they were not in the prognostic trees model.

Fig 4—Prognostic tree for survival of patients with primary cutaneous malignant melanoma: men with non-ulcerated lesions

were close to the corresponding observed numbers; only at the fourth year after diagnosis was there a slight suggestion of conservative underestimation of survival in all four subgroups of patients.

As a further indication of the prognostic accuracy of the trees, table 2 gives the estimated five year rate of disease free survival for potential patients with malignant melanoma and a wide range of prognostic factors that were calculated with the full survival analysis, based on the proportional hazards model, and the prognostic trees in figures 1-4. For example, a man with an ulcerated lesion 3.5 mm thick was predicted to have a 48% probability of surviving at least five years based on the proportional hazards model and a 52% probability based on the prognostic trees—a good agreement. Table 2 shows that, in general, the performance of the trees was close to the "correct" result based on the more technically correct proportional hazards model.

Conclusions

These prognostic trees are simple guides to the likelihood of survival from malignant melanoma. They performed well with the test sample of 300 patients both in predicting the number of survivors at up to four years and in their predictive value relative to the full proportional hazards model. They are currently used in our clinic as an aid to counselling patients after diagnosis of melanoma and enable advice to be directed more specifically to individual patients. With greater awareness of melanoma in the community, and thus in some cases earlier presentation, these trees will probably need to be revised about every five years, and we plan to do this. In collaboration with colleagues outside Glasgow we are also testing the robustness of the trees in other geographic areas.

The one disadvantage, in a sense, with the trees is the need to split each prognostic factor into two possible groups at each stage of a tree's construction. This is fine for variables such as site that could take only two values (axial or extremity), but produces slightly arbitrary categorisations of continuous variables such as tumour thickness, age, etc. This means that in figure 1, for example, a woman with an ulcerated lesion 4 mm thick (that is, just above the cut off point) gets a substantially different prognosis from a similar patient with a lesion 3.9 mm thick. Clearly this is rather arbitrary for patients near the cut off point, and care should be taken with such patients, with perhaps the more optimistic route being explored first. However, this is somewhat balanced for by the

Key messages

- Prognosis for patients with primary invasive cutaneous malignant melanoma can be assessed with considerable accuracy at the time of excision of the primary tumour
- We developed four prognostic trees to predict prognosis in four subgroups of patients—men and women with ulcerated or non-ulcerated primary tumours—in whom the possible prognostic variables interacted differently
- Prospects for five year disease free survival ranged from 100% for women with non-ulcerated lesions to only 34% for men with ulcerated lesions
- Validation of the trees by comparison with results for 300 patients showed excellent agreement between actual status of patients in the relevant subgroups and their predicted status
- These prognostic trees are a simple and accurate method of predicting prognosis of patients with malignant melanoma

fact that the trees use such variables more than once and more variables in total than the proportional hazards model to compensate for the rough categorisation of continuous variables such as tumour thickness. For example, in the prognostic prediction for women with non-ulcerated lesions (fig 2) tumour thickness is used twice, while age, level of invasion, and site are also included.

In spite of these differences with the full and

somewhat more complicated approach of the survival analysis, the trees seem to be effective prognostic indicators for the vast majority of patients. These tree diagrams can be used in a clinic as a quick and simple means of assessing an individual patient's prospects of survival after diagnosis of a malignant melanoma.

We thank the secretaries of the Scottish Melanoma Group, Miss E Salt and Mrs J Stewart, for their help.

Funding: We thank the Scottish Hospital Endowment Research Trust for financial support and the Cancer Research Campaign for funding the Scottish Melanoma Group.

Conflict of interest: None.

- 1 Clark WH, Elder DE, Guerry D, Braitman LE, Trock BJ, Schultz D, *et al*. A model predicting survival in stage I melanoma based on tumour progression. *J Natl Cancer Inst* 1989;81:1893-904.
- 2 Balch CM, Soong SJ, Shaw HM, Urist MM, McCarthy WH. An analysis of prognostic factors in 8500 patients with cutaneous melanoma. In: Balch CM, Houghton AN, Milton GW, Sober AJ, Soong SJ, eds. *Cutaneous melanoma*. 2nd ed. Philadelphia: Lipincott, 1992:165-87.
- 3 Vossaert KA, Silverman MK, Kopf AW, Bart RS, Rigel DS, Friedman MD, *et al*. Influence of gender on survival in patients with stage I malignant melanoma. *J Am Acad Dermatol* 1992;26:429-40.
- 4 Rigel DS, Friedman RJ, Kopf AW, Silverman MK. Factors influencing survival in melanoma. *Melanoma/Skin Cancer Update* 1991;9:631-42.
- 5 MacKie RM, Aitchison TC, Sirel JM, Watt DC. Prognostic models for subgroups of melanoma patients from the Scottish Melanoma Group Database 1979-1986, and their subsequent validation. *Br J Cancer* 1995;71:173-6.
- 6 Brieman L, Friedman JH, Olshen RA, Stone CJ. *Classification and regression trees*. Belmont, CA: Wadsworth International, 1984.
- 7 Everitt BS. The analysis of survival data. In: *Statistical methods for medical investigation*. Oxford: Oxford University Press, 1989:83-98.

(Accepted 11 October 1995)

Commentary: Prognostic models: clinically useful or quickly forgotten?

Jeremy C Wyatt, Douglas G Altman

We are all familiar with using single items of patient data such as age or smoking history to help in making difficult clinical decisions.¹ Prognostic models are more complex tools for helping decision making that combine two or more items of patient data to predict clinical outcomes.² They are of potential value when doctors are making difficult clinical decisions (such as ordering invasive tests or selecting which patients should benefit from scarce resources),³ conducting comparative audit,⁴ or selecting uniform groups of patients for clinical trials.⁵ Another prognostic model appears in this week's *BMJ*⁶ to join the hundreds published every year.⁷ However, apart from exceptions such as the Glasgow coma scale⁸ and APACHE III,⁹ few of these models are routinely used to inform difficult clinical decisions.

It might be argued that doctors never prognosticate, working always in the present, but studies of medical decision making show this is untrue.^{10,11} Some doctors might claim that they can foretell patient outcomes better than any statistical model, but again there is contrary evidence.^{9,12} A journal editor's view might be that most published models reflect preliminary work and need further research before clinical adoption.¹³ Finally, some models predict events that are of no clinical relevance or do not generate predictions in time to inform clinical decisions, suggesting that their developers wished merely to publish journal articles, not build clinically useful tools.¹⁴

We believe that the main reasons why doctors reject published prognostic models are lack of clinical credibility and lack of evidence that a prognostic model can support decisions about patient care (that is, evidence of accuracy, generality, and effectiveness). We examine each of these issues in turn.

Clinical credibility of the model

However accurate a model is in statistical terms, doctors will be reluctant to use it to inform their patient management decisions unless they believe in the model and its predictions. Some prerequisites for clinical credibility include:

- All clinically relevant patient data should have been tested for inclusion in the model. For example, it would be foolish to omit smoking history from the variables screened when constructing a model to predict the probability of acute myocardial infarction
- It should be simple for doctors to obtain all the patient data required, reliably and without expending undue resources, in time to generate the prediction and guide decisions. Data should be obtainable with high reliability, particularly in those patients for which the model's prediction are most likely to be needed. This is not always the case¹⁵
- Model builders should try to avoid arbitrary thresholds for continuous variables. For example, it is unlikely that the prognosis for a woman with an ulcerated melanoma 3.9 mm thick would be very different if it was 4 mm thick. (Aitchison *et al*⁶ discuss this issue in their paper.⁶)
- The model's structure should be apparent and its predictions should make sense to the doctors who will rely on them, as only then will the law treat users of the model as "learned intermediaries" in a case of alleged negligence.¹⁶ This requirement means that the statistical modelling method must be correctly applied and not transgress the method's assumptions; these were checked in only a fifth of one series of published models.¹⁷ It also makes "black box" models such as

Medical Informatics,
Biomedical Informatics
Unit, Imperial Cancer
Research Fund, PO Box
123, London WC2A 3PX
Jeremy C Wyatt, consultant

ICRF Medical Statistics
Group, Centre for
Statistics in Medicine,
Institute for Health
Sciences, Oxford OX3 7LF
Douglas G Altman, head