help numb the pain afterwards. When you wake up you will be breathing oxygen. All this is absolutely routine. Is that all right?" This might not satisfy all my patients who would question me about details. These would be answered but matters would be amplified only as required by the patient.

The professional conduct committee appears to have ruled that my approach would have been too paternalistic.

Outcome

Mitchell has indicated what has happened in the respondent's hospital and that the decision has been applied to day units there. Some of the reactions may be excessive, but there is a need for guidance to indicate the limits to which every anaesthetist shall go in seeking consent. The professional conduct

committee's decision has considerable resource implications for the NHS.

I assume that it was the generic "suppository" that was relevant, not the specific drug content. The absence of knowledge about the intention to use it resulted in legal assault. Nevertheless, the action was benign in intent. The penalty of being found guilty of serious professional misconduct beyond reasonable doubt (the standard of proof for a criminal) is surely out of proportion. We shall not know the reasons for this verdict, as the GMC is shy of open debate. Perhaps there are no grades of professional misconduct. In the absence of a penalty other than admonition there can be no appeal. Meanwhile, it seems to me that this entire case was provocative at first, meddlesome in the middle, and outrageous at the end.

1 Mitchell J. A fundamental problem of consent. *BMJ* 1995;310:43-6.

Lesson of the Week

Hypocalcaemic stridor and infantile nutritional rickets

J J A Train, R W Yates, M R J Sury

Infants with vitamin D deficiency and rickets may present with stridor and laryngospasm due to severe hypocalcaemia

Nutritional deficiency of vitamin D is considered to be a rare cause of rickets in industrialised countries, mainly because vitamin D is added routinely to formula feeds for infants.^{1,2} Human breast milk, however, may be deficient in vitamin D, and exclusively breast fed infants who have not received vitamin supplements are at risk. Otherwise well nourished infants may not have the obvious or typical bony deformities of rickets but may present with the symptoms of hypocalcaemia. We describe two infants who presented with stridor and intermittent sudden airway obstruction due to laryngospasm. Both infants had severe hypocalcaemia due to nutritional deficiency of vitamin D.

Case reports

CASE 1

A 5 month old black boy weighing 7.5 kg was taken to the accident and emergency department at Queen Elizabeth Hospital for Children with a one month history of intermittent laboured and noisy breathing. The boy's respiratory distress had often been accompanied by generalised rigidity, convulsions, and cyanosis and had usually occurred when he was crying. He was taken to the hospital because he had become apnoeic at home and had been given mouth to mouth resuscitation by his mother. Just after he was admitted he became distressed and stridulous. Convulsions followed, which were treated with rectal paraldehyde. He had enlarged tonsils, and a nasopharyngeal airway seemed to relieve the airway obstruction partially. Initially the convulsions were thought to be caused by hypoxaemia, and a tonsillectomy was proposed. The plasma calcium concentration, however, was 1.15 mmol/l, and vitamin D deficiency and rickets were diagnosed from the results of biochemical tests and an x ray examination. The boy was successfully treated with intravenous calcium gluconate followed by oral calcium and vitamin D supplements.

CASE 2

A 7 month old black boy weighing 7.5 kg was brought to the same hospital one year later. His general

practitioner had referred him for investigation of upper airway noises and recurrent respiratory infections, which he had had for three months. According to his parent, a sibling had had his "narrowed windpipe" surgically corrected.

The boy was tachypnoeic and had both inspiratory and expiratory stridor. Coarse chest crepitations were heard on auscultation, and a chest x ray film showed atelectasis of the left upper lobe. The stridor became worse. Examination of the oropharynx caused crying followed by complete upper airway obstruction and generalised convulsions, which were successfully treated with continuous positive airway pressure and rectal diazepam. Examination of the larynx under anaesthesia was proposed. Tetanic spasms of the arms were observed, however, and hypocalcaemia was suspected. The plasma calcium concentration was 0.9 mmol/l and the QTc interval (rate corrected) of 0.52 s (normal range 0.35-0.44 s) in an electrocardiogram was prolonged. An intravenous infusion of calcium gluconate was started and vitamin D supplements were given. Biochemical investigations and x ray examinations confirmed vitamin D deficiency and rickets.

RESULTS OF BIOCHEMICAL TESTS

The table shows the results of the biochemical tests in both cases, which confirmed the diagnoses of vitamin D deficiency. Both boys had been exclusively breast fed. Typical features of rickets were seen in the x ray films for both infants, including splaying of long bone metaphyses, expansion of the joints, and widening

Results of biochemical tests done in two black infants aged 5 and 7 months who were admitted to hospital with stridor and intermittent sudden airway obstruction. Reference ranges are given in parentheses

Biochemical test	Case 1	Case 2
Calcium (mmol/l) (2.41-2.77)	1.15	0.9
Albumin (g/l) (35-55)	42	41
Alkaline phosphatase (U/l) (55-265)	471	1406
Phosphate (mmol/l) (1.2-2.1)	1.78	1.57
Magnesium (mmol/l) (0.74-1.0)	0.72	0.54
Vitamin D metabolite:		
25-Hydroxycholecalciferol (nmol/l) (7.5-75)	<2.5	
Cholecalciferol (nmol/l) (15-100)		6
Parathyroid hormone (ng/l) (5-45)	118	195

Department of Anaesthesia, Hospitals for Sick Children, Great Ormond Street, London WC1 3JN

J J A Train, senior registrar
M R J Sury, consultant

Department of Paediatrics, Queen Elizabeth Hospital for Children, London E2 8PS

R W Yates, senior house officer

Correspondence to: Dr Sury.

BMJ 1995;310:48-9

ing of the anterior ends of ribs. Acute respiratory symptoms resolved within hours of treatment being started. Plasma calcium concentrations gradually increased and were in the normal range by 10 days in case 1 and 13 days in case 2.

Discussion

Vitamin D deficiency is a rare cause of hypocalcaemia and tetany,² and, although the causes of paediatric stridor are well known, with 80% of cases being due to infection,³ hypocalcaemia may be overlooked.⁴ Fortunately, in both of these cases severe hypocalcaemia was diagnosed in time for surgical intervention to be prevented.

Both infants were at risk of developing rickets: they had been exclusively breast fed, were black, and lived in an inner city area.^{1,5} Typically, term infants with an inadequate nutritional intake of vitamin D may present at between 3 and 6 months of age and are at risk, particularly during the winter.^{2,5} Vitamin D supplements are recommended for lactating mothers and for children up to the age of 2 years.⁵⁻⁷

In infants and small children the features of vitamin D deficiency include lethargy, recurrent respiratory infections, stridor, tetany, convulsions, and rickets.^{1,8} Infants who are too young to walk will not develop the limb deformities of rickets, such as genu varum. Hypocalcaemia reduces myocardial contractility and may result in arrhythmias and left ventricular failure. Cardiac tetany, although rare, is often fatal. An electrocardiogram with a prolonged QTc interval may indicate hypocalcaemia before biochemical investigations are done.

Hyperventilation reduces the arterial carbon dioxide partial pressure and causes the plasma pH to increase. The plasma concentration of ionised calcium then

decreases and may cause carpopedal spasm and tetany. Laryngospasm related to exercise was described in a child with hypocalcaemia, and a similar mechanism was suggested.⁹ Crying, and consequent hyperventilation, may worsen existing stridor of any aetiology but may also cause tetany and laryngospasm when hypocalcaemia is present.

If infants are exclusively breast fed and their mothers are not given dietary supplements of vitamin D then the infants may develop rickets, become severely hypocalcaemic, and present with acute respiratory symptoms. The plasma calcium concentration should be measured urgently. Infants with severe hypocalcaemia may suddenly develop laryngospasm and should be treated urgently in a department where, if necessary, they can receive prompt tracheal intubation.

We thank Mr D Albert and Professor C B S Wood for their permission to report these cases.

- 1 Barness LA. Nutritional disorders. In: Nelson HE, Behrman MD, Vaughan VC, eds. *Textbook of paediatrics*. 12th ed. Philadelphia: W B Saunders, 1983:179-84.
- 2 Harrison HE. Tetany. In: Nelson HE, Behrman MD, Vaughan VC, eds. *Textbook of paediatrics*. 12th ed. Philadelphia: W B Saunders, 1983:249-53.
- 3 Maze A, Bloche E. Stridor in pediatric patients. *Anesthesiology* 1979;50:132-45.
- 4 Sharief N, Matthew DJ, Dillon MJ. Hypocalcemic stridor in children: how often is it missed? *Clinical Pediatrics* 1991;30:51-2.
- 5 Department of Health and Social Security. *Present day practice in infant feeding: third report*. London: HMSO, 1988. (Reports on health and social subjects, 32.)
- 6 Polnay L, Hull D. *Community paediatrics*. 1st ed. Edinburgh: Churchill Livingstone, 1985:110-7.
- 7 Department of Health Committee on Medical Aspects of Food Policy. *Dietary reference values for food, energy and nutrients for the United Kingdom*. London: HMSO, 1991:123-7. (Reports on health and social subjects, 41.)
- 8 Rizen MF. Anesthetic implications of concurrent diseases. In: Miller RD, ed. *Anesthesia*. 3rd ed. New York: Churchill Livingstone, 1990:816-7.
- 9 Hidalgo HA, Davis SH. Intermittent stridor and hypocalcaemia in childhood. *Pediatric Pulmonology* 1989;7:110-1.

(Accepted 7 April 1994)

A PATIENT WHO CHANGED MY PRACTICE

One child's death saved many others

Many years ago a baby of about 18 months was admitted to hospital under my care having taken 10 of his mother's imipramine tablets prescribed for her depression at becoming pregnant against her wishes. This was before the days of child resistant containers. Imipramine was then a relatively new drug, and the baby had convulsions and cardiac irregularities, which are recognised side effects. I got the fits under control, but the cardiac irregularities were beyond me. My paediatric cardiology colleague and later an adult cardiologist came to help. But it was to no avail and the baby died after about 10 hours. He was the only child under my care who died from medicinal poisoning.

All the time his mother was in and out of his cubicle. "Nobody told me it could do him any harm" she kept saying. I said to myself, "What on earth are we doing, allowing this to happen?" So I did a study of accidental poisoning in childhood, which was published in the *BMJ*. This led to my being put on the Central Health Services Council Committee on the hospital management of acute poisoning. Although the Medicines Commission refused to include the prevention of poisoning in the committee's remit, we managed to include a recommendation in the report that the Central Health Services Council should examine the prevention of poisoning in childhood.

Further pressure for action resulted in the establishment of a British Standards Institution (BSI) committee on the testing of child resistant containers, which included the use of young children and adults as testers along the lines of work in America. But the Medicines Commission refused to use this standard. It said that if a child was used as a tester for a possible child resistant container and was

poisoned subsequently the commission might be sued.

This seemed to put a stop to any further development. Many months later, however, there was a meeting on non-accidental injury at the Department of Health and Social Services chaired by Dr David Owen (now Lord Owen) when he was Under Secretary of State. At the end of the meeting Dr Owen asked me if I was the person interested in poisoning—how he knew remains a mystery—and asked me for my views on the Medicine Commission's comment on the BSI standard. I said I thought it was disastrous because without the use of a proper standard of test the label of being a child resistant container might be applied to any container, effective or otherwise. Dr Owen's emphatic reply was "So do I" and within a short space of time aspirin and paracetamol sold over the counter for children had, by regulation, to be sold in child resistant containers which had passed the BSI test. A year later this was extended to include aspirin and paracetamol sold over the counter for adults. Before the introduction of the regulations about 7000 children a year were admitted to hospital but within three years the number had come down to below 2000, while admissions from other drugs remained unchanged.

This effective piece of accident prevention made me look at the prevention of other types of accidents and thus to the formation of the Joint Committee on Childhood Accident Prevention under the aegis of the Medical Commission on Accident Prevention. This joint committee subsequently became the Child Accident Prevention Trust.

One death from poisoning is one too many, but it certainly altered my life.—HUGH JACKSON is a retired consultant paediatrician in Newcastle upon Tyne