those attempting such projects could read about the experience of others.

SASHA SHEPPERD
Research officer
CRISPIN JENKINSON
Deputy director

Health Services Research Unit,
Department of Public Health and Primary Care,
University of Oxford, Oxford OX2 6HE

PATRICK MORGAN
Consultant in public health medicine
Northamptonshire Health Authority,
Northamptonshire NN1 5DN

- 1 Black N. Experimental and observational methods of evaluation. *BMJ* 1994;309:540. (20-27 August.)
- 2 McWhinney IR, Bass MJ, Donner A. Evaluation of palliative care service: problems and pitfalls. *BMJ* 1994;309:1340-2. (19 November.)
- 3 Sheldon TA. Please bypass the PORT. *BMJ* 1994;309:142-3. (16 July.)

Identifying relevant studies for systematic reviews

EDITOR,—We agree with Kay Dickersin and colleagues' recommendations regarding the need for improved reporting of randomised controlled trials by authors and improved indexing of such trials in electronic databases.¹ We wish to make two additional points on the basis of our experience of searching for randomised controlled trials related to stroke.

Firstly, there is a need for improved indexing in Medline (and other databases) of the medical subjects as well as the terms used to identify trials. For example, the MeSH term that covers stroke is CEREBROVASCULAR-DISORDERS but this is imprecise as it also covers many conditions not related to stroke (such as vascular dementia and migraine). In addition, the term CEREBROVASCULAR-DISORDERS is not used consistently for all stroke trials, especially for trials relating to prevention and rehabilitation. After studying the text and MeSH headings of several hundred stroke trials, which we had identified using a variety of methods, we have had to add 13 further MeSH or free text terms to maximise the sensitivity of our search. This further reduced the precision: the search with maximal sensitivity (87% of the articles in Medline that related to stroke trials) had a precision of only 10%. When this search was applied to all journals in Medline over six years about 10 000 articles were retrieved, each of which had to be assessed.

Secondly, given the problems with electronic searching of Medline and the practical difficulties of organising hand searching of all journals likely to include relevant trials (we think that at least 300 journals have included stroke trials), we suggest that several overlapping search strategies should be used to ensure that as many as possible of the available randomised controlled trials are included in systematic reviews. The Cochrane Stroke Review Group uses several such strategies: hand searching 15 major journals and the proceedings of major meetings on stroke; electronic searching of Medline, Embase, the Index to Scientific and Technical Proceedings (a database of conference proceedings available through the Bath Information and Data Services), and two dissertation databases; searching of the Ottawa stroke trials registry; reviewing the bibliographies of trials and other relevant articles; and contacting drug companies and colleagues. Each of these methods has retrieved trials that would have been missed if a single search strategy had been used.

CARL COUNSELL
Clinical research fellow
HAZEL FRASER
Administrator,
Cochrane Collaboration Stroke Review Group

Neurosciences Trials Unit,
Department of Clinical Neurosciences,
University of Edinburgh,
Western General Hospital NHS Trust,
Edinburgh EH4 2XU

- 1 Dickersin K, Scherer R, Lefebvre C. Systematic reviews: identify relevant studies for systematic reviews. *BMJ* 1994; 309:1286-91. (12 November.)

Pressure sores

Clinical trials best way of assessing different mattresses

EDITOR,—R K Vohra and C N McCollum's review on pressure sores contains two misconceptions.¹ Firstly, the authors state that measurement of the interface pressure is the best method of comparing the efficacy of pressure relieving supports, whereas clinical trials have shown that it is a poor indicator.^{2,3} Deep periosteal pressures are considerably higher than the interface pressure,⁴ and animal studies have repeatedly shown that the initial ischaemic necrosis that causes deep sores occurs in subcutaneous tissues, not in the skin.⁵

Secondly, the authors share the common confusion concerning the different actions of low pressure and alternating pressure supports: "in a comparison of alternating air, static air, and water mattress overlays on sacral and heel pressures . . . mean pressures were significantly higher for the alternating air mattress than the other surfaces; they should therefore be avoided."

Low pressure mattresses are soft supports that aim at distributing the weight as widely as possible and thus at preventing high pressures over bony prominences, which cause distortion of tissue and ischaemia. In contrast, alternating pressure mattresses are designed to be sufficiently firm to lift the patient off the bed and to support him or her while adjacent cells inflate and deflate underneath the body, constantly changing the areas of high pressure. They mimic the alternate high and low pressures that occur in normal people as a result of changes of position in response to pain due to pressure, which permit reactive hyperaemia and reoxygenation of the tissues and thus prevent ischaemic necrosis. Averaging the pressures in alternating pressure mattresses to enable them to be compared with low pressure supports is therefore meaningless.

Only clinical trials can show which system works best for different types of patient. A recent randomised trial in a district general hospital comparing alternating pressure mattresses with similarly priced constant low pressure mattresses (for example, fibre fills, slit foam, static air, water, and low air loss overlays) showed the alternating pressure mattresses to be considerably more effective.¹

M R BLISS
Consultant geriatrician

Homerton Hospital,
London E9 6SR

- 1 Vohra RK, McCollum CN. Pressure sores. *BMJ* 1994;309:853-7. (10 October.)
- 2 Gebhardt KS, Bliss MR, Winwright PL. A randomised controlled trial to compare the efficacy and cost of alternating pressure and constant low pressure supports for preventing pressure sores in hospital [abstract]. *Age Ageing* 1994;23(suppl 2):9.
- 3 Bliss MR. A randomised controlled trial of seven pressure relieving mattress overlays for preventing pressure sores in elderly patients [abstract]. *Age Ageing* 1994;23(suppl 2):20.
- 4 Le KM, Madsen BL, Barth PW, Ksander GA, Angell JB, Vistnes LM. An in depth look at pressure sores using monolithic silicon pressure sensors. *Plast Reconstr Surg* 1984; 74:745-54.
- 5 Salcido R, Donorrio JC, Fisher SB, Le Grand EK, Dickey K, Carney JM, et al. Histopathology of pressure ulcers as a result of sequential computer controlled pressure sessions in a fuzzy rat model. *Advances in Wound Care* 1994;7(5):23-40.

Carers should provide informed, cohesive approach

EDITOR,—I hope that R K Vohra and C N McCollum's review on pressure sores will be read by the people who chair curriculum committees of medical schools.¹ Pressure sores are seldom included as a specific topic for instruction to

medical students. This accounts for widespread ignorance on the subject among doctors both in hospitals and in the community. It has always struck me as extraordinary that a condition that affects between 5% and 10% of all patients in hospital should not be a matter of top priority for teaching of medical students.

One point that the review fails to emphasise is the need to establish satisfactory preventive measures in the community before patients are discharged from hospital. One of the commonest causes of the high rate of recurrence of pressure sores is the failure of communication between carers in hospital and carers in the community.

As a surgeon with an interest in pressure sores, I seldom agree to close a pressure sore until I know that future prevention of the same sore is assured. This often means that special equipment has to be purchased and so proves expensive, but, in the long term, prevention of sores is much cheaper than treatment and the cost of providing the equipment for prevention is equivalent to that of only a two or three week stay in hospital. Demarcation disputes often delay the decision on who should fund the equipment: the hospital believes that the community should do so and the community believes that the hospital should. It would be in everybody's interest if each health authority set aside money for the provision of equipment for patients with pressure sores that could be called on by both hospitals and the community.

Finally, I agree with the authors that a wide range of topical dressings and applications is marketed with, usually, little evidence of efficacy. This can be summarised in the aphorism "it matters far more what you put a pressure sore on than what you put on a pressure sore."

H P HENDERSON
Consultant plastic and hand surgeon

Leicester Royal Infirmary NHS Trust,
Leicester LE1 5WW

- 1 Vohra RK, McCollum CN. Pressure sores. *BMJ* 1994;309: 853-7. (1 October.)

Assessing risk of suicide

Samaritans' scoring system helps develop judgment

EDITOR,—H G Morgan's article on the role of doctors in preventing suicide emphasises the need for training in assessing the risk of suicide and responding appropriately.¹ Samaritan volunteers in training are taught always to ask about suicidal feelings and to assess the risk in detail. A rough and ready scoring system is sometimes used to help develop sound judgment based on thorough inquiry; it is mainly useful for training purposes.

The scoring system was based partly on a booklet about assessing the risk of suicide published by the New York Suicide Prevention Center (which was much too detailed and elaborate for day to day use) and partly on advice from the Los Angeles Suicide Prevention Center, which emphasised the importance of whether there is a suicide plan and, if so, its nature and intended timing in the assessment of the immediate risk. Sudden deaths (by hanging, shooting, or jumping), which are final, are distinguished from slower deaths (by overdosing, for example), in which rescue is possible if the chosen place and time favour it. Other risk factors—historical, social, and medical—are then scored as features that increase the immediate risk or warn of longer term risk for those not in imminent danger (table). The table thus offers reminders about what to ask: Any plan? What? When? Where? Are the means available? Ever tried before? How seriously? Preparations (making a will, giving things away, etc)?

The score that results offers guidance for an appropriate response: a score of 7 or 8 in the first

part of the table or of ≥ 20 in total indicates that the person is in imminent danger and should not be left alone; 14-19 (high risk) indicates that another meeting should be fixed soon; 6-11 (moderate risk) indicates that another meeting should be fixed; and 1-5 (slight risk) or 0 in the first part of the table (no suicidal intent) indicates that the volunteer should hear the person out and let him or her go unless there are other reasons to meet again.

Scoring table used to assess risk of suicide

Choose one figure to score suicide plan	
Chief indicator of immediate risk:	
Imminent sudden death	8
Imminent slow method of suicide	7
Future sudden death planned	6
Future slow method of suicide planned	5
Planning suicide "gamble"	4
Planning suicidal gesture	3
Definite suicidal thoughts but no plan	2
Toying vaguely with idea of suicide	1
No suicidal thoughts	0
Add points for every relevant item, mostly the long term factors	
Previous suicidal acts or gestures	≤ 4
Recent broken relationship	3
Isolation. Rejection	3 (each)
No hope. Loss of faith	3 (each)
Depressive illness	3
Dependence on alcohol or drugs	2
Possession of means of suicide	2
Putting affairs in order	2
Over 60. Male. Ill. Chronic pain	1 (each)

The volunteers then apply the table to clinical examples (described or used in role play), assess the risk, and select the appropriate response. The method is crude, and precision is not claimed; for example, some workers might rate depressive illness higher than items such as isolation or rejection. When applied to the clinical examples, however, the scoring table produces results that look fairly convincing, and as a training tool it works well.

ROY M VINING
General practitioner

Lowestoft,
Suffolk NR33 0BQ

1 Wilkinson G, Morgan HG. Can suicide be prevented? *BMJ* 1994;309:860-2. (1 October.)

Deal with self harm in prisons

EDITOR,—There seems to be sense in what both Greg Wilkinson and H G Morgan say about the prevention of suicide.¹ Setting targets for reducing suicide is a facile and superficial way of attempting to measure the efficacy of mental health services, ignoring as it does the huge sociocultural influences over this behaviour. In an age of mass unemployment and social disintegration the concept of "anomie" as described by Durkheim a century ago is surely relevant.²

On the other hand, we should not ignore the situations in which it may be possible to prevent suicide. A recent study has shown that suicide rates in young men aged 15 to 19 have risen in recent years.³ While the authors suggest that factors such as unemployment and poverty may contribute to this, some of these deaths occur in custodial settings. Suicide rates in prison are several times those in the general population. Judge Tumim has drawn attention to this and identified factors in the prison environment that could heighten the risk for vulnerable prisoners.⁴ Unsympathetic regimes with poor training of staff and low morale can have a deleterious effect, and their responses to self harm can be counterproductive and harmful. Morgan is surely right in emphasising the importance of establishing a therapeutic alliance with those at risk of self harm and suicide, but the absence of this is apparent in many custodial settings. A disproportionate number of acts of self harm in prisons occur among young prisoners. Liebling suggested that self harm in prisoners often occurs in a "poor coping" group, who find

adjustment to prison life difficult.⁵ Factors such as bullying may be particularly important in young prisoners.

To deal with self harm in prisons the approach must move beyond the identification of those at risk. Addressing the factors in the prison environment that precipitate suicide attempts and the attitudes of staff and prisoners to this problem could be highly beneficial. Allied with an increased emphasis on "certain basic clinical skills: a sympathetic ear . . . a quiet confidence," this might lead to safer custodial care of a highly vulnerable group of young people. This is an important and pressing social issue worthy of further study.

R P ROWLANDS
Senior registrar in psychiatry

Middlewood Hospital,
Sheffield S6 1TP

- 1 Wilkinson G, Morgan HG. Can suicide be prevented? *BMJ* 1994;309:860-2. (1 October.)
- 2 Durkheim E. *Suicide*. London: Routledge and Kegan Paul, 1970. (First published 1897.)
- 3 McClure GMG. Suicide in children and adolescents in England and Wales 1960-1990. *Br J Psychiatry* 1994;165:510-4.
- 4 Home Office. *Report of a review by her majesty's chief inspector of prisons for England and Wales of suicide and self-harm in prison establishments in England and Wales*. London: HMSO, 1990.
- 5 Liebling A. *Suicides in prison*. London: Routledge, 1992.

Antidepressants and suicide

EDITOR,—Erkki Isometsä and colleagues¹ supported the main conclusion of our toxicological study of suicides over two years in Sweden²—namely, that underprescribing and therapeutic failure are greater problems than toxicity with antidepressants. Isometsä and colleagues studied 1397 suicides in Finland during 1987-8; 57% of the victims were depressed. Antidepressants were prescribed to only a third of those with major depression (3% at adequate doses). Altogether 1348 subjects were investigated toxicologically in the same Finnish study, and Erkki Vuori and colleagues confirm our toxicological finding that less than 16% of the patients who committed suicide were taking antidepressants at the time of death.³ They report that antidepressants were detected in 38 of the 1083 people who committed suicide by means other than self poisoning. If all the remaining 265 people were taking antidepressants, a maximum proportion of 22% ((38+265)/1348) is reached. As, however, only a third of the subjects with major depression had been prescribed antidepressants, it is likely that less than a third of the 265 who died of self poisoning were positive for antidepressants, which suggests that at most 9% of the Finnish patients who committed suicide were being treated with antidepressants at the time of death.

Vuori and colleagues have doubts about the validity of our "post-mortem study on antidepressant effect." A weaker antidepressant effect is one possible explanation of our finding that moclobemide and mianserin had a higher association with suicide than did other antidepressants. In the case of moclobemide this is supported by results from prospective clinical trials.⁴ We have discussed alternative explanations in a previous reply to correspondence⁵ and in a paper.⁶ The purpose of studies like ours, however, is to provide early signals on risks with new drugs (for example, rare interactions, sudden deaths, suicides, and therapeutic failures, which are seldom found in pre-marketing trials or reported in spontaneous adverse reporting systems).⁷⁻⁹ A meta-analysis of the clinical trials of fluoxetine¹⁰ was of too low statistical power to be able to detect a possible 50% increased risk of suicide.¹¹ That would have required 42 000 subjects. Alternative approaches have therefore been proposed for postregistration surveillance.^{7,11} Results from non-experimental studies must, however, be interpreted cautiously and validated by further studies.⁵ Also, experi-

mental studies, which are often unable to show differences in effect, must be interpreted critically with regard to their low power (type II error).^{2,4,11}

GÖRAN ISACSSON
Assistant chief physician

Department of Clinical Neuroscience and Family Medicine,
Division of Psychiatry, Karolinska Institute,
Huddinge University Hospital,
Huddinge,
Sweden

PER HOLMGREN
Chemist

Department of Toxicology,
National Laboratory of Forensic Chemistry,
Linköping,
Sweden

DANUTA WASSERMAN
Head

Centre for Suicide Research and Prevention,
Karolinska Hospital,
Stockholm,
Sweden

UIP BERGMAN
Chief physician

Department of Medical Laboratory Sciences and Technology,
Division of Clinical Pharmacology, Karolinska Institute

- 1 Isometsä E, Henriksson M, Heikkinen M, Aro H, Lönnqvist J. Suicide and the use of antidepressants. *BMJ* 1994;308:915. (2 April.)
- 2 Isacsson G, Holmgren P, Wasserman D, Bergman U. Use of antidepressants among people committing suicide in Sweden. *BMJ* 1994;308:506-9. (19 February.)
- 3 Vuori E, Lönnqvist J, Klaukka T. Antidepressants and suicide. *BMJ* 1994;308:1573. (11 June.)
- 4 Gram LF, Isacsson G, Bergman U. Moclobemide. *Lancet* 1994;343:679-80.
- 5 Isacsson G, Holmgren P, Wasserman D, Bergman U. Suicide and the use of antidepressants. *BMJ* 1994;308:916. (2 April.)
- 6 Isacsson G, Redfors I, Wasserman D, Bergman U. Choice of antidepressants: questionnaire survey of psychiatrists and general practitioners in two areas of Sweden. *BMJ* 1994;309:1546-9. (10 December.)
- 7 Bergman U. Pharmacoepidemiological perspectives. *J Clin Epidemiol* 1992;45(3):313-7.
- 8 Henry JA. The safety of antidepressants. *Br J Psychiatry* 1992;160:439-41.
- 9 Mehtonen O-P, Aranko K, Mäklonen L, Vapaatalo H. A survey of sudden death associated with the use of antipsychotic or antidepressant drugs: 49 cases in Finland. *Acta Psychiatr Scand* 1991;84:58-64.
- 10 Beasley CM Jr, Dronseif BE, Bosomworth JC, Saylor ME, Rampey AH Jr, Heiligenstein JH, et al. Fluoxetine and suicide: a meta-analysis of controlled trials of treatment for depression. *BMJ* 1991;303:685-92.
- 11 Li-Wan-Po A. Fluoxetine and suicide: meta-analysis and Monte-Carlo simulations. *Pharmacoepidemiology and Drug Safety* 1993;2:79-84.

Patient confidentiality in New Zealand

EDITOR,—Charles Essex's news article concerning the confidentiality of patients' medical files in New Zealand is a one sided representation of the facts.¹ The Northern Region Health Authority in New Zealand is responsible for paying various subsidies to medical practitioners in its region. Health Benefits Limited is the agent of the regional health authority that makes these payments and also investigates any apparent inappropriate claims. The statutory authority under which Health Benefits Limited conducts these investigations is section 22G of the Health and Disability Services Act. This section provides that records of claiming practitioners may be inspected and copied to determine whether claims have been made correctly or otherwise.

In many cases certain types of claims may not be made in conjunction with other claims for the same services. It is, however, occasionally necessary to view a patient's records to ascertain whether double claiming is justified. This particularly occurs, for example, when a claim under the maternity benefit schedule is made at the same time and for the same patient as a claim under the general medical services benefit schedule. In the case referred to in Essex's article, claiming patterns for the practitioner concerned seemed inappropriate to both the region and Health Benefits Limited. The police have laid charges against the practitioner, who in due course will appear in court.