a direct approach—for example, isolation or molecular identification of a potentially infecting organism from thyroid gland and injection into an experimental animal to test for possible induction of autoimmune thyroid diseases.

ORIANA LERI
Researcher
MARIA TERESA SINOPOLI
Doctor
MARIA ANNA DI PRIMA
Researcher

Department of Infectious and Tropical Diseases,
University "La Sapienza",
Rome,
Italy

ANNA PAGGI
Associate professor

Department of Internal Medicine,
University "La Sapienza"

- 1 Duclos Vallée JC, Johanet C, Trinchet JC, Deny P, Laurent MF, Duron F, *et al.* High prevalence of serum antibodies to hepatitis C virus in patients with Hashimoto's thyroiditis. *BMJ* 1994;309:846-7. (1 October.)
- 2 Tran A, Quaranta JF, Beusnel C, Thiers V, De Souza M, François E, *et al.* Hepatitis C virus and Hashimoto's thyroiditis. *Eur J Med* 1992;1:116-8.
- 3 Tomer Y, Davies TF. Infection, thyroid disease and autoimmunity. *Endocr Rev* 1993;14:107-20.
- 4 Paggi A, Di Prima MA, Paparo BS, Pellegrino C, Faralli AR, Sinopoli MT, *et al.* Anti 70 kDa heat shock protein antibodies in sera of patients affected by autoimmune and non autoimmune thyroid disease. *Endocr Res* (in press).

Enalapril and β blockers in chronic renal failure

EDITOR,—T Hannedouche and colleagues report a randomised controlled trial of enalapril and β blockers in patients with non-diabetic chronic renal failure.¹ The importance of hypertension and progression of renal insufficiency is well established.^{2,3} The authors' conclusion that enalapril was superior to β blockers in preventing end stage renal failure in patients with chronic renal insufficiency was partly based on the statement that "no differences in blood pressure were seen at any point of the follow up between the two groups." Table III of their paper, however, shows blood pressure only at one year intervals for a study consisting of three year follow up.

Table IV shows that the patients in the conventional group had a significantly lower incidence of adequate control of blood pressure when taking only one drug—namely, the β blocker. As far as we can tell, additional antihypertensive drugs were added at three month intervals. The conventional group may thus have had poorer control of blood pressure for the first six to nine months of follow up. We wonder if the authors can provide comparisons of blood pressure, especially during the first year of follow up, at three month intervals. Without this information one cannot exclude better control of blood pressure in the enalapril group as the explanation for the drug's beneficial effect. This is especially true as the length of follow up is only three years and even a three to six month period of better control may alter the rate of progression of renal disease.

DERYA BORA HAZAR
Renal fellow
LI TSO
Renal fellow

Renal Unit, Massachusetts General Hospital,
Jackson 825, Boston,
MA 02114,
USA

- 1 Hannedouche T, Landais P, Goldfarb B, El Esper N, Fournier A, Godin M, *et al.* Randomised controlled trial of enalapril and β blockers in non-diabetic chronic renal failure. *BMJ* 1994;309:833-7. (1 October.)
- 2 Brazy PC, Stead WW, Fitzwilliam JF. Progressive renal disease: role of race and antihypertensive medications. *Kidney Int* 1990;37:1113-9.
- 3 Hannedouche T, Albouze G, Chauveau P, Lacour B, Junger P. Effects of blood pressure and antihypertensive treatment on progression of advanced chronic renal failure. *Am J Kidney Dis* 1993;21(suppl 2):131-7.

Computed tomography for single seizures

EDITOR,—Ronald A Schoenenberger and Sabine M Heim suggest that cerebral imaging after a single seizure may be unnecessary if a score of ≥ 3 is elicited with the CAGE questionnaire (a screening questionnaire on alcohol misuse).¹ Their study raises several related issues that merit further comment.

Focal cerebral lesions thought to be responsible for the presenting seizures were found in 34% of cases. In these cases there was presumably a focal onset before generalisation, which was not evident on ictal clinical assessment. It follows, therefore, that the clinically generalised seizure was in fact a partial seizure with secondary generalisation in an appreciable minority of cases. This fact alone should lower thresholds for imaging in this clinical scenario.

While I accept the validity of the CAGE questionnaire, using it after obtaining a detailed alcohol and drug history, as should be the case, may elicit unreliable answers, especially in a period of postictal confusion. False positive and false negative responses must occur. Moreover, if a seizure is precipitated by alcohol withdrawal (and this is not synonymous with a score of ≥ 3 on testing the CAGE questionnaire) there are immediate clinical implications, not the least of which is the possible need for detoxification.

The possibility of behavioural disturbance during this period is high. Given the frequent interrelation among cerebral trauma, disordered haemostasis, and alcohol misuse, the ratio of risk to benefit in this subgroup of patients with a single seizure seems to be strongly in favour of cerebral imaging irrespective of scores obtained with the CAGE questionnaire.

J R C BOWEN
Neurology registrar

Princess Alexandra's Hospital,
Wroughton,
Wiltshire SN4 0QJ

- 1 Schoenenberger RA, Heim SM. Indication for computed tomography of the brain in patients with first uncomplicated generalised seizure. *BMJ* 1994;309:986-9. (15 October.)

Seasonal variations in fatal pulmonary embolism

Several mechanisms contribute

EDITOR,—Peter Wilmschurst draws attention to the increased incidence of coronary heart disease and stroke associated with cold weather.¹ This conclusion is drawn largely from epidemiological data from studies that were not based on postmortem examinations. We are surprised that he does not mention another major cause of cardiovascular mortality, pulmonary embolism.

Fatal pulmonary embolism causes over 20 000 deaths annually in Britain and accounts for a tenth of all deaths in hospital.² It also has a seasonal variation, with both postoperative and "spontaneous" emboli being significantly more common in winter.^{3,4} The cause of many deaths due to pulmonary embolism may be registered as myocardial infarction in the absence of routine postmortem examination, and this may contribute to the seasonal increase in deaths registered as due to ischaemic heart disease.

The seasonal increase in fatal pulmonary embolism may partly be explained by the changes in coagulation described in the editorial, but other factors, such as peripheral vasoconstriction and reduced activity resulting in a fall in blood flow in the lower limbs, may also contribute.⁵ We agree with Wilmschurst that cold weather causes an increase in preventable cardiovascular deaths. The

mechanisms of this increased death rate may include an increase in numbers of cerebrovascular, coronary, and thromboembolic episodes. In the absence of a large study based on postmortem findings it is difficult to determine the contribution of each mechanism and the preventive measures required.

E A COOKE
Research fellow
M A McNALLY
Senior registrar
R A B MOLLAN
Professor of orthopaedic surgery

Department of Orthopaedic Surgery,
Queen's University of Belfast,
Musgrave Park Hospital,
Belfast BT9 7JB

- 1 Wilmschurst P. Temperature and cardiovascular mortality. *BMJ* 1994;309:1029-30. (22 October.)
- 2 Sandler DA, Martin JF. Autopsy proven pulmonary embolism in hospital patients: are we detecting enough deep vein thrombosis? *J R Soc Med* 1989;82:203-5.
- 3 Wroblewski BM, Siney PD, White R. Fatal pulmonary embolism after total hip arthroplasty. Seasonal variation. *Clin Orth* 1992;276:222-4.
- 4 Gallerani M, Manfredi R, Ricci L, Grandi E, Cappato R, Calo G, *et al.* Sudden death from pulmonary thromboembolism: chronobiological aspects. *Eur Heart J* 1992;13:661-5.
- 5 McNally M, Mollan RAB. Total hip replacement, lower limb blood flow and venous thrombogenesis. *J Bone Joint Surg [Br]* 1993;75:640-4.

Malignant phase hypertension does not vary by season

EDITOR,—In his editorial on temperature and cardiovascular mortality Peter Wilmschurst comments that the difference in temperature between summer and winter in Britain should result in a difference in blood pressure of 5 mmHg.¹ Malignant phase hypertension represents a syndrome of high blood pressure that is associated with a high mortality if left untreated. We recently reported that we had not noted a decline in the incidence of this condition between 1970 and 1993.²

To investigate whether the presentation of malignant phase hypertension varied by season we reviewed the month of presentation of 315 patients (211 men, 104 women; mean age 49.4 (SD 12.7) years) with the condition who were referred to our hospital between 1965 and 1994. There was no clear relation between the number presenting and season, although the largest number of patients first presented in March (n=35) and the lowest number of patients first presented in August (n=17) (figure). When patients who presented in March were compared with those who presented in August there were no significant differences in mean age (49.6 (13.3) v 46.2 (8.3) years; unpaired *t* test, *P*=0.27), mean duration of known hypertension before presentation (17.5 (43.8) v 6.4 (14.7) months; *P*=0.18), and mean diastolic blood pressure (144.3 (19.8) v 140.0 (30.5) mmHg; *P*=0.61). There was, however, a trend towards a high mean systolic blood pressure (234.2 (28.0) v

Number of patients presenting with malignant phase hypertension by month (n=315)