colleagues' study among the total migrant population of 54 was black (n=38); this group included Caribbean (n=19) and black African (n=14) subjects. Countries with a majority black population do not feature at all in the study most quoted when similar worldwide prevalences of schizophrenia are claimed.² Limited data exist on prevalence in some African and Caribbean countries, but these studies do not use standardised criteria. Because the provision of services is different in the developing world underreporting may well occur.^{3,4}

As the incidence of schizophrenia in Britain is thought to be declining⁵ it is important that studies of migrant populations relate to the countries of origin as well as the host countries. While I am not trying to suggest that biological factors can fully explain the increased incidence of psychosis in migrants, I do not think that they can be ignored.

NICHOLAS AIREY
Clinical medical officer

Tone Vale Hospital,
Norton Fitzwarren,
Somerset TA4 1DB

- 1 King M, Coker E, Leavey G, Hoare A, Johnson-Sabine E. Incidence of psychotic illness in London: a comparison of ethnic groups. *BMJ* 1994;309:1115-9. (29 October.)
- 2 Sartorius N, Jablensky A, Korten A, Ernberg G, Anker M, Cooper J, et al. Early manifestations and first-contact incidence of schizophrenia in different cultures. *Psychol Med* 1986;16: 909-28.
- 3 Fuller Torrey E. Prevalence studies in schizophrenia. *Br J Psychiatry* 1987;150:598-608.
- 4 Burke A. First admissions and planning in Jamaica. *Social Psychiatry* 1974;9:39-45.
- 5 Der G, Gupta S, Murray R. Is schizophrenia disappearing? *Lancet* 1990;335:513-6.

Conclusions of study are unwarranted

EDITOR,—Michael King and colleagues draw some unwarranted conclusions from their study on psychosis in ethnic groups.¹ On the basis of their finding of high rates of both schizophrenia and affective psychoses in black and Asian patients they suggest that "the current focus on African Caribbeans and schizophrenia [is] misleading." High rates of serious mental illness in British African Caribbean people have been noted in studies over three decades, and this has often included affective illness.^{2,3} In contrast, no consistent picture has emerged for Asians. The authors rightly avoid multiple statistical comparisons of small numbers but then extrapolate from their results specific conclusions about African Caribbeans with schizophrenia, of whom there were just nine in their sample.

In discussing the limitations of their work King and colleagues imply that knowledge of the predictive validity of psychiatric diagnosis in ethnic minorities awaits the completion of their planned follow up study. They may not be aware that, for African Caribbean patients with schizophrenia, the predictive validity has already been established.⁴

The phrase "all ethnic minority groups are vulnerable" is offered as a clinical implication; worst of all, the paper concludes that the "personal and social pressures of belonging to any minority group in Britain are important determinants of the excess of psychotic disorders." These views may be attractive to many commentators, but the latter in particular remains without foundation. While some widespread environmental factor seems to precipitate extra cases of schizophrenia in African Caribbean families at risk,⁵ circumstantial evidence points to prenatal viral infection.⁶ At present no empirical data or plausible mechanism exist to relate racial prejudice and disadvantage to population rates of psychosis.

PHILIP A SUGARMAN
Senior lecturer in forensic psychiatry

United Medical and Dental Schools of
Guy's and St Thomas's Hospitals,
London SE1 7EH

- 1 King M, Coker E, Leavey G, Hoare A, Johnson-Sabine E. Incidence of psychotic illness in London: comparison of ethnic groups. *BMJ* 1994;309:1115-9. (29 October.)
- 2 Hemi LK. Psychiatric morbidity of West-Indian immigrants. *Social Psychiatry* 1967;2:95-100.
- 3 Harrison G, Owens D, Holton A, Neilson D, Boot D. A prospective study of severe mental disorder in Afro-Caribbean patients. *Psychol Med* 1988;18:643-57.
- 4 Sugarman PA. Outcome of schizophrenia in the Afro-Caribbean community. *Soc Psychiatry Psychiatr Epidemiol* 1992;27:102-5.
- 5 Sugarman PA, Craufurd D. A family history study of schizophrenia in the Afro-Caribbean community. *Br J Psychiatry* 1994;164:474-80.
- 6 Eagles JM. The relationship between schizophrenia and immigration. Are there alternatives to psychosocial hypotheses? *Br J Psychiatry* 1991;159:783-9.

Social observations on childhood

EDITOR,—As the producer of Clifford Yorke's talks on BBC Radio 3, *Childhood and Social Truth*, I wish to respond to Mark Berelowitz's review.¹ Berelowitz says that Yorke does not explain why the children of stable families have become unstable parents. In fact, his talks concentrate on answering this question. The reason is that parents often fall victim to fashionable ideas and social pressures and therefore act in ways that are contrary to their good understanding of children's needs. Berelowitz is disappointed that the talks do not offer new theories of childhood. This shows that he has missed the point, which is that we should pay much more attention to what we know.

Berelowitz misrepresents the contents of the talks. Yorke does not say that it hardly matters whether a child suffers from divorce or marital disharmony. He makes it clear that divorce is likely to be much worse. He certainly does not say that the whole of parenting consists of teaching the child to manage frustration. He mentions frustration to show how regressive "doing your own thing" is. Yorke gives a vivid picture of the harmony with oneself and others to which most parents aspire.

Many criticisms in the review are of the brief for the series rather than its contents. Berelowitz thinks that the talks should have been commissioned to "encourage parents"; to give a blueprint for reversing poverty, illness, and poor schooling in Britain and pestilence, war, and famine abroad. He wanted more treatment of the role of fathers within a stable household. But there are already plenty of manuals for parents and politicians. I commissioned a set of social observations by an authority on normal child development. The purpose of the broadcasts was to stimulate thought among listeners, and that thought was to be about what individuals can do to mitigate the weaknesses of society.

It is not easy to understand why a doctor should be less able than some dozen newspaper journalists to digest and summarise Yorke's arguments correctly.

LOUISE GREENBERG
Chief producer, arts, science, and features

BBC Radio,
London W1A 1AA

- 1 Berelowitz M. Everybody's doing it wrong. *BMJ* 1994;309:1309. (12 November.)

Circumcision

EDITOR,—Disappointingly, the discussion concerning the personal view¹ and book² on the joy of uncircumcising failed to emphasise the central issue with a clear statement that the practice of social circumcision is unacceptable in a civilised society. It is unacceptable not because of the increased surgical risks,² surgical complications,³ harmful long term outcome,⁴ difficulty in reversing the process,¹ or increased sexual dysfunction⁵ but

primarily because systematic ritual genital mutilation is both unnecessary and unjust.

Parents pursuing personal, social, and religious beliefs may request such an operation, and surgeons may then carry it out seemingly without question and sometimes for not entirely altruistic reasons. When such a permanently mutilating procedure is proposed and performed without clinical need there seems to be no individual culpability; surely a systematic deformation on any other part of the body would have been deemed to be a form of ritual child abuse. Furthermore, society has traditionally turned a blind eye to this problem, and seldom is any comment or criticism made by politicians and parliamentarians; social pundits and religious gurus; organisations entrusted with the welfare of children and the prevention of cruelty; or medical mandarins in the higher echelons of the General Medical Council, Royal Colleges of Surgeons and Physicians, and BMA.

Is it not time for all members of our society to reflect on this issue and to give a considered judgment on whether social circumcision could be deemed justifiable? While the jury is out—and it could be out for a long time—the least we can do is to stop performing such operations on children until they are old and mature enough to give informed consent.

S MULICK
Consultant in rehabilitation medicine

Department of Rehabilitation Medicine,
Luton and Dunstable Hospital NHS Trust,
Luton LU4 0DZ

- 1 The joy of uncircumcising. *BMJ* 1994;309:676-7. (10 September.)
- 2 Kirby RS. The Joy of Uncircumcising! Restore your birthright and maximize sexual pleasure [book review]. *BMJ* 1994;309: 679. (10 September.)
- 3 Eason JD, McDonnell M, Clark G. Male ritual circumcision resulting in acute renal failure. *BMJ* 1994;309:660-1. (10 September.)
- 4 Warren J. Circumcision and uncircumcision. *BMJ* 1994;309: 1020. (15 October.)
- 5 Bigelow J. Circumcision and uncircumcision. *BMJ* 1994;309: 1160. (29 October.)

Triglyceride concentration and coronary heart disease

EDITOR,—We wish to respond to two letters concerning our paper.^{1,2} Despite the intuitive appeal of lowering triglyceride concentrations along with cholesterol concentrations when both are raised, there is no evidence that this strategy is more effective than targeting treatment at cholesterol alone. Even treatments that raise triglyceride concentrations reduce the risk of coronary heart disease in hypercholesterolaemic subjects. In the Lipid Research Clinics coronary primary prevention trial, treatment with diet and cholestyramine resulted in a 19% reduction in the risk of coronary heart disease despite an increase in serum triglyceride concentrations averaging 2.5%.³ Oestrogen treatment is also associated with increased triglyceride concentrations and a reduced risk of coronary heart disease.⁴ Clinical trials have not established which subgroups of hypercholesterolaemic patients, if any, benefit from a reduction in triglyceride concentrations.

A M Cruickshank notes that the rare patient with severe hypertriglyceridaemia is at risk of pancreatitis⁵; this association, however, is insufficient justification for universal screening of triglyceride concentrations. Certainly, screening would detect some hypertriglyceridaemic subjects who would otherwise be overlooked. But severe hypertriglyceridaemia, in the range associated with pancreatitis, is usually obvious from its physical manifestations and visual inspection of the lipaemic blood specimen. Seldom will the result of a screening test be the sole clue to a very high triglyceride concentration.