consumption carries no benefit in terms of mortality. Pending further studies, it would be prudent from a health promotion perspective not to advocate alcohol consumption as a procedure for extending life but to assume that the dose-response relation for mortality and alcohol consumption is linear with a positive slope.

DAVID F MARKS
Professor of psychology

Health Research Centre,
School of Psychology,
Middlesex University,
Enfield,
Middlesex EN3 4SF

1 Doll R, Peto R, Hall E, Wheatley K, Gray R. Mortality in relation to consumption of alcohol: 13 years' observations on male British doctors. *BMJ* 1994;309:911-8. (8 October.)
2 Cox BD, Blaxter M, Buckle ALJ, Fenner NP, Golding JF, Gore M, et al. *The health and lifestyle survey*. London: Health Promotion Trust, 1987.

Moderate drinking also improves health

EDITOR,—Richard Doll and colleagues convincingly show the beneficial effect of mild to moderate alcohol consumption on the risk of death from ischaemic heart disease and from several other causes.¹ No published report seems to exist, however, on the effect of sensible alcohol consumption on health status. We describe the relation between alcohol use and quality of life related to health, based on the findings of a survey that we undertook in West Glamorgan in 1993.

The method and main findings of the study, including the aging effect of smoking, have been fully described elsewhere.^{2,3} In essence, a random sample of adults (n=827, response rate 82%) were interviewed in their own home by trained health visitors who used a structured questionnaire. This contained the short form 36 and a variety of questions on smoking habits, alcohol intake, and history of chronic illness. On alcohol consumption, participants were asked to indicate whether they drank alcohol and, specifically, whether they had consumed any alcohol in the past week. Subjects who answered "yes" were further prompted to quantify their consumption. Four categories of drinkers were derived: non-drinkers; mild drinkers (≤ 14 units a week for women and ≤ 21 units a week for men); moderate drinkers (15-34 units a week and 21-49 units a week respectively); and severe drinkers (> 34 units a week and > 49 units a week respectively).

Of the 827 participants, 337 were non-drinkers, 403 mild drinkers, 66 moderate drinkers, and 21 severe drinkers. Mean scores obtained with the short form 36 in the four categories were determined for each of the eight parameter scores, and the differences between drinkers and non-drinkers were compared. Multiple regression was used to adjust for age, sex, social class, smoking habits, and the presence of chronic illness.

Our results showed that the mild to moderate drinkers reported a significantly better health

Mean difference in SF-36 scores between mild drinkers and non-drinkers and between moderate drinkers and non-drinkers (adjusted for age, sex, social class, smoking habits, and presence of chronic illness)

	Mild drinkers v non-drinkers	Moderate drinkers v non-drinkers
Physical functioning	4.7**	8.7***
Role limitations due to physical problems	6.4**	10.0**
Bodily pain	4.9**	2.5
General health perception	3.7**	1.5
Vitality	3.3*	2.9
Social functioning	6.9***	0.7
Role limitations due to emotional problems	2.2	5.2
Mental health	1.1	0.1

*P<0.1, **P<0.05, ***P<0.01.

experience than did the non-drinkers: they perceived themselves to be more physically active and to experience fewer role limitations due to physical problems (table). In addition, the mild drinkers experienced significantly less bodily pain and perceived themselves to be more healthy, to have more vitality, and to be generally more sociable than did the non-drinkers. We have not presented the results for the severe drinkers because the sample was too small to yield any reliable estimate.

Thus mild to moderate alcohol consumption seems to confer not only longevity but also a better quality of life related to health. The biblical advice by the apostle Paul to Timothy urging him to stop drinking only water and start using a little wine⁴ is thus scientifically sound and valid. The implication for prevention is not to pursue a population strategy aimed at lowering the average alcohol consumption as this is likely to produce less health gain (and may even produce health loss) but to target preventive efforts at high risk drinking—that is, pursue the high risk strategy.⁵

RONAN A LYONS
Consultant
SU VUILO
Senior registrar
STEPHEN MONAGHAN
Senior registrar
BEVERLEY N C LITTLEPAGE
Director

Department of Public Health Medicine,
West Glamorgan Health Authority,
Swansea SA1 5AQ

1 Doll R, Peto R, Hall E, Wheatley K, Gray R. Mortality in relation to consumption of alcohol: 13 years' observation on male British doctors. *BMJ* 1994;309:911-8. (8 October.)
2 Lyons RA, Lo SV, Littlepage BNC. Perception of health amongst ever-smokers and never-smokers: a comparison using the SF-36 health status questionnaire. *Tobacco Control* 1994;3: 213-5.
3 Lyons RA, Lo SV, Littlepage BNC. Comparative health status of patients with 11 common illnesses in Wales. *J Epidemiol Community Health* 1994;48:388-90.
4 Holy Bible. 1 Timothy v, 23. (New International Version.)
5 Rose G. *The strategy of preventive medicine*. Oxford: Oxford University press, 1992.

The conclusions we draw

EDITOR,—Richard Doll and colleagues' finding that all cause mortality in men is lowest among those who drink 21 units of alcohol a week can be interpreted differently depending on the observer's stance.¹

The teetotaler—It's so unfair!
The average drinker—The doctors were wrong: the safe limit isn't 21, it's 63. So back to life as normal: drink what you like—in moderation.

The heavy drinker—Drinking four and half pints of beer a night has about the same risk as not drinking at all. I like a few pints and don't mind a little extra risk, so I'll stick to six pints (or seven etc).

The insurance company—We must add an extra question to proposals for life insurance asking whether prospective customers drink 21 units of alcohol a week and if not why not. If they don't we'll weight their premiums.

The health fanatic—This is a better excuse than "rehydrating myself" for drinking one and a half pints of beer every night after a game of squash.

The standpoint of the authors seems to be that of the health fanatic, with length of life rather than quality and enjoyment being the prime objective. Most doctors whom I know would prefer the second viewpoint—that of the average drinker—and would drink within reason while accepting a risk no greater than that of teetotalers. Overall, the main conclusion that many people will draw from the study is that drinking alcohol is not as bad as some people say.^{2,3}

T M REYNOLDS
Consultant chemical pathologist

Burton Hospital,
Burton upon Trent,
Staffordshire DE13 0RB

1 Doll R, Peto R, Hall E, Wheatley K, Gray R. Mortality in relation to consumption of alcohol: 13 years' observations on male British doctors. *BMJ* 1994;309:911-8. (8 October.)
2 Department of Health. *Action against alcohol misuse*. London: HMSO, 1991.
3 Secretary of State for Health. *The health of the nation*. London: HMSO, 1992.

Length of femoral neck and rates of hip fractures

Measurements may not be reliable

EDITOR,—Too little attention has been given to the relation between anatomical structure and risk of fracture. Ian R Reid and colleagues report the relation between the length of the femoral neck and rates of hip fracture,¹ but they in fact measured the projected length of the femoral neck. This varies with the degree to which the femur is rotated around its shaft axis. In neutral position the neck usually points somewhat anteriorly and is thus shortened on the frontal projection. As the femur is rotated inward its projected length increases. Frontal radiographs of the hip are usually obtained with the femur rotated inward, but practice varies, and it cannot be assumed that identical projections were obtained in independent routine radiographs obtained with an interval of 40 years. The authors must therefore show that, on average, the projections of the femoral neck were identical in 1950 and 1990.

For correct measurement of the length the long axis of the femoral neck must be parallel to the film. For this, however, anteversion of the femoral neck must be corrected for. Because of the great variability of this, with a range of nearly 30° in most populations, correct positioning is impossible without three dimensional radiography. A computer program is now available for measuring the length of the femoral neck on dual x ray absorptiometry of the hip. This measurement is made with the foot rotated inward, but no individual differentiation is made for the degree of anteversion of the femoral neck. Therefore this measurement may not give a reliable individual assessment of the length of the femoral neck.

ARNE HØISETH
Radiologist

Sentrum Røntgeninstitut,
Oslo,
Norway

1 Reid IR, Chin K, Evans MC, Jones JG. Relation between increase in length of hip axis in older women between 1950s and 1990s and increase in age specific rates of hip fracture. *BMJ* 1994;309:508-9. (20-27 August.)

Author's reply

EDITOR,—I agree that the length of the hip axis measured from radiographs depends on femoral rotation and that this cannot be directly assessed from the plain radiographs used in our study. We were aware of this potential problem and excluded films that showed incorrect positioning or in which there was appreciable disease of the hip joint, which might have compromised hip positioning. During both periods pelvic radiographs were routinely obtained with the legs internally rotated. As Arne Høiseth points out, it is impossible to be sure that positioning was the same at two periods so widely separated in time, but there is no reason to believe that any change in radiographic technique occurred. Indeed, inspection of the two sets of films suggests that the degree of internal rotation was not different between the groups.

IAN R REID
Associate professor

Department of Medicine,
University of Auckland,
Auckland,
New Zealand