haemorrhage, but this investigation is not done when there are concerns about raised intracranial pressure. Where safe, a subdural tap should be done as the analysis of cerebrospinal fluid is helpful in the dating of injuries.

During the shaking episode the infant is often held by the thorax⁴ and the compression forces on the ribs may result in fractures. Alternatively, the child may be held by the shoulders and upper arms or feet.⁷ The squeezing and violent movement associated with the shaking may cause the typical fracture patterns of child abuse.⁸ Brain damage is often multifactorial with direct shaking injury to the brain being compounded by hypoxic and ischaemic injury, infarction, coning, and the pressure effects of subdural haematomas and impact trauma.⁷

Computed tomography in suspected cases of shaking injury is essential. Subdural haematomas, sometimes of different ages; subarachnoid blood; intracerebral and intraventricular bleeding; cerebral oedema; diffuse loss of differentiation between grey and white matter; and cerebral laceration and contusional tears may all be found in shaken infants.⁹ Skull radiology is essential as fractures may be missed on computed tomography.¹⁰ Magnetic resonance imaging supplements computed tomography by showing small subdural collections or identifying subdural haematomas of different ages and shaking injuries that are not visible on computed tomography.¹¹ Owing to its availability computed tomography remains the primary imaging procedure. More recently, high resolution ultrasonography performed through a patent fontanelle has been shown to be very sensitive in showing shearing injuries and subdural and subarachnoid fluid.¹² Late results of brain injury include multicystic encephalomalacia, obstructive or communicating hydrocephalus, cerebral atrophy, infarctions, and gliosis.

The most severe shaking injuries tend to occur in younger infants as the head is relatively large in relation to the body. As body weight increases and neck muscles strengthen the incidence of brain damage due to shaking falls and is rare after the second year of life.

The outcome depends on the severity of the shaking injury; morbidity and mortality are high when the infant is comatose on presentation.^{3,5} Long term sequelae include profound mental retardation, spasticity, motor dysfunction, blindness, convulsions, and hydrocephalus.⁵ In one series of children shaken to unconsciousness 60% died or had profound mental retardation, spastic quadriplegia, or severe motor dysfunction. Others who had convulsions, irritability, or

lethargy but no lacerations, cerebral infarction, or severely raised intracranial pressure had subtle neurological sequelae or persistent convulsions.¹³

Who shakes their child? The shaking may represent a response to tension and frustration generated by the infant's incessant crying, which may be exacerbated by ignorance of appropriate infant care. People experiencing stress may be more prone to impulsive and aggressive behaviour.¹⁴

Until now social and medical effort has focused mainly on diagnosing and treating the shaken infant, with little emphasis on prevention. Studies in the United States have shown that although between a quarter and a half of the public did not know of the dangers of shaking infants, they retained information given in an awareness campaign.¹⁵

We do not know how many infants are shaken in Britain each year, but over 100 deaths occur from child abuse or neglect and in many of these children death is due to brain injury. Increased public awareness of the dangers of shaking should reduce this number and reduce the disabilities of survivors.

HELEN CARTY
Consultant radiologist
JANE RATCLIFFE

Consultant in paediatric intensive care

Alder Hey Children's Hospital,
Liverpool L12 2AP

Drs Carty and Ratcliffe have been involved in the campaign to increase public awareness of the dangers of shaking babies.

- 1 Billmire ME, Myers PA. Serious head injury in infants: accident or abuse? *Pediatrics* 1985;75:340-2.
- 2 Caffey J. On the theory and practice of shaking infants: its potential residual effects of permanent brain damage and mental retardation. *Am J Dis Child* 1972;124:161-9.
- 3 Hadley MN, Sonntag VKH, Rekeate HL, Murphy A. The infant whiplash-shake injury syndrome: a clinical and pathological study. *Neurosurgery* 1989;24:536-40.
- 4 Kleinman P. Diagnostic imaging in infant abuse. *Am J Roentgenol* 1990;155:703-12.
- 5 Committee on Child Abuse and Neglect, American Academy of Pediatrics. Shaken baby syndrome: inflicted cerebral trauma. *Pediatrics* 1993;92:872-5.
- 6 Levin AV. Ocular manifestations of child abuse. *Ophthalmology Clinics of North America* 1990;3:249-64.
- 7 Brown JK, Minns RA. Non-accidental head injury, with particular reference to whiplash shaking injury and medico-legal aspects. *Dev Med Child Neurol* 1993;35:849-69.
- 8 Carty HML. Fractures caused by child abuse. *J Bone Joint Surg Br* 1993;75:849-57.
- 9 Han BK, Towbin RB, De Courten-Myers, McLaurin RL, Ball WS. Reversal sign on CT: effect of anoxic/ischemic cerebral injury in children. *Am J Roentgenol* 1990;154:361-8.
- 10 Saulsbury FT, Alford BA. Intracranial bleeding from child abuse. The value of skull radiographs. *Pediatr Radiol* 1982;12:175-8.
- 11 Sato Y, Yuh WT, Smith WL, Alexander RC, Kao SC, Ellerbroek CJ. Head injury in child abuse: evaluation with MR imaging. *Radiology* 1989;173:6537.
- 12 Jaspan T, Narborough G, Punt JAG, Lowe J. Cerebral contusional tears as a marker of child abuse—detection by cranial sonography. *Pediatr Radiol* 1992;22:237-45.
- 13 Sinai SH, Ball MR. Head trauma due to child abuse: serial computerised tomography in diagnosis and management. *South Med J* 1987;80:1505-12.
- 14 Dykes LJ. The whiplash shaken infant syndrome: what has been learned. *Child Abuse Negl* 1986;10:211-21.
- 15 Showers J. "Don't shake the baby": the effectiveness of a prevention program. *Child Abuse Negl* 1992;16:11-8.

New treatments for multiple sclerosis

May delay deterioration

Encouraging progress has been made in the development of treatments that reduce disease activity in multiple sclerosis. The relation between disability and such activity (as reflected in the relapse rate or the appearance of new lesions on magnetic resonance imaging) is complex, but recent evidence suggests that it may be possible to delay, though not yet abolish, the progress of irrecoverable neurological deficit.

The results of three large randomised controlled trials in ambulant patients with little or moderate disability have been announced during the past 18 months. Only one has been fully reported in the medical literature,¹ but two others were widely discussed at a joint meeting of the American Neurological Association and the Association of British

Neurologists held last October. Because much uncertainty exists about the potential role of these very expensive treatments it is important that the present position is clearly understood. In the review that follows, the figures for the two recent reports have been derived from information provided by the sponsors of the trials^{2,3}; the data on which the conclusions of these two studies are based have yet to be subjected to peer review.

In a trial of interferon beta-1b 372 patients with relapsing and remitting disease were allocated to placebo or a low dose (1.6 MIU) or high dose (8 MIU) of the drug, which was given by subcutaneous injection on alternate days. After two years the relapse rate (the primary end point) was significantly

lower in patients receiving the high dose than in those receiving placebo (1.27/year *v* 0.84/year, $P=0.0001$). This effect was maintained at three years.

A more striking effect was seen with the secondary end points of appearances on magnetic resonance imaging.⁴ After two years the total mean area of abnormality in annual T2 weighted scans had increased by 20% in the placebo group, compared with a reduction of 0.1% in the high dose group. In a subgroup of patients scanned at intervals of six weeks the median number of active lesions per year was 0.5 in patients treated with high dose interferon beta-1b compared with 3.0 in patients treated with placebo.

Changes were not, however, accompanied by any effect on disability as measured by the Kurtzke expanded disability status scale.¹ The natural variability of multiple sclerosis or the fact that the trial studied too few patients or was too short may explain the failure to detect genuine clinical benefit. Alternatively, interferon beta may have been modifying those acute elements of the pathological process that lead to relapse but not those that determine disability. These are poorly understood, and currently we do not know whether changes evident on standard magnetic resonance imaging predict clinical outcome.⁵ Hence the need for a trial with disability as a primary outcome measure.

The results of such a trial were reported last October.² Interferon beta-1a (which differs from interferon beta-1b in its mode of preparation and chemical structure) was given once a week by intramuscular injection. The trial was conducted on 301 patients with relapsing and remitting or relapsing and progressive multiple sclerosis who had little or no disability. They were randomly assigned to either treatment with 6 MIU interferon or placebo. The primary end point was the time to progress from baseline by one unit on the expanded disability status scale. Kaplan-Meier survival curve analysis showed a significant delay in deterioration in the treated group.

As with interferon beta-1b, the annual relapse rate was lower in the treated group (0.9/year *v* 0.62/year, $P=0.003$). Gadolinium-pentetic acid (DTPA) enhancement to assess pathological activity showed that treatment halved the mean number of active lesions in the treated group after two years. The change in the total mean area of abnormality in T2 weighted scans, however, did not differ between the placebo and treated groups, unlike in the trial of interferon beta-1b.

In both trials of interferon beta, some patients experienced minor side effects, including transient flu-like symptoms and minimal irritation at the injection sites. In neither trial were serious side effects encountered. A slight excess of episodes of depression with interferon beta-1b¹ was not found with interferon beta-1a.² No information is yet available about the risks of long term (longer than five years) continuous use of interferon beta in large numbers of patients.

Reduces relapses

Taken together, these two trials lead to the conclusion that two forms of interferon beta favourably modify at least some aspects of the pathological process in relapsing and remitting and relapsing and progressive multiple sclerosis and perhaps reduce the relapse rate by about a third. In addition, interferon beta-1a may have a favourable effect on the clinical course; large trials with disability as a primary end point are currently being conducted to assess whether interferon beta-1b also does. No information exists about the effect of interferon beta in patients with a progressive course from the onset of their disease (primary progressive multiple sclerosis), or with a progressive course after an initial relapsing and remitting phase (secondary progressive multiple sclerosis).

The results of a two year trial of copolymer 1 (a synthetic polypeptide with some resemblance to myelin basic protein) were also reported at October's meeting. A total of 251 patients with relapsing and remitting multiple sclerosis were randomised to either placebo or active treatment, which consisted of 20 mg copolymer 1 daily by subcutaneous injection for two years. The primary end point was the relapse rate, which was significantly lower in the treatment group than the placebo group (1.74 *v* 1.37, $P=0.007$). Progression of disability was a secondary outcome measure. The proportion of patients who progressed by ≥ 0.5 on the expanded disability status scale over two years was significantly lower in the treated group (32%) than the placebo group (46%). A change of 0.5 on the expanded disability status scale at the low levels of disability of patients recruited to this trial is of only marginal clinical importance. No serious side effects occurred. The data from magnetic resonance imaging are not yet available. Further details of the trial are required before the effectiveness of copolymer 1 can be assessed.

The results of these trials provide grounds for optimism about developing a treatment for multiple sclerosis that will ameliorate its course. Caution, however, is still necessary. The effects of interferon beta and copolymer 1 on more disabled patients is unknown, and trials are under way. The data on which the conclusions from the trials of interferon beta-1a and copolymer 1 are based have not yet been published in peer reviewed journals, and many data in both trials remain to be analysed. The size of the effect in those patients who seem to benefit from interferon beta and the types of patients who might benefit from this treatment remain to be determined. Without this information the widespread use of interferon beta cannot yet be recommended.

Several new therapeutic strategies have been suggested on the basis of better understanding of the pathogenesis of the multiple sclerosis lesion. They include agents that produce general⁶ or selective⁷ immunosuppression or that interfere with the action of inflammatory cytokines. Serial magnetic resonance imaging enhanced with gadolinium-DTPA provides a powerful means of screening for an effect of such agents in relatively few patients (about 60 divided between treated and placebo groups) over a matter of months.⁸ Results of such studies of antibodies to CD4 cells and the immunosuppressant mitozantrone will be available later this year. When positive effects are found definitive (phase 3) large scale studies with disability as the end point will still have to be mounted, but progress may now be faster than before.

W I McDONALD
Professor

National Hospital for Neurology and Neurosurgery,
London WC1N 3BG

The Institute of Neurology receives a grant from Schering AG, manufacturers of interferon, beta-1b, for participating in a trial of this drug.

- 1 IFNB Multiple Sclerosis Study Group. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. I. Clinical results of a multicenter, randomized, double-blind, placebo-controlled trial. *Neurology* 1993;43:655-61.
- 2 Biogen's interferon-beta-1a reduces progression of disability in multiple sclerosis. First European presentation of phase III trial results. *Clinical Courier* 1994;12:1-4.
- 3 Teva Pharmaceutical Industries. *Copolymer-1. The search for a specific therapy for MS*. TPI: 1994.
- 4 Paty DW, Li DKB. The UBC MS/MRI Study Group and the IFNB Multiple Sclerosis Study Group. Interferon-beta-1b is effective in relapsing-remitting multiple sclerosis. II. MRI analysis results of a multicenter, randomized, double-blind, placebo-controlled trial. *Neurology* 1993;43:662-7.
- 5 McDonald WI, Miller DH, Thompson AJ. Are magnetic resonance findings predictive of clinical outcome in therapeutic trials in multiple sclerosis? The dilemma of interferon- β . *Ann Neurol* 1994;36:14-8.
- 6 Sipe JC, Romine JS, Koziol JA, McMillan R, Zyffro J, Beutler E. Cladribine in treatment of chronic progressive multiple sclerosis. *Lancet* 1994;344:9-13.
- 7 Moreau T, Thorpe J, Miller D, Moseley I, Hale G, Waldmann H, *et al*. Preliminary evidence from magnetic resonance imaging for reduction in disease activity after lymphocyte depletion in multiple sclerosis. *Lancet* 1994;344:298-301.
- 8 Nauta JJP, Thompson AJ, Barkhof F, Miller DH. Multiple resonance imaging in monitoring the treatment of multiple sclerosis patients: statistical power of parallel-groups and crossover designs. *J Neurol Sci* 1994;122:6-14.