clinically relevant examples than the purely scientific questions posed by Montgomery and colleagues.

PETER MILLS
Consultant cardiologist

Royal London Hospital,
London E1 1BB

1 Montgomery H, Hunter S, Morris S, Naunton-Morgan R, Marshall RM. Interpretation of electrocardiograms by doctors. *BMJ* 1994;309:1551-2. (10 December.)

Formal teaching can reduce serious errors in interpretation

EDITOR,—We were not surprised by the results of Hugh Montgomery and colleagues' study of doctors' ability to interpret electrocardiograms.¹ We have audited the ability of senior house officers to interpret electrocardiograms in an accident and emergency department.² We found that only 114 of 245 traces were interpreted correctly but that management was seriously affected by misinterpretation in only eight cases. After formal teaching and the issue of guidelines serious errors of misinterpretation fell by half to four out of 242 (1.6%; confidence interval 0.7% to 4%).

We also tested the ability of cardiographers, cardiac technicians, nurses in the coronary care unit, anaesthetic registrars, and medical senior house officers to interpret electrocardiograms; we used a standard set of 20 electrocardiograms. All the subjects were asked to report the rate, rhythm, QRS axis, and abnormalities. Answers were graded 1-4, where 1=complete concordance; 2=minor disagreement (for example, first degree heart block); 3=disagreement (for example, left ventricular hypertrophy); and 4=appreciable disagreement potentially affecting management (for example, ST segment elevation). All the paramedical staff were better than the doctors and made fewer serious errors (table; $\chi^2=27.2$, $P<0.01$).

We believe that these results are explained by the fact that formal training in interpretation of electrocardiograms is given to all the groups of staff that we tested except the doctors. It is an indictment of medical education that doctors can interpret electrocardiograms less well than paramedical staff. Should this interpretation be part of the undergraduate curriculum? We believe that it should certainly be part of postgraduate training for doctors in clinical specialties. It is particularly important when decisions about giving thrombolytic treatment to patients with acute myocardial infarction depend partly on interpretation of the electrocardiogram. If general practitioners begin to use thrombolysis in the community they must be capable of recording and interpreting an electrocardiogram.

P A WOODMANSEY
Research registrar
T WHITE
Medical student
K S CHANNER
Consultant cardiologist

1 Montgomery H, Hunter S, Morris S, Naunton-Morgan R, Marshall RM. Interpretation of electrocardiograms by doctors. *BMJ* 1994;309:1551-2. (10 December.)

2 White T, Woodmansey PA, Ferguson DG, Channer KS. Improving the interpretation of electrocardiographs in an accident and emergency department. *Postgrad Med J* (in press).

Grade given to interpretation by different groups of staff of sample of 20 electrocardiograms (ECGs)*

Grade*	Registrar (n=6)	Nurse in coronary care unit (n=6)	Cardiac technician (n=6)	Cardiographer (n=5)	Senior house officer (n=6)
1	53	46	51	51	47
2	13	19	8	6	10
3	18	27	43	26	33
4	36	21	23	17	30
Total ECGs	120	120	120	100	120

*See text for details of grades.

Continuous ambulatory electrocardiography in elderly people

EDITOR,—I J Räihä and colleagues' paper on the predictive value of continuous ambulatory electrocardiographic monitoring in elderly people digresses and therefore arrives at several misleading and clinically unsound recommendations.¹ In a retrospective study the authors found several abnormalities on Holter monitoring that were associated with a high odds ratio for the risk of cardiac death over 10 years. The subjects had a mean age of about 75.

The predictive value—as distinct from the odds ratio—is a statistical term indicating the likelihood that a patient in whom the result of a test is abnormal will have the disease in question (or, in this case, a poor outcome). With respect to the three factors found by Räihä and colleagues to be associated with a poor outcome, 95 patients had >100 ventricular ectopic beats during the day, of whom 32 died of cardiac causes over the next 10 years; 88 had multifocal ventricular ectopic beats at night, of whom 31 died of cardiac causes; and 79 had sinus pauses of >1.5 s during the day, of whom 34 died of cardiac causes. Thus the predictive value that a given patient with one of the identified factors on Holter monitoring would die of heart disease over the ensuing 10 years was only slightly greater than 1 in 3.

Even if we assume that these markers, which were defined arbitrarily and post hoc, are indeed valid, what possible benefit could there be in determining that a given 75 year old has a 1 in 3 chance of dying of heart disease during the next 10 years? To imply that Holter monitoring should be performed routinely to stratify the risk in elderly people and to recommend that "patients with these findings should be evaluated for silent ischaemic heart disease and sick sinus syndrome" seems the height of folly. The cardiological community generally accepts that giving antiarrhythmic drugs to patients with ventricular extrasystoles does more harm than good, and Räihä and colleagues themselves state that "the benefit of treating silent ischaemia in elderly patients is unclear." Adherence to the authors' clinical recommendations would not only be economically wasteful but also give rise to much unnecessary anxiety in a population already so concerned by its declining sense of wellbeing.

MAYER BASSAN
Consultant cardiologist

Jerusalem Heart Clinic of Kupat Holim,
Jerusalem,
Israel

1 Räihä IJ, Piha SJ, Puukka P, Sourander LB. Predictive value of continuous ambulatory electrocardiographic monitoring in elderly people. *BMJ* 1994;309:1263-7. (12 November.)

Weight in infancy and coronary heart disease in adult life

EDITOR,—C H D Fall and colleagues' paper assumes that low weight at 1 year is a result of undernutrition and that greater weight is a result of better nutrition.¹ Unfortunately, weight gain in the first year of life is not as straightforward as this.

When weight profiles of individual babies are examined in the early months considerable shifts across centile lines take place in both upward and downward directions as the baby leaves behind intrauterine influences on growth and grows under the direction of his or her own genetic machinery.² In a study of weight gain over the first year in 66 normal term infants only 12% kept roughly in the same "centile channel" in which they were born; 28% showed downward crossing of centiles, while most showed upward crossing of centiles.³ The greatest shift occurred in the first three months.

Some babies of low weight at 1 year will therefore be those who were light at birth, for physiological or pathological reasons, and who maintain this phenotype. Some will have become small, having crossed various numbers of centile channels. These have not all necessarily failed to thrive but have simply recaptured their true growth phenotype. At the same time, there are those of higher weight who have continued a high weight profile at birth while others have reached this weight, having caught up from being relatively light at birth. It is therefore important to know more about the detailed patterns of growth in the babies in the authors' study who weighed ≤ 18 lb (≤ 8.2 kg) and ≥ 26 lb (≥ 11.8 kg). Is metabolic disadvantage relevant to those whose weight continues in a low channel or in those whose weight has fallen from a higher centile and, if so, which type? Likewise, what of the protection conferred on those who are of higher weight at 1 year?

There is further confusion. In some of the earlier studies by the authors low birth weight was linked with later risks of high blood pressure and ischaemic heart disease.⁴ Yet a typical pattern of early weight gain in growth retarded neonates is to catch up—to reach a higher centile than they were at birth.² This phenomenon is also found in some infants of appropriate weight for gestation. Some heavier children at 1 year will therefore probably be those who have caught up from intrauterine growth retardation. On the one hand, these are considered to be at risk of later cardiovascular disease, yet we now hear that they seem to be advantaged in being less vulnerable to later coronary artery disease.

It is an intriguing possibility that early growth may somehow be related to programming of later disease. But the literature on the subject contains contradictions that need to be explained.

D P DAVIES
Professor of child health

University Department of Child Health,
University of Wales College of Medicine,
Cardiff CF4 4XN

1 Fall CHD, Vijayakumar M, Barker DJP, Osmond C, Duggleby S. Weight in infancy and prevalence of coronary heart disease in adult life. *BMJ* 1995;310:17-9. (7 January.)

2 Davies DP, Williams T. Is weaning babies in clinics worthwhile? *BMJ* 1983;286:860-7.

3 Davies DP, Martan R. Growth and growth charts. *BMJ* 1980;280:1188.

4 Barker DJP, Osmond C, Golding J, Kuh D, Wadsworth MEJ. Growth in utero, blood pressure in childhood and adult life and mortality from cardiovascular disease. *BMJ* 1989;298:564-7.

Simple aspiration for spontaneous pneumothorax

EDITOR,—Editor's Choice in the issue of 19 November describes the clinical trial reported by John Harvey and Robin J Prescott—of simple aspiration versus intercostal drainage for spontaneous pneumothorax in patients with normal lungs¹—as "methodologically strong." I challenge that view.

Firstly, patients "with signs of a tension pneumothorax" are said to have been excluded from the trial, presumably because intercostal tube drainage is considered to be mandatory in all such cases. But the criteria for that diagnosis are not stated, and at

least some of the 28 patients with "complete collapse" of the lung might well have been shown to have a tension pneumothorax if intrapleural pressures had been measured. Unrecognised tension pneumothorax might therefore have been responsible for some of the seven failures of aspiration, and that figure could have been even higher if most patients (18/28) with complete lung collapse had not been treated by intercostal drainage.

Secondly, although the authors state that "28 out of 35 aspirations were successful," the corresponding figures for intercostal drainage are not provided. Why not?

Thirdly, it is methodologically unsound, and perhaps ethically questionable, to include in a clinical trial of this type patients with vastly different indications for therapeutic intervention. Those with only a "small rim" pneumothorax should not have been exposed to the chance of being submitted to the unnecessary trauma of intercostal drainage (as one of them was) when they probably required no active treatment at all. Conversely, it could have been dangerous not to resort immediately to intercostal drainage in patients with complete lung collapse when the aspiration of 3 litres of air without x ray evidence of re-expansion of the lung strongly suggested the persistence of a fistulous communication between the lung and pleura. The doctors concerned should also have considered the remote but lethal possibility of a pneumothorax developing on the opposite side.

Fourthly, two of the benefits attributed to aspiration—less pain than with intercostal drainage and a shorter stay in hospital—were surely too obvious to warrant statistical analysis. The observation that intercostal drainage increased the need for subsequent pleurectomy (now an obsolete procedure) would surprise most thoracic physicians and surgeons but could perhaps be explained by the higher frequency of complete lung collapse (18/38 *v* 10/35) in the intercostal drainage group.

The authors are right to advocate aspiration in suitable cases as a less uncomfortable alternative to intercostal drainage. They have not, however, provided convincing evidence that the results of a statistical comparison of two ill matched groups should usurp individual clinical decisions on whether the initial treatment of spontaneous pneumothorax should be aspiration or intercostal drainage.

IAN W B GRANT
Former consultant physician

Nether Balchandy,
By Pitlochry,
Perthshire PH16 5JT

1 Harvey J, Prescott RJ. Simple aspiration versus intercostal tube drainage for spontaneous pneumothorax in patients with healthy lungs. *BMJ* 1994;309:1338-9. (19 November.)

Author's reply

EDITOR,—When the British Thoracic Society's research committee began the study in 1985 some doctors declined to take part because they thought that all pneumothoraces should be treated by simple aspiration. An equal number declined to take part because they thought that all pneumothoraces required intercostal tube drainage. It was because of these uncertainties that a well designed, randomly selected study was undertaken. Rather than "reinventing the wheel," we were testing which of the two commonly used wheels was the right one to use.

Since our aim was to test current clinical practice, patients with signs of tension pneumothorax were specifically excluded from the trial. Most doctors do not now measure intrapleural pressures and would not diagnose tension pneumothorax simply on the basis of complete collapse. Whatever the numbers might indicate, the distribution of patients between the two randomly

selected arms showed no greater variation than would be expected by chance, and the interpretation of our findings is not affected by the greater number of patients with complete collapse in the intercostal drainage group. Aspiration was successful in six of 10 patients with complete collapse, so that even in this group an initial attempt at aspiration would not be contraindicated. We do not agree, therefore, that routine thorascopic surgery is indicated as the initial treatment for all patients with spontaneous pneumothorax who present with complete collapse, as Anthony P C Yim proposes, although it may in future be more commonly used for recurrent episodes.

Whether intercostal drainage is successful depends on how success is defined. Two of the 38 patients treated by intercostal drainage required pleurectomy during the same admission and a further five within the next year because of recurrent pneumothorax, whereas none of the patients in whom aspiration either failed or was successful required pleurectomy during the subsequent year. The size of pneumothorax is not, therefore, an "alternative explanation for the higher rate of pleurectomy in the group treated by intercostal drainage."

We accept that minimal intervention thorascopic surgery is now rapidly replacing pleurectomy, but it was not in widespread use during the years of this study. We also agree that "small rim" pneumothoraces would not usually require any drainage procedure. Only four of the 73 patients in our study had small rim pneumothoraces, and they were judged by the doctors to merit a drainage procedure.

As we mentioned, the results of this study have been widely discussed with respiratory physicians and thoracic surgeons and form the basis of our guidelines.³ Given the successful outcome measures, a less painful and less traumatic procedure that requires a shorter stay in hospital seems to be the best initial management for all spontaneous pneumothoraces in patients with normal lungs.

JOHN HARVEY
Consultant physician

Lung Function Unit,
Southmead Hospital,
Bristol BS10 5NB

ROBIN PRESCOTT
Senior lecturer in medical statistics

Department of Public Health Sciences,
Edinburgh University Medical School,
Edinburgh EH8 9AG

1 Bradley G. Simple aspiration for spontaneous pneumothorax. *BMJ* 1994;310:255-6. (28 January.)

2 Currie D. Simple aspiration for spontaneous pneumothorax. *BMJ* 1994;310:256. (28 January.)

3 Miller AC, Harvey J. Guidelines for the management of spontaneous pneumothorax. *BMJ* 1993;307:114-6.

Reform of GMSC not needed

EDITOR,—I believe that David Roberts is wrong in his assertion that morale in general practice is related to the method of election of the General Medical Services Committee and in his conclusion that the committee is undemocratic.¹ I have been a member of the GMSC since 1980 and have served on the agenda committee for the annual representative meeting for several years. I carried out the detailed work on the section of the agenda that included the motion about the method of election to the GMSC. My original decision was adopted after debate by the agenda committee, and my view of the motion's importance was upheld by the annual representative meeting, which decided not to suspend standing orders to debate it.

There were two main reasons for the decision not to raise the issue at the meeting. The first concerns the internal working of the constitution of the BMA² and the necessity of maintaining a united front to represent all doctors. This requires

that the GMSC is allowed to operate in a largely autonomous way and also that amendment of its constitution and electoral arrangements is carried out in a way acceptable to the conference of local medical committees. The second and much more important reason relates to democracy and the representation of ordinary general practitioners. Roberts fails to appreciate that the sovereign body in general practice affairs is the local medical committee, not the GMSC. All general practitioners have ample opportunity to seek to influence the decisions of their local medical committee; they have the opportunity to seek election to it and from there to seek election to the GMSC.

The role of the GMSC is to carry out the collective will of local medical committees. These committees raise the quota that maintains the General Medical Services Defence Fund Limited, which gives freedom of action to the GMSC while maintaining the local medical committees' control over it. As the NHS becomes more fragmented and local deals attain increasing importance, the prime need for every general practitioner is an effective local medical committee. I believe it is essential that the role of local medical committees is not sidelined by the adoption of a spurious argument about democracy.

Finally I believe that my office of secretary of a local medical committee requires me to vote against an extraordinary general meeting or referendum if Roberts puts such a resolution to the BMA's council.

J DAVID WATTS
Secretary

Ayrshire and Arran Local Medical Committee,
Dundonald,
Ayrshire KA2 9HE

1 Roberts D. Reform of GMSC needed. *BMJ* 1994;309:1585. (10 December.)

2 Watts JD. Strains in the constitutional framework of the BMA? *BMJ* 1988;296:1549.

Doctors and management

EDITOR,—In describing how managers and doctors view each other Jenny Simpson makes no reference to the inherent suspicion among doctors that managers are not totally committed to health care.¹ Whatever changes occur in the health service, doctors will still be there caring for their patients in what they believe to be the best way; most doctors would never contemplate leaving the NHS, no matter how dire the situation. Managers, however, have a skill that is in demand throughout the economy. A better offer of remuneration could mean that they find themselves in the upper echelons of a trust one week and in the offices of a manufacturer of machine tools or a telecommunications company the next. Many health service managers are dedicated to the NHS, but they must communicate this to doctors in word and deed if they are to dispel this commonly held prejudice.

BRENDAN HARRINGTON
Lecturer in child health

St Mary's Hospital,
Manchester M13 0JH

1 Simpson J. Doctors and management—why bother? *BMJ* 1994;309:1505-8. (3 December.)

Failure of consultant expansion

Consultant expansion depends on local cooperation

EDITOR,—Surely Stephen Brearley's editorial on the failure of consultant expansion underestimates the importance of teamwork in producing changes in major issues at the local level.¹ If people wish to move towards a consultant led service then the consultants in individual hospitals will need to work closely with their purchasing authorities;