Nursing is underrepresented

EDITOR,—Fiona Godlee's controversial articles on the World Health Organisation expose its strengths and weaknesses but make scant mention of the contribution of nursing to its operations and organisation.¹ This is particularly regrettable given the articles' attention to the medical dominance of the organisation. We can conclude only that the absence of any substantive discussion of nursing reflects the perceived invisibility of nursing within the organisation more generally.

A recent study of nurses in government and non-government organisations across the world chronicled the fluctuating fortunes of the nursing policy base at the WHO since its inception in 1948 and the consequences of nursing's diminished voice in international matters of health.² The contemporary history of the WHO's operations as represented by Godlee suggests the retreat of professionals into enclaves of skill, bureaucratic shelters from the sully of the real world. Yet much of the success of the WHO's programmes has been due to its traditional reliance on nursing and midwifery skill and staff for their implementation in the field. Credit for the successes of programmes such as the expanded immunisation, family health, emergency relief, AIDS, and mental health programmes do not reflect the major contributions of nurses and midwives. Indeed, it is almost a condition that in joining such programmes nurses are expected to shed any overt allegiance to nursing.

How many initiatives of the WHO have failed because they have taken insufficient account of the conditions under which nursing and other front line workers can contribute effectively to health care in the field? How much more could be achieved with appropriate nursing leadership than is at present?

It is worth speculating whether the fate of nursing reflects the declining faith in health professionals as advocates and activists of international health more generally and, in the light of the World Bank's recent initiatives in the health field, a swing of the pendulum towards favouring economists as the technical "fix its" of the future.⁴ Rather than mirroring the status patterns and values of Western capitalism more generally, the WHO could set a moral and economic example by putting its own hierarchical house in order. By taking such positive action it would ensure the greater representation of nursing skills throughout the organisation and the more cost effective deployment of skilled staff.

JANE ROBINSON
Professor

Department of Nursing and Midwifery Studies,
Medical School,
University of Nottingham,
Nottingham NG7 2UH

ANNE MARIE RAFFERTY
Harkness fellow

Centre for Health Services Research and Policy Analysis,
University of Pennsylvania,
PA, USA

1 Godlee F. WHO in crisis. *BMJ* 1994;309:1424-8. (26 November.)

2 Splane R, Splane V. *Chief nursing officer positions in national ministries of health: focal points for nursing leadership*. San Francisco: University of California San Francisco Press, 1994.

3 World Health Organisation Study Group. *Nursing beyond the year 2000*. WHO Tech Rep Ser 1994; No 842.

4 World Bank. *World development report 1993: investing in health*. Oxford: Oxford University Press, 1993.

A new strategy is needed

EDITOR,—Fiona Godlee is right: the World Health Organisation is in crisis.¹ In the past it came to occupy a position of leadership in major global undertakings such as the eradication of smallpox, the work resulting in the code on breast milk substitutes, the programme on essential drugs, and the primary health care movement. It grasped the importance of these issues, brought its own

resources to bear, and harnessed resources from the outside. It became the instrument by which the global community could agree on common goals and achieve collective action.

But today's challenges are not being met in a similar manner. Health services are breaking down in many countries as a result of insufficient resources. Widespread privatisation programmes are jeopardising access to health care for poor people in developing and industrialised countries alike. The AIDS pandemic threatens to wipe out the progress with regard to health in many parts of the world. The World Bank, in particular through its report *Investing in Health*,² is increasingly taking a leading role in health. Meanwhile, the WHO is occupied by trying to recycle its "health for all" strategy.

The WHO must recapture its normative functions, including not only the obvious ones such as the setting of standards and classification of diseases but also the dissemination of information, advocacy for health, promotion of research, and, above all, its role as the global leader in health and health policy.^{3,4} But it needs also to redefine its role. The most important policy issue that faces it is the very notion of health. In its constitution "health" is the main concept. Yet "disease" has been the focus of action for all the 47 years of the WHO's existence. Medical knowledge has developed, and diagnostic and therapeutic tools have been refined to new technological heights. The WHO has taken as its role the application of this knowledge to specific disease problems.

The time has come to make health and health development the main focus of interest. That would entail focusing more on the aetiology of health and the factors in our environment that are necessary for a healthy life than on pathogenic vectors. It would also mean an end to the medical monopoly on health. Other professionals, such as economists, anthropologists, sociologists, and health planners, would come into the forefront to join with the medical profession in the battle for health.

Doing the right things has to be accompanied by measures to improve internal efficiency. The most urgent reforms in this regard are the strengthening of health policy and analytical functions at country level and the abolition of an obsolete regional structure.

The member states of the WHO must now take the lead. It is up to them to define and implement a new strategy in the battle for world health. If they succeed it will also mean the survival of the WHO.

BO STENSON
Senior fellow

Karolinska Institutet,
Department of International Health and Social Medicine,
S-171 77 Stockholm,
Sweden

1 Godlee F. WHO in crisis. *BMJ* 1994;309:1424-8. (26 November.)

2 World Bank. *World development report 1993: investing in health*. Oxford: Oxford University Press, 1993.

3 Nordic UN Project. *The United Nations: issues and options*. Stockholm: Almqvist och Wiksell, 1991.

4 Stenson B, Sterky G. What future WHO? *Health Policy* 1994;28: 235-56.

August: the killing season

Last week in July is a poor comparison

EDITOR,—Paul Aylin and F Azeem Majeed use an incorrect comparison to prove the hypothesis that the newly qualified house officers who start their posts in the first week of August may be associated with an increase in deaths in hospital—the "killing season."¹ They compare details of all deaths occurring in the first week of August with those occurring in the last week of July; finding no significant difference, they conclude that newly qualified house officers have been falsely accused of increasing the number of deaths in hospital.

Junior medical staff often take annual leave during the last week of July (despite attempts to prevent this), and their places are either taken by locum doctors or remain vacant. It would not be surprising to find that this week was a time of increased mortality, and therefore it is no comfort to know that the first week in August is no different. Perhaps a better comparison would have been a week in the middle of the six months or, even better, the mortality averaged over the whole six months.

PAUL SIKLOS
Consultant physician

West Suffolk Hospital,
Bury St Edmunds,
Suffolk IP33 2QZ

1 Aylin P, Majeed FA. The killing season—fact or fiction? *BMJ* 1994;309:1690. (24-31 December.)

Other measures of quality are needed

EDITOR,—Paul Aylin and F Azeem Majeed point out that they did not have data on admission rates or bed occupancy during the weeks of their study into whether the changeover of house officers in hospitals in August results in an increase in potentially avoidable deaths.¹ They were therefore unable to calculate death rates. Death rates are a crude measure of suboptimal care, and measures of quality such as adverse events or complication rates should also be looked at.² We suggest that the quality of care should be audited regularly throughout the year, so that the adequacy or otherwise of care during the changeover in August is fully assessed.

JOHN LEWIS
Senior registrar
PETER JENKINS
Consultant psychiatrist
MARION GRAY
Senior house officer
CAROL BANNISTER
Research registrar

South Gwent Consultation Liaison Psychiatry Team,
St Cadoc's Hospital,
Caerleon,
Gwent NP6 1XQ

1 Aylin P, Majeed FA. The killing season—fact or fiction? *BMJ* 1994;309:1690. (24-31 December.)

2 Smith R, ed. *Audit in action*. London: BMJ Publishing Group, 1992.

Effects of health publicity on prevalence of smoking

EDITOR,—Donald Reid draws attention to the potentially class free effects of health promotion in the electronic media, but he misrepresents the term "health publicity" in our analysis by referring to it as "everything else except price."¹ The model we used to assess different influences on cigarette smoking over the past 20 years included price, income, a specific linear trend, and an error term for random variation.²

Smoking has not declined steadily over this period but, for most of the groups we considered, has oscillated about a downward trend. This cumulative linear effect, not explained by price or income, we interpreted as including the net effects of health publicity and other social trends, including social acceptability and restrictions in the workplace and public places; we denoted this by the term "health publicity." The term would include, but not differentiate, effects of health publicity against smoking in the electronic media. It represents a constant annual decrease in smoking for each group, while the error term includes variation not explained by the three other variables. We conclude that in Britain in the 20 years to 1990 there was no such decreasing trend in cigarette smoking by women in socioeconomic groups IV