Rotterdam symptom checklist, the instrumental help for daily living scale, the activities of daily living scale, and the social experiences scale. Satisfaction was assessed with the Rand corporation's patient satisfaction questionnaire.

Our trial failed to attain the intended sample size too and we stopped the intake of patients. McWhinney and colleagues reported problems with patient recruitment, attrition, and the exposure of the control group to services equivalent to the home support team. In our case the major problem was patient accrual. As it happened, the number of eligible patients was nearly half that of previous years. To improve patient intake we invited on several occasions 11 associated centres to refer their patients. During two years only 22 patients entered the trial, of whom nine came from associated centres. Furthermore, 22 eligible patients refused to participate in the trial. This was suspected early on in the trial, and the reason for refusal and the characteristics of these patients were registered. Eight patients did not want chemotherapy, 10 refused either continuous infusion or randomisation, and four were not asked by the physician because of medical reasons. In contrast with McWhinney and colleagues' experiences, attrition did not have a major role—four patients died before the first evaluation at two months, and all other patients completed the questionnaires. A possible explanation for this difference could be that the patients in McWhinney and colleagues' trial had a more advanced disease than our patients.

Despite inclusion criteria allowing only patients with disseminated colorectal cancer, without metastasis to the brain, and with normal results of haematology tests, with normal liver and kidney function, World Health Organisation performance status of 2 or more, and no serious comorbidity to participate, our study population turned out to be rather heterogeneous. Intersubject variability of the measures of quality of life was great. Individual plots of quality of life scores over time seemed stable when the clinical situation was stable and seemed to decline around the time of disease progression. No apparent relations between quality of life and satisfaction were found. Small sample size and great intersubject variability do not allow for further statistical analysis.

In conclusion, randomised trials in palliative cancer care are difficult, but not impractical. Our trial failed because of difficulties in patient accrual, but other common problems, such as attrition, contamination of exposure, or selection of adequate outcome measures did not have a major role. Future randomised trials in palliative cancer care should consider multicentre participation and the possibility of patient entry without limitations to type of primary cancer. Moreover, the care intervention should be well defined, and adequate outcome variables should be selected.

GERDY RINCK
Junior clinical epidemiologist
JOS KLEIJNEN
Clinical epidemiologist

Department of Clinical Epidemiology and Biostatistics,
Academic Medical Centre,
University Hospital of Amsterdam,
PO Box 22700,
1100 DE Amsterdam,
Netherlands

TRUDI G A M VAN DEN BOS
Professor of social medicine and chronic diseases
Institute of Social Medicine,
Academic Medical Centre

HANNEKE J C J M DE HAES
Associate professor of medical psychology
Department of Medical Psychology,
Academic Medical Centre

EGBERT SCHADE
Professor of family medicine
Department of General/Family Medicine,
Academic Medical Centre

CEES H N VEENHOF
Associate professor of medical oncology
Division of Medical Oncology,
Department of Internal Medicine,
Academic Medical Centre

- 1 McWhinney IR, Bass MJ, Donner A. Evaluation of a palliative care service: problems and pitfalls. *BMJ* 1994;309:1340-2. (19 November.)
- 2 Ovesen L, Allingstrup L, Hannibal H, Mortensen EL, Hansen OP. Effect of dietary counseling on food intake, body weight, response rate, survival and quality of life in cancer patients undergoing chemotherapy: prospective randomized study. *J Clin Oncol* 1993;11:2043-9.

- 1 Dillner L. Scottish death rates published with health warning. *BMJ* 1994;309:1599-600. (17 December.)
- 2 Goldacre MJ. Hospital inpatient statistics: some aspects of interpretation. *Community Medicine* 1981;3:60-8.
- 3 Goldacre M, Seagroatt V, Hawton K. Suicide after discharge from psychiatric inpatient care. *Lancet* 1993;342:283-6.
- 4 Seagroatt V, Goldacre MJ. Measures of early post-operative mortality: beyond hospital fatality rates. *BMJ* 1994;309:361-5.
- 5 Gill L, Goldacre MJ, Simmons H, Bettley G. Computerised linking of medical records: methodological guidelines. *J Epidemiol Community Health* 1993;47:316-9.

Interpreting hospital death rates

Can be difficult

EDITOR,—We agree that hospital death rates for myocardial infarction can be difficult to interpret.¹ In a guide to hospital statistics and their interpretation we reported hospital death rates in Oxford, which ranged from 8% (27/341) to 73% (73/100).² The hospitals were anonymised because our code of practice is not to identify individual hospitals in publications.² The hospital with a death rate of 8% admitted only convalescent medical patients: its statistics were selectively based on survivors. The hospital with a death rate of 73% was a geriatric hospital: many of its patients were admitted, with conditions other than myocardial infarction, for long stay care. Those whose deaths were certified as due to myocardial infarction contributed not only to the numerator but also to the denominator because they had a fatal infarct while in hospital.

Circumstances such as these should be easy to identify locally, but a more important problem is that of defining a hospital admission when patients with myocardial infarction die close to the time of admission. There may be considerable variation among hospitals in whether such patients have been formally admitted and therefore included in the hospital's statistics.² We also know currently of a hospital computer system in which, when the patient's disposal is unknown to the coder, the default option is that the patient is coded as alive. A useful check in studies of hospital death rates is to identify death certificates that specify that the patient died in hospital and see whether a corresponding hospital record abstract exists, and vice versa. Although problems like these arise, however, they should be addressed in the analysis and interpretation of hospital statistics rather than be regarded as reasons for not auditing hospital care.

England's problem has been not so much the lack of means to link records but the incompleteness of its data since the Körner changes. For example, in the Oxford region from 1979 to 1986 (when collection of data for hospital statistics was managed by the Oxford record linkage study) 97% of hospital activity analysis records (1 557 753/1 609 588) included diagnostic data. From 1987 to 1990 the figure dropped to 79% (1 095 711/1 383 973), and it has only gradually crept back towards earlier levels of completeness. The completeness of clinical data in the national hospital dataset in England fell from the mid-1980s, and the hospital inpatient enquiry and the mental health enquiry—previously rich sources of information—have not been available since 1985.

Mortality after care in hospital can be studied in England.^{3,4} Statistical information about hospital episodes, including data needed to link records, exists in patient administration systems in most hospitals. Death certificates can be made available locally for linkage: copies are held by district departments of public health. Software to match records for linkage has been developed to run on microcomputers, and copies of it can be made available to anyone in the NHS in England.⁵

MICHAEL GOLDACRE
Honorary director
LEICESTER GILL
Chief computer scientist

Oxford Record Linkage Study,
Department of Public Health and Primary Care,
University of Oxford,
Anglia and Oxford Regional Health Authority,
Oxford OX3 7LF

Report was intended to promote discussion

EDITOR,—The stated intention of the Clinical Resources and Audit Group in publishing its report on clinical outcomes is to promote discussion among clinicians to determine what lessons can be learnt to improve the treatment of patients—a worthy and proper aim.¹ Contrary to the group's advice, however, the data provided have been put into league tables. This has caused considerable and quite unnecessary anxiety to people living in Fife. Had the opportunity been afforded to trusts to comment on a draft of the document before publication this anxiety would have been avoided by the correction of a simple error—that is, the inclusion of elderly patients in the figures for myocardial infarction. Furthermore, the appropriate and more satisfactory figure would have been attributed to the appropriate trust—namely, the Queen Margaret Hospital Acute NHS Trust.

When it was first suggested that these tables should be published medical directors in Scotland expressed considerable disquiet, being concerned that wrong messages would be given to the public and that the primary purpose, of promoting open discussion among clinical staff, would be undermined. This view is now well justified. The expectation is that purchasers will use the information in the report to influence the contracting process. Therefore if the public is not to be caused unnecessary anxiety, and if the document is to have credibility for clinicians and in the contracting process, basic errors must be eliminated before any further publication.

N W HARRY
Medical director

Fife Healthcare NHS Trust,
Leven,
Fife KY8 5RG

- 1 Dillner L. Scottish death rates published with health warning. *BMJ* 1994;309:1599-600. (17 December.)

Asthma trends

EDITOR,—Helen Booth and colleagues suggest that differences in the output of DeVilbiss No 40 and No 45 nebulisers¹ may have influenced our estimates of the increase in the prevalence of airway hyperresponsiveness in children studied 10 years apart, in 1982 and 1992.² We expressed airway responsiveness both as a dose-response ratio (defined as the percentage fall in forced expiratory volume in one second divided by the dose administered) and as airway hyperresponsiveness (defined as a 20% fall in forced expiratory volume in one second at a dose of 3.92 μmol histamine). As reported, we normalised the dose-response ratios measured in 1992 to take account of methodological differences in collecting the data in 1982 and 1992 and so that we could more accurately compare the dose-response ratios measured in 1982 with those measured in 1992. To normalise the 1982 values we added a further constant to the dose-response ratio so that the cumulative frequency response curve for normal subjects studied in 1992 matched that for normal subjects studied in 1982. This adjusted for systematic methodological changes between the years, including nebuliser outputs, brand of spirometer,