Local research ethics committees and multicentre drug trials

EDITOR,—The role of local research ethics committees in multicentre trials has been discussed at length in the *BMJ* and other journals. I understand the frustration of researchers at having to jump through so many hoops, but I sincerely hope that any central committees which do emerge reflect the practice of the best of the local committees and do not become a soft option for gaining approval for poor quality projects, either in hospital or general practice.

The ethics committee that I chair recently received a response from a major pharmaceutical company that shows scant regard for the carefully considered opinion of a local research ethics committee and a gung-ho attitude to ethics committee approval in general. The study in question was to compare antibiotics in the treatment of otitis media and chest infections in children. We coopted the expert opinion of a consultant microbiologist and on his advice wrote to the participating general practitioner, the following letter.

Dear Dr —

Re: A multicentre, open label, parallel group study in general practice of the comparative efficacy and tolerance of Orelox (cefepodoxime proxetil), Augmentin (co-amoxiclav), or Distacolor (cefaclor) in the treatment of otitis media and chest infections in children.

The ethics committee has carefully considered your protocol but was unable to approve it for the following reasons:

- (1) The inclusion criteria are extremely vague and most of the infections treated would undoubtedly be viral in origin.
- (2) Neither of the comparative drugs reflects the current recommendation for treatment of such infections in general practice in Leeds—that is, ampicillin.
- (3) Any data on bacteriological efficacy are likely to be very limited due to a combination of poor selection (see above) and poor techniques for microbiological sampling, with prolonged transport time to the central laboratory. Cough swabs are an extremely inefficient method of sampling lower respiratory tract secretions and an ear swab fails to sample the site of infection in otitis media in the absence of a perforated ear drum.
- (4) The list of causative pathogens in the protocol includes commensal organisms such as corynebacteria and viridans streptococci which are probably not pathogens in this setting.

The study seems to be of limited scientific value in an area where a good range of therapeutic options already exists. The vague criteria for patient inclusion are particularly worrying in view of the well recognised tendency to overprescribe antibiotics to children with these types of infection.

Yours sincerely

We would have been quite prepared to consider the drug company's response to these criticisms, which we believe are valid, and we would have been prepared to change our minds concerning approval if the problems could have been resolved to our satisfaction. However, instead of a substantive reply we received the following.

Dear Dr —

Re: A multicentre, open label...

Thank you for your letter concerning your ethics committee's consideration of our protocol.

I note your committee's comments on our protocol with considerable interest. Fortunately, there are

Advice to authors

We prefer short letters that relate to a recently published article and we are unlikely to publish letters longer than 400 words and containing over five references. Letters may be shortened. Your letters should be typed with double spacing and include a word count. All authors need to sign the letter and provide one current appointment and address. We encourage you to declare any conflict of interest. Please enclose a stamped addressed envelope if you require an acknowledgment.

sufficient LRECs [local research ethics committees] elsewhere in the country for us to be able to proceed with this study without the participation of Dr — as a trialist.

I understand that the Department of Health is currently engaged in drawing up new guidelines for local ethics committees in relation to their role in the consideration of multicentre studies. When this happens, we will hopefully see less of the failures of communication that have occurred during your consideration of this protocol.

I regret that this protocol submission to your committee has, in effect, been a waste of time. However, I would like to thank you for taking the trouble to give us your comments and look forward to doing business with you again in the future.

Yours sincerely

The only concern of the company seemed to be that its project would be allowed to take place in some centres. It appeared to show no interest in the valid questions that we posed, and we got the impression from the terms of its letter that it hoped that once there was a central committee the kind of objections raised by us could be put aside.

R P F DEAR
Chairman, clinical research
(ethics) committee

St James's University Hospital,
Leeds LS9 7TF

Emergency care in general practice

Many general practitioners are unable to provide safe emergency care

EDITOR,—Anthony Avery and Mike Pringle's editorial on emergency care in general practice is a counsel of perfection divorced from reality.¹ The reality is that, no matter how well they are trained and equipped, many general practitioners are not able to deliver a safe level of emergency care to their patients. This is particularly true in inner city areas, where the pressure of work in the surgery often precludes routine visiting, let alone the possibility of abandoning overcrowded waiting rooms to attend emergencies. Heavy traffic and one way systems often compel general practitioners to perform house visits on foot. The threat of violence in many council estates in inner cities has led doctors to abandon their black bags, quite apart from defibrillators and electrocardiographs, to avoid theft. There is an urgent need for a nationwide reactive and effective emergency service, staffed by properly trained paramedics, to

respond to medical emergencies at home while general practitioners get on with the work in which they are expert.

ANDREW DICKER
General practitioner

London SW1V 4PY

¹ Avery A, Pringle M. Emergency care in general practice. *BMJ* 1995;310:6. (7 January.)

British Heart Foundation recommends early defibrillation

EDITOR,—Anthony Avery and Mike Pringle emphasise the need for general practitioners to be adequately equipped and trained to provide optimal care in medical emergencies.¹ They emphasise the value of early defibrillation in the management of cardiac arrest, which in many cases could be provided by an appropriately equipped general practitioner. In recent years the British Heart Foundation has helped provide general practitioners with defibrillators, and the results show the value of this policy.

The foundation's records show that among 83 patients who suffered a cardiac arrest in the presence of a general practitioner equipped with a defibrillator, defibrillation was successful in 63. Fifty six reached hospital alive, of whom 47 were ultimately discharged home. In 71 cases the cardiac arrest had been either ventricular fibrillation or ventricular tachycardia and therefore potentially correctable. Defibrillation was often effective even when cardiac arrest had occurred before the general practitioner's arrival, particularly when it was attempted within 10 minutes of collapse. Eighteen of 36 patients who had received cardiopulmonary resuscitation from a bystander before the doctor's arrival survived to leave hospital. Even when cardiopulmonary resuscitation had not been given by a bystander, an appreciable proportion—seven of 33—survived when defibrillation was performed within the same time limit.

Our results show the importance of defibrillation provided by general practitioners. All front line ambulances should now carry a defibrillator, and our results endorse the British Heart Foundation's guidelines on managing patients with myocardial infarction.² These guidelines recommend a dual response by both the ambulance service and a general practitioner in such cases, so that a defibrillator is available as soon as possible.

M C COLQUHOUN
General practitioner

Court Road Surgery,
Malvern WR14 3BL

B L PENTECOST
Medical director

British Heart Foundation,
London W1H 4DH

¹ Avery A, Pringle M. Emergency care in general practice. *BMJ* 1995;310:6. (7 January.)

² Weston CFM, Penny WJ, Julian DG, on behalf of the British Heart Foundation Working Group. Guidelines for the management of patients with myocardial infarction. *BMJ* 1994;308:767-71.

Carrying heavy equipment may be impractical

EDITOR,—General practitioners are exhorted to carry nebulisers, electrocardiographs, and defibrillators while on call.¹ How can I, a woman weighing 54 kg, be expected to carry an electrocardiograph