So far the only justification for the laparoscopic method is a possible marginal lessening of post-operative discomfort, but is this so important when one considers the complexity and risks of the procedure compared with those of the mesh technique?

M J NOTARAS
Consultant surgeon

London Hernia Centre,
London NW6 2QA

- 1 Nixon S. It's all right now. *BMJ* 1995;310:198. (21 January.)
- 2 Amid PI, Shulman AG, Lichtenstein IL. Critical scrutiny of the open "tension-free" hernioplasty. *Am J Surg* 1993;165:369-71.
- 3 Iles JDH. Mortality from elective hernia repair. *Journal of Abdominal Surgery* 1969;11:87-95.
- 4 Welsh DRJ, Alexander MAJ. The shouldice repair. *Surg Clin North Am* 1993;73:451-69.

Sedation in fiberoptic bronchoscopy

Further study is needed

EDITOR,—We agree with M Q F Hatton and colleagues that it is unnecessary to sedate a patient for a procedure for which local anaesthesia may be used satisfactorily.¹ Sedation is safe provided that the administrator has had specific training and that guidelines for the selection of patients, monitoring, resuscitation facilities, and use of supplemental oxygen are followed. The drugs and techniques used should carry a wide margin of safety.² Hatton and colleagues state that sedation for bronchoscopy is associated with major complications.^{3,4} We note that many of these complications were attributed to inadequate resuscitation equipment, the failure to give supplemental oxygen, or the use of opiates with benzodiazepines (a synergistic combination with a narrow safety margin).² Patients' comfort and their tolerance of bronchoscopy may be influenced by several factors, including the depth and quality of sedation and the effectiveness of airway anaesthesia.

Active agent or placebo was used to produce light sedation. Did the authors use a recognised objective assessment of sedation—for example, the Ramsay score⁵—to titrate the level of sedation? The failure of doctors to identify whether active treatment had been given (18% for phenoperidine and 47% for midazolam) suggests that many patients were not sedated. It is not surprising that subtherapeutic doses failed to make the procedure more tolerable.

We note that patients who received phenoperidine said that they would be disinclined to repeat the procedure. We agree that the high incidence of nausea, vomiting, and dysphoria and the lack of hypnosis associated with synthetic opiates might produce poor quality sedation and contribute to this disinclination. Did the authors record the amount of topical local anaesthetic given, and was a similar quantity used for all groups?

We suggest that a further study with an objective assessment of sedation and control of compounding variables, such as the amount of local anaesthesia, is required to answer whether sedation helps in fiberoptic bronchoscopy.

M R J PARKER
Intensive care research fellow
C J EDAY
Lecturer in anaesthesia
A H COOTE
Anaesthetic research fellow

Bristol Royal Infirmary,
Bristol BS2 8HW

- 1 Hatton MQF, Allen MB, Vathenen AS, Mellor E, Cooke NJ. Does sedation help in fiberoptic bronchoscopy? *BMJ* 1994; 309:1206-7. (5 November.)
- 2 Royal College of Surgeons of England. *Report of the working party on guidelines for sedation by non-anaesthetists*. London: RCS, 1993.
- 3 Credle WF, Smiddy JF, Elliott RC. Complications of fiberoptic bronchoscopy. *Am Rev Respir Dis* 1974;109:67-72.

- 4 Simpson FG, Arnold AG, Purvis A, Belfield PW, Muers MF, Cooke NJ. Postal survey of bronchoscopic practice by physicians in the UK. *Thorax* 1986;41:311-7.
- 5 Ramsay MAE, Savage TM, Simpson BRI, Goodwin R. Controlled sedation with alphaxalone-alphadolone. *BMJ* 1974;iii: 656-8.

Doses that produce amnesia should be given

EDITOR,—I have considerable reservations about M Q F Hatton and colleagues' conclusion that routine sedation has little part to play in patients undergoing a single diagnostic bronchoscopy.¹ The description of the patients' willingness to undergo a repeat procedure, as indicated on a visual analogue scale, makes it difficult to compare their results with others but presumably means that a substantial number of patients would not be happy to have the test repeated. I have been impressed by the number of patients who, when discussing a bronchoscopy, have said something to the effect of "I do not mind what you do to me as long as you put me to sleep." Certainly, previous studies have shown that up to a quarter of patients would not be prepared to have a repeat bronchoscopy.^{2,3}

My colleagues and I have recently published our results of using higher doses of midazolam (mean 0.24 mg/kg).⁴ This dose produces complete amnesia, and only one of 123 patients was not prepared to have the procedure repeated. Although there are concerns about complications associated with sedation, we have been using this technique for several years without serious problems. This is probably because we give the midazolam in small incremental doses over several minutes. If any problems arise the sedation can be rapidly reversed with intravenous flumazenil. Surely, if many un-sedated patients find the procedure so unpleasant that they are not prepared to have it repeated then sedation in doses that produce amnesia should be used.

T J WILLIAMS
Consultant physician

Kettering General Hospital,
Kettering,
Northamptonshire NN16 8UZ

- 1 Hatton MQF, Allen MB, Vathenen AS, Mellor E, Cooke NJ. Does sedation help in fiberoptic bronchoscopy? *BMJ* 1994; 309:1206-7. (5 November.)
- 2 Gomm SA, Jones M, Hilton AM, Barker PW. Comparison of rigid and fiberoptic bronchoscopy: the patient's view. *Thorax* 1990;45:333P.
- 3 MacFarlane JT, Storr A, Ward MJ, Roderick Smith WH. Safety, usefulness and acceptability of fiberoptic bronchoscopy in the elderly. *Age Ageing* 1981;10:127-31.
- 4 Williams TJ, Nicoulet I, Coleman E, McAleney C. Safety and patient acceptability of intravenous midazolam for fiberoptic bronchoscopy. *Resp Med* 1994;88:305-7.

Intravenous sedation is inappropriate in most minor procedures

EDITOR,—M Q F Hatton and colleagues' suggestion that intravenous sedation has little part to play in fiberoptic bronchoscopy has far reaching implications in all the surgical specialties.¹ I believe that intravenous sedation has little if any role in most minor procedures.

Many such procedures are carried out under sedation by junior medical staff, who often draw on their experience in casualty departments. The aim of using sedation under these circumstances is to make the procedure more comfortable for the patient. I have often, however, seen escalating doses of hypnotics being given without effect when analgesia would have been more appropriate. What makes such procedures unpleasant is not awareness but pain. There is a widespread, fundamental failure to distinguish between analgesia and sedation.

I believe that the root of this evil rests in casualty departments, where many doctors start their surgical training. There, many procedures—for example, manipulation of Colles' fractures—are

routinely carried out under intravenous sedation. Many casualty departments have policies forbidding senior house officers to use intravenous regional anaesthesia (Bier blocks) because of potential complications. Instead they encourage the use of intravenous sedation. I have never found sedation to be of any value in facilitating the manipulation of such fractures unless doses sufficient to induce general anaesthesia are used. This is because the patient wakes up as soon as pain is elicited by movement at the fracture site. All too often doctors accept patients' short lived pain and distress, hiding behind the theoretical side effect of amnesia. Even more surprising is that they are encouraged to do so by the nursing staff.

The relief of pain and suffering is the basis of our work. Surgical trainees and, indeed, all doctors and nurses should be encouraged to learn effective methods of analgesia through their reading, consultation with anaesthetic staff, and supervision by senior colleagues. Let us eradicate this barbaric practice of inappropriate intravenous sedation and banish midazolam from the cabinet.

FRASER W H SUTHERLAND
Senior house officer III

Department of Cardiac Surgery,
Glasgow Royal Infirmary,
Glasgow G4 0SF

- 1 Hatton MQF, Allen MB, Vathenen AS, Mellor E, Cooke NJ. Does sedation help in fiberoptic bronchoscopy? *BMJ* 1994; 309:1206-7. (5 November.)

Patients must be monitored

EDITOR,—M Q F Hatton and colleagues suggest that the routine use of intravenous sedation has little part to play when patients undergo a single diagnostic procedure.¹ We agree that the blanket administration of intravenous sedatives to all patients undergoing endoscopic procedures is inappropriate, but we believe that these drugs have a useful role in helping a small number of patients to accept an often distressing procedure. In this context it is important to emphasise monitoring and the correct dose of drugs to ensure the safety of patients. In view of guidelines on sedation by non-anaesthetists issued by a working party of the Royal College of Surgeons we are concerned that the authors fail to mention if, or how, patients were monitored during the procedure.²

In simple sedation, reassurance and hypnotic suggestion, together with the use of drugs, produce a state of depression of the central nervous system during which communication is maintained so that patients respond to instruction. If this is lost then the patient has, for all intents and purposes, received a general anaesthetic with all its attendant risks. The authors do not define the term "light sedation." We find this misleading, as by implication a state of deep sedation must exist. What features delineate clearly the transition between the two? Does not deep sedation simply constitute a state of general anaesthesia as defined above? Although we are told that the dose of drug was titrated to achieve light sedation, most patients received the same total dose. In view of the varying susceptibility of patients to intravenous sedatives, it would be surprising if all patients achieved the same degree of light sedation.

Increasing numbers of patients are undergoing endoscopic procedures as day cases and as a result are exposed to the hazards of intravenous sedation. We applaud the authors for questioning and discouraging the routine use of intravenous sedatives in such procedures but would emphasise that, when they are used, the guidelines suggested by the Royal College of Surgeons should be followed.

GARETH THOMAS
Senior registrar in gastroenterology
CHRISTINE McBETH
Registrar in anaesthetics

University Hospital of Wales,
Cardiff CF4 4XW

- 1 Hatton MQF, Allen MB, Vathenen AS, Mellor E, Cooke NJ. Does sedation help in fiberoptic bronchoscopy? *BMJ* 1994; 309:1206-7. (5 November.)
- 2 Royal College of Surgeons of England. *Report of the working party on guidelines for sedation by non-anaesthetists*. London: RCS, 1993.

No grounds for abandoning sedation

EDITOR,—M Q F Hatton and colleagues suggest that in the case of fiberoptic bronchoscopy "routine sedation has little part to play in patients undergoing a single diagnostic procedure."¹ The patients in their study received both lignocaine and intravenous atropine before placebo or a sedative agent if the order reported was the order of administration. The taste of lignocaine and the side effects of atropine, the administration of which is questionable, are themselves distressing to some patients. It would have been more appropriate to give the sedative before these drugs, thus allowing the sedation to encompass the whole procedure. In addition, the operators found the procedure easier with sedation. Might this not have resulted in improved biopsy technique and yield?

There are no grounds for abandoning sedation on the basis of the findings of this study.

S P HANLEY
Consultant physician

Department of Chest Diseases,
North Manchester General Hospital,
Manchester M8 6RL

- 1 Hatton MQF, Allen MB, Vathenen AS, Mellor E, Cooke NJ. Does sedation help in fiberoptic bronchoscopy? *BMJ* 1994; 309:1206-7. (5 November.)

Immunisation against chickenpox

Pregnant women should be screened

EDITOR,—Lainie Friedman Ross and John D Lantos recommend that immunisation against chickenpox should be confined to those at high risk.¹ While they acknowledge the risk to susceptible pregnant women and their fetuses, however, they seem to include only immunocompromised children in their high risk category.

Even without the possible effects of a universal immunisation programme the average age at infection is showing a natural shift upwards,² and significant numbers of females of childbearing age are being infected.³ Antenatal patients are already screened for syphilis and for antibody to rubella. Would it not be prudent to screen those without a definite history of chickenpox for immunity and to offer immunisation to susceptible mothers postnatally?

D S G SLOAN
Consultant in public health medicine

Department of Public Health,
Argyll and Clyde Health Board,
Paisley PA2 7BN

- 1 Friedman Ross L, Lantos JD. Immunisation against chickenpox. *BMJ* 1995;310:2-3. (7 January.)
- 2 Miller E, Vandien J, Farrington P. Shift in age in chickenpox. *Lancet* 1993;341:307-8.
- 3 Sloan DSG, Burlison A. Shift in age in chickenpox. *Lancet* 1992;340:974.

Good argument exists for universal vaccination

EDITOR,—We wish to make three comments in response to Lainie Friedman Ross and John D Lantos's editorial on immunisation against chickenpox.¹ Firstly, the implicit general proposition that a disease must be potentially fatal or at least a source of serious morbidity to justify prevention by vaccination is hardly tenable in a profession whose objective is to improve quality of

life. If a vaccine has been proved to be safe the ideal circumstances is for it to be available to all citizens at possible risk of the disease.

Secondly, the posited detrimental outcome of the possible waning of the effect of childhood vaccination is based on an assumption that "illness and teratogenicity" will be more common and more severe in vaccinated patients than unvaccinated or previously infected patients. There is no evidence to support this view from studies of live attenuated varicella zoster vaccines.^{2,3}

Finally, one of the main consequential problems of chickenpox is herpes zoster in adults and particularly neuralgia after herpes zoster. In the experience of one of us (GRBS) this is one of the most intractable problems not related to cancer that arises in clinical practice; it seriously prejudices the last years of elderly—usually quite elderly—patients: GRBS has failed to find a therapeutic strategy that appreciably relieves neuralgic symptoms. Present thinking would argue that if chickenpox was prevented then herpes zoster would not occur. We contend that an effective inactivated varicella zoster vaccine might be preferred to a live vaccine, and as there is already one report of herpes zoster after vaccination⁴ it is difficult to exclude the possibility that the vaccine virus or a recombinant vaccine field virus might result in herpes zoster in later later life. A mixed regimen with a primary inactivated vaccine—to prevent colonisation of the ganglia by live attenuated virus—followed by a live attenuated virus vaccine might be a useful compromise.⁵

On these grounds we believe that there is a good argument for universal vaccination against chickenpox and that until further evidence is available, inactivated vaccines may be the safer option.

GORDON R B SKINNER
Senior lecturer
JUDITH DAVIES
Research associate

Department of Infection,
University of Birmingham,
Medical School,
Birmingham B15 2TT

- 1 Friedman Ross L, Lantos JD. Immunisation against chickenpox. *BMJ* 1995;310:2-3. (7 January.)
- 2 Andre FE. Worldwide experience with the Oka-strain live varicella vaccine. *Postgrad Med* 1985;61(suppl 4):113-20.
- 3 Gershon AA. Live attenuated varicella vaccine. *J Infect Dis* 1985;152:859-62.
- 4 Williams DL, Gershon AA, Gelb LD, Spraker MK, Steinberg S, Ragab AH. Herpes zoster following varicella vaccine in a child with acute lymphocytic leukemia. *J Pediatr* 1985;106:259-61.
- 5 Davies JA, Hallworth JA, Mcleish P, Randall S, Martin BAB, Buchan A, et al. Characterisation and immunogenicity of a candidate subunit vaccine against varicella zoster virus. *Med Microbiol Immunol* 1994;183:105-17.

Authors' reply

EDITOR,—We agree with D S G Sloan that women of childbearing age are at high risk of serious chickenpox. In fact, both male and female adolescents and young adults who are not immune are at high risk. We question, however, whether the upward shift in the average age at infection is a true trend. Although epidemiological studies of military recruits in the United States have found a rising rate of seronegativity for varicella zoster virus,¹ this can be explained both by the increased number of recruits from island nations or territories where chickenpox has always occurred at older ages^{2,3} and by known associations between seronegativity for the virus and race and sex.⁴ Sloan's citation of the advancing age at infection in the British population⁵ does not address the question of whether it is more than a reflection of the increasing number of immigrants from tropical countries or other changing demographic features.

We also agree with Sloan's recommendation that voluntary immunisation should be offered to susceptible mothers postnatally. We would go one step further and recommend that varicella vaccine

should be given to all susceptible people at high risk, including (but not limited to) immunocompromised people and their families, health care workers, those taking care of small children, women of childbearing age before pregnancy, military recruits, and college students. Our recommendation is not, as Gordon R B Skinner and Judith Davies suggest, to deny varicella vaccine to any group, only to reject routine immunisation of healthy young children at low risk.

Finally, Skinner and Davies's concern about neuralgia after herpes zoster in elderly people is valid, but the commercial vaccines that are available do not prevent herpes zoster.⁶ If a varicella vaccine offering lifelong immunity and preventing herpes zoster becomes available then we will reconsider our recommendations against universal vaccination.

LAINIE FRIEDMAN ROSS
Associate professor of paediatrics

MacLean Center for Clinical Medical Ethics,
University of Chicago,
Chicago,
IL 60637,
USA

JOHN D LANTOS
Associate professor of paediatrics

Department of Paediatrics,
La Rabida Children's Hospital,
Chicago,
IL 60649,
USA

- 1 Struwing JP, Hyams KC, Tueller JE, Gray GC. The risk of measles, mumps, and varicella among young adults: a serosurvey of US Navy and Marine Corps recruits. *Am J Public Health* 1993;83:1717-20.
- 2 Longfield JN, Winn RE, Gibson RL, Juchau SV, Hoffman PV. Varicella outbreaks in army recruits from Puerto Rico. Varicella susceptibility in a population from the tropics. *Arch Intern Med* 1990;150:970-3.
- 3 Maretic Z, Cooray MPM. Comparisons between chickenpox in a tropical and a European country. *J Trop Med Hyg* 1963;66:311-5.
- 4 Miller E, Vurdien J, Farrington P. Shift in age in chickenpox. *Lancet* 1993;341:308-9.
- 5 Garnett GP, Grenfell BT. The epidemiology of varicella-zoster virus infections: the influence of varicella on the prevalence of herpes zoster. *Epidemiol Infect* 1992;108:513-28.

Spontaneous fractures in cerebral palsy

EDITOR,—Last year a patient of ours presented to hospital with a chest infection. She was 19 years old, had cerebral palsy, and at the time was chronically malnourished because of feeding difficulties. On the day after the girl's admission her mother noticed a change in her behaviour and that she was sitting on a bean bag in a position that her mother thought was uncomfortable. Later it was realised that her thighs were swollen. Subsequent x ray films showed an oblique fracture of the left femur and a greenstick fracture of the right femur. The fractures were recent and the bones demineralised and thinned.

Had this child had only one fracture we might have considered that it was a spontaneous fracture. Probably, however, she had been put on the bean bag by a member of staff who did not appreciate her physical deformity and tried to put her in a "good" position. Her mother, visiting shortly afterwards, was aware of her daughter's mood change—something difficult for staff meeting the child for the first time to appreciate.

Given the history of this case, we suggest that S Lingam and Jane Joester may be incorrect in describing as spontaneous the fractures that they report.¹ Our patient was cared for at home, and her father commented that she liked her legs to be passively moved, flexed, and extended. Forced abduction and external rotation to achieve a "good" sitting position probably caused the fractures.

We have previously seen a 12 year old boy with microcephaly and spina bifida and an unexplained fracture of the femur; he was malnourished and had thin bones. His fracture occurred while his