adequate measures to ensure the safety of those who perform abortions or work in facilities where the procedure is performed. Doctors and other health care workers who dedicate their lives to meeting the health care needs of Americans should be ensured the right to work in a safe environment. Violence against health care workers is unacceptable. The American Medical Association has long condemned acts of terrorism against health care workers and will continue to do so.

JAMES S TODD
Executive vice president

American Medical Association,
515 North State Street,
Chicago, IL 60610,
USA

1 Ironies in the abortion war. *BMJ* 1995;310:267. (28 January.)

Suing for unpaid overtime payments

EDITOR,—In response to the news headline stating that three junior doctors are suing Raigmore Hospital Trust for unpaid overtime payments¹ I would like to point out that doctors at the Luton and Dunstable Hospital NHS Trust have recently been successful in their claim for increased overtime payments. Juniors demanded that each additional duty hour worked since they started work at the trust should be paid at class 2 instead of class 3 rates. After negotiation the claim was met in full by the trust, which had agreed to abide by the BMA's guidelines for working conditions. Each junior received about £1000 extra pay per six months worked.

Luton and Dunstable NHS Trust has probably set a precedent in this matter, and juniors all over Britain can probably pursue similar retrospective claims for money they are entitled to.

RICHARD STONE
House officer

Royal London Hospital,
London E1 1BB

1 British junior doctors sue for unpaid overtime. *BMJ* 1995;310:348. (11 February.)

Electoral reform of GMSC

EDITOR,—As a member of the General Medical Services Committee of many years' standing who may have to answer to an electorate of ordinary general practitioners, J David Watts has one view of the need for changes in the method of election to the committee.¹ I, as one of those ordinary doctors, have another view.² It is perhaps understandable that they differ considerably. I am grateful to Watts, however, for telling us why Leicestershire's motion at the BMA's annual representative meeting last year was deliberately placed low on the order paper.

What Watts seems not to understand is that the current, somewhat archaic system, unchanged in "its constitution and procedures" in any major way for over 80 years,^{3,4} has finally given way under the pressures of the late 20th century. A committee that represents doctors and directly negotiates with the government on behalf of doctors should be directly answerable to those doctors for its performance. I find it intriguing and revealing that Watts calls my argument in favour of democracy "spurious."

Watts's argument that the local medical committees raise the quota for the defence fund is weakened by the fact that they raise it from general practitioners and that it is the general practitioners who are expressing their dissatisfaction with the present system by encouraging me in my efforts to reform the voting procedures of the GMSC. These doctors feel remote and unable to influence events on a national scale. Direct elections would give

them that influence, would keep GMSC members' eyes more closely on the ball, and would allow a greater range of candidates for seats on the committee.

Under my proposed procedure, local medical committees would remain untouched and would still be able to exert an influence on the GMSC. Members of local medical committees would be able to stand, with other general practitioners, for election from the larger GMSC constituencies. The difference is that all general practitioners would be able to express their vote for them. Is this suggestion "spurious democracy"? I maintain that it is far better that the GMSC carries out the "collective will" of all general practitioners rather than just that of local medical committees.

For the GMSC to resist the first step on this road to democracy—a referendum of all general practitioners carried out by the Electoral Reform Society—is to show either a Luddite mentality or electoral cowardice. In the meantime, general practitioners' morale remains low, recruitment is minimal, and retirements are increasing, and the government remains supreme. The GMSC cannot absolve itself from some responsibility for this.

Finally, it is perhaps as well for the future of general practice that neither Watts nor the GMSC can influence any requisition for an extraordinary general meeting of the BMA. A referendum would be preferable, but, one way or the other, ordinary general practitioners will speak.

DAVID ROBERTS
General practitioner

Welford,
Northampton NN6 6HG

1 Watts JD. Reform of GMSC not needed. *BMJ* 1995;310:469 (18 February.)

2 Roberts D. Reform of GMSC needed. *BMJ* 1994;309:1585. (10 December.)

3 GMSC. *GMSC 75 1913-1988. The General Medical Services Committee and local medical committees.* London: GMSC, 1988:16.

4 Ellis N. *GMSC and LMCs.* London: GMSC, 1985:2.

Junior doctors' hours

Position of the "English clause" is uncertain

EDITOR,—During the negotiations that led to the "heads of agreement" on junior doctors' hours in 1991 I gained general agreement from those concerned—namely, the junior doctors, the Joint Consultants Committee, the medical royal colleges, and the Department of Health—for what at that time became known as the "English clause." This was contained in paragraph 6 of the guidance from the Conference of Medical Royal Colleges on the training implications of the heads of agreement.¹ It read:

2. The arrangements for those in General Professional Training (GPT) are different from those in Higher Specialist Training (HST). In certain situations and under defined circumstances higher specialist trainees should continue to be available for duties in excess of 72 hours maximum average per week (see RCS guidelines).

The Royal College of Surgeons' guidelines referred to (paragraph 11) were as follows:

At a meeting of the Technical Group on Junior Doctors' Hours on 16 January 1991 the following statement was agreed, which has particular relevance to surgery.

"In certain situations higher specialist trainees should continue to be available for duties in excess of a 72 hour maximum average per week where it would be of benefit to their training and where they wish to do so, providing the proper supporting staff structure exists and providing the duties were not harmful either to the trainees or to patients. If management has concerns about such an agreement, they should consult the relevant Royal College."

The main reason for introducing this clause, which to my knowledge has never been rescinded,

was the widely held professional view that the quality of training among higher surgical trainees (registrars and senior registrars) depended on adequate exposure to emergencies and that this would be unlikely if their contracted hours were reduced to 72 a week. In addition, there was concern over the practicality of organising 1 in 5 rotas in surgical units in smaller hospitals, particularly as partial shift systems were not deemed appropriate for higher trainees where continuity of care of patients is more demanding and important.

NHS trusts and health authorities have been urged by the minister for health, Mr Gerald Malone, to ensure that no doctor should be contracted to work more than 72 hours on an on call rota in hard pressed posts or to work for more than 56 hours a week; changes in contracts must genuinely reflect reductions in rostered hours and in the intensity of hours of work. Thus it seems that, unless higher surgical trainees can be classified as being in posts that are not hard pressed, they will not be able to opt for continuing to be contracted for an average 83 hours a week (equivalent to a 1 in 4 rota) and the spirit and intention of the English clause will be effectively thwarted.

An unambiguous statement from the NHS Executive on the current position of the English clause is indicated and would, I know, be appreciated by employing authorities, postgraduate deans, consultants, and higher trainees.

TERENCE ENGLISH
Past president, Royal College of Surgeons (1989-92)
Papworth Hospital,
Cambridge CB3 8RE

1 NHS Management Executive. *Hours of work of doctors in training: guidance from the medical royal colleges and their faculties in the UK.* Leeds: NHSME, 1991.

NHS Executive's reply

EDITOR,—I can confirm that the English clause is very much part of the new deal. I wrote to chief executives and general managers of trusts and authorities in February last year with a plan of action for reducing hours. My letter specifically restated the criteria for those wishing to adopt the English clause. Since then I have reminded our task forces of those criteria and, indeed, we began to monitor the take up of the clause during the reporting round last September. I later wrote to the chairpeople of task forces to begin action on the monitoring exercise in February this year and emphasised the need to ensure that the clause was well understood. This was the first occasion on which its effect could be gauged fully.

Ministers tend, when making statements about the new deal, to dwell on the broader themes. But their letters are followed up by officials with detailed discussions of the implementation arrangements, who have not neglected the English clause.

GRAHAM WINYARD
Medical director

NHS Executive,
Leeds LS2 7UE

National survey of hospital patients

Has a management dominated agenda

EDITOR,—We ask patients about their satisfaction with health services in order to achieve greater responsiveness to their needs and thereby improve standards of care. Stephen Bruster and colleagues seem to have taken a limited view of the value of seeking patients' views.¹ Their focus on standards set out in the patient's charter and national charter inevitably leads to a management dominated agenda in the formulation of the questions. The investigators provide no evidence that standards

set out in the charter or the adapted American questionnaire they used cover the concerns of British patients.

It would be surprising if the satisfaction and views of diverse types of patients could be covered adequately by a general questionnaire. Bruster and colleagues' conclusion that many of the problems related to communication is not surprising since at least 21 of the 35 direct questions listed in table II are about communication, although they purport to be about other aspects of care. Among the questions on satisfaction, the patients were asked about the helpfulness and courtesy of staff, but we should expect more than this from health professionals. No questions were asked about practical aspects of care (for example, help with going to the toilet) or more intangible but equally important aspects of care, such as reassurance. Surprisingly, patients were not asked their views on the outcome of their treatment, a long recognised flaw in research into satisfaction among patients' and a major omission if the investigators really do regard patients as "expert witnesses in the care process."

Ensuring the validity of a questionnaire seeking to establish patients' satisfaction is vital if some of the problems the investigators refer to (for example, focus on easily measured items, hotel aspects of care) are not to be repeated, in this case on a grand scale. Methods of finding out what concerns patients are well developed and include the use of in depth interviewing and content analysis' as a prelude to designing questionnaires for larger scale use. In stroke care we have developed questionnaires to establish patients' and carers' satisfaction that cover concerns identified only from careful, in depth interviewing.^{4,5} Despite the large scale of Bruster and colleagues' survey we urge against its findings or methods being used as a gold standard.

PANDORA POUND
Research sociologist
SHAH EBRAHIM
Professor of clinical epidemiology

Department of Public Health,
Division of Population and Health Care Sciences,
Royal Free Hospital School of Medicine,
London NW3 2PF

- 1 Bruster S, Jarman B, Bosanquet N, Weston D, Erens R, Delbanco TL. National survey of hospital patients. *BMJ* 1994;309:1542-6. (10 December.)
- 2 Cleary PD, McNeil BJ. Patient satisfaction as an indicator of quality care. *Inquiry* 1988;25:25-36.
- 3 Fitzpatrick R, Boulton M. Qualitative methods for assessing health care. *Quality in Health Care* 1994;3:107-13.
- 4 Pound P, Gompertz P, Ebrahim S. Patients' satisfaction with stroke services. *Clinical Rehabilitation* 1994;8:7-17.
- 5 Pound P, Gompertz P, Ebrahim S. Development and results of a questionnaire to measure carer satisfaction after stroke. *J Epidemiol Community Health* 1993;47:500-5.

Authors' reply

EDITOR,—Pandora Pound and Shah Ebrahim suggest that patients are asked about their satisfaction in order to achieve greater responsiveness to their needs. One of the outcomes of our survey was that asking about satisfaction produces little useful information. Asking about patients' experiences, however, can help to improve standards of care. In our paper the results of the questions concerning satisfaction were shown simply to illustrate that answers to these types of questions are positive despite the obvious problems reported in questions related to patients' actual experiences.

Pound and Ebrahim's interpretation that we took a limited view of patients' satisfaction is, we think, due to a misunderstanding. The paper certainly focused on the standards set out in the patient's charter and national charter. We chose to highlight these as we thought that they would be of interest to a general readership, particularly as no national data had been published on them since the patient's charter was first published. The survey, however, had a much broader scope than was

reported in the paper. With a limit of 2000 words it is impossible to cover all aspects of the survey. In the paper we described 75 of the 221 questions that were asked, so about two thirds of the results were not reported. We hope to publish more specific results in the future on the management of pain, planning of discharges, communication, and nursing implications of the survey—and these will illustrate the practical aspects of the survey, which included questions on help with using the toilet, bathing, and eating and questions relating to the reassurance of patients.

A considerable amount of qualitative work, including focus groups with patients, was used in designing the questionnaire. The number of problems reported by patients in this survey suggests that most of the questions we asked concerned topics of real concern to them. In addition, answers to open ended questions in the survey did not highlight any major concerns that were not addressed directly in the questionnaire.

Regarding the gold standard, we indicated that the survey was a national one, with over 5000 randomly selected patients interviewed face to face and with an 86% response rate, which could be used as a national reference dataset.

STEPHEN BRUSTER
Research assistant
BRIAN JARMAN
Professor of primary health care
NICK BOSANQUET
Professor of health policy

Department of General Practice,
St Mary's Hospital Medical School,
Lisson Grove Health Centre,
London NW8 8EG

BOBERENS
Research director

Social and Community Planning Research,
London EC1V 0AX

Non-immunisation of children

Statistics on vaccination coverage may be a poor measure of practice performance

EDITOR,—Neil Simpson and colleagues cite homoeopathic and religious reasons for the non-immunisation of children.¹ My experience of inner city practice, however, is that repeated failure to respond to requests for vaccination, without active refusal, is also an important reason for "negative consent." The fact that in 40% of cases in Simpson and colleagues' study the parents did not respond or gave no clear reason for non-vaccination or the children were having delayed vaccinations is consistent with this.

I compared the experience of an inner city practice in giving the preschool booster to its target cohort with that of a suburban practice in Leicester for the quarter ending September 1994. I did this by examining the medical records and vaccination returns to the health authority. These cohorts were typical of the respective practices. The two practices are similar in their accessibility, the

vaccination clinics offered, the method of calling children, and the enthusiasm for vaccination, but their experience differs considerably. When the cumulative percentage vaccination coverage after successive call ups is compared the inner city practice is at a disadvantage because its population is less compliant (table). Not only does a lower proportion respond but those who do respond require more call ups, showing resistance to vaccination rather than refusal.

Among the children who remained unvaccinated active refusal was not a major problem (one child from the inner city practice) but poor compliance was. This manifested as persistent failure to attend (five children) or having had late primary courses (eight children), which delayed call up until after the cohort had been assessed for target payment. Unlike the suburban practice, the inner city practice found it essential to do home visits to achieve the top target. Four of the cohort were vaccinated in this way, but seven visits were necessary as patients were frequently out. In our experience most of the parents who regularly fail to bring children for vaccination are compliant with "on the spot" vaccination; this is consistent with findings elsewhere.²

The reasons for a poor response are complex but are probably partly related to increased mobility (3.5% of the suburban cohort compared with 13.5% of the inner city cohort joined the practices after the age of 3½). Employment status, receipt of social security benefits, single parenthood, immigration, and poor English have been identified as factors.^{2,3}

There is a role for health visitors in performing on the spot vaccination. Consideration should be given to including children in whom primary courses have been late in later cohorts to increase their chance of vaccination. Poor compliance greatly increases the effort required to reach top targets. It is clear that the value of statistics on vaccination coverage alone as a measure of practices' performance is doubtful.

DAVID J SHEPHERD
General practitioner

Saffron Group Practice,
Leicester LE2 6UL

- 1 Simpson N, Lenton S, Randall R. Parental refusal to have children immunised: extent and reasons. *BMJ* 1995;310:227. (28 January.)
- 2 Jones K, Fasher B, Hanson R, Burgess M, Isaacs D, Joshi P, et al. Immunisation status of casualty attenders: risk factors for non-compliance and attitudes to "on the spot" immunisation. *J Paediatr Child Health* 1992;28:451-4.
- 3 Whiting CM, Schabas RF, Ashley MJ. Hib vaccine coverage in children attending day-care/nursery school in East York. *Can J Public Health* 1990;81:152-5.

The Society of Homoeopaths has no official policy on vaccination

EDITOR,—I wish to correct misleading information concerning the Society of Homoeopaths given in Neil Simpson and colleagues' paper on parental refusal to have children immunised.¹ The society

Vaccination coverage after successive call ups in suburban and inner city practices

	Suburban practice (n=142)		Inner city practice (n=168)	
	No vaccinated	Cumulative response (%)	No vaccinated	Cumulative response (%)
Attendance data available:				
1st Call	106	77.4	84	61.3
2nd Call	25	95.6	24	78.8
3rd Call	3	97.8	13	88.3
4th Call	2	99.3	6	92.7
7th Call	0	99.3	1	93.4
No response	1		9	
Attendance data unavailable:				
Joined practice after first call for preschool booster*	5		23	
Vaccinated urgently as was near end of quarter	0		5	
Not called because of late primary course	0		3	
No unvaccinated	1		17	

*The vaccination coverage of these children is worse than that for each cohort overall, implying that these children do not bias the attendance data that are available.