Editors of journals have the responsibility to ensure that material that is published is soundly based, conducted, and interpreted; if it is, then the source of the funding is immaterial. If editors have any doubts about any of these then they do not publish the paper; again, the source of the funding is immaterial.

If those who take money from commercial sources are required to identify their interests, what of other potential conflicts? For example, I might be a member of the Conservative party and consider that it was individual behaviour that accounted for most ill health and interpret the results of my study accordingly. Alternatively, as a member of the Labour party I would have an interest in showing how urban deprivation and poverty were the cardinal determinants of health. Smith is surely not suggesting that all conflicts of interest should be enumerated at the end of a paper; if they were, the disclaimer might take up considerably more space than the subject matter.

Another point needs consideration—namely, the source of funds for medical research. Without the support of commercial organisations many medical scientists would be unable to carry out any research, and to deny them the opportunity to publish their work because they have received funding from commercial or other sponsors seems indefensible. At present the Medical Research Council is not able to attend to everyone's needs, nor are other important sources of funding such as the Wellcome Trust. Many of my colleagues undertake work that is supported by drug companies, and I doubt that their judgment is adversely influenced by this. In case it is, however, might not an agreeable alternative be for all the commercial sponsors to put money into a central fund that was administered by an independent board, which could allocate the money irrespective of the outcomes hoped for by the primary sources of the funds?

JAMES BIRD
Consultant physician

St Mary's Hospital,
London W2 1NY

1 Smith R. Questioning academic integrity. *BMJ* 1994;309:1597-8. (17 December.)

Most people are not disgusted by the Portman group

EDITOR,—To bracket the Portman Group with an organisation such as Forest is grossly unfair, as anyone familiar with the group's activities will know.¹ It is untrue to say that most people are "disgusted" by our activities. On the contrary, our participation in a wide range of prevention projects has been welcomed by everyone who recognises that the drinks industry has a role in tackling alcohol misuse. Indeed, it has won over many people who were sceptical. Most of our work in research, education, and practical measures that target particular problems such as drinking and driving is done in partnership with government departments and non-government organisations. Richard Smith may be disgusted but most people are not.

I very much regret the controversy caused by the Portman Group's plan to publish a critical commentary on *Alcohol Policy and the Public Good*, not because I think that we acted improperly but because that impression has been erroneously given. There is genuine disagreement about how best to prevent alcohol misuse, and we have as much right as anyone else to argue for or against a particular approach. The fact that we wished to obtain expert contributions to present our view should be to our credit. We do not believe that we know all the answers but we do believe in a balanced debate. Surely it is not suggested that only one side in the debate should have access to expert advice.

Unfortunately, the controversy has played into the hands of those who want to see confrontation between the industry and the health community. I am determined to prevent that because I am convinced that collaboration without illusions is the better way. Smith advises doctors and academics to "consider very carefully the ethical implications of all their actions." I have always accepted that the Portman Group should do no less.

JOHN RAE
Director

Portman Group,
London W1M 7AA

1 Smith R. Questioning academic integrity. *BMJ* 1994;309:1597-8. (17 December.)

Referral of women with breast lumps

EDITOR,—Justine Foster¹ disagrees with J M Dixon and R E Mansel's suggestion that general practitioners should not aspirate breast lumps.² I can understand that a general practitioner finds that aspiration of a cyst provides instant relief, and it is true that a surgeon will aspirate the cyst anyway. A patient who is presenting with a first breast lump, however, is more likely to have a solid lump than a cyst. In this situation, many general practitioners will not obtain a satisfactory aspirate for cytological examination since it requires a high level of practice to obtain cells in most smears. The best opportunity for getting an early diagnosis is then hampered. Patients resent having the test repeated unnecessarily.

If a patient is known to have cysts then the balance of probability is different and it seems sensible for the general practitioner to aspirate a new lump on the assumption that it may well be another cyst and easily relieved.

B M HOGGIN
Consultant surgeon

Hove,
Sussex BN3 4GF

1 Foster J. Referral of women with breast lumps. *BMJ* 1995;310:331. (4 February.)

2 Dixon JM, Mansel RE. ABC of breast diseases: symptoms, assessment, and guidelines for referral. *BMJ* 1994;309:722-6. (17 September.)

Appropriate prescribing in asthma

PACT data have drawbacks

EDITOR,—Jeannette Naish and colleagues describe the variations in the costs of prescribing respiratory drugs among general practices in one family health services authority.¹ While it is possible that better asthma care results in higher prescribing costs, the use of costs from prescribing analysis and cost (PACT) data to derive the volume of prescribing for asthma has two major drawbacks.

Firstly, PACT data are not linked to indication, and therefore one cannot assume that all bronchodilators or inhaled steroids prescribed are for use in asthma. For example, aggregate data from a research database based on general practice computer systems in the former Northern region show that 15% (1354/8985) of patients receiving inhaled steroids in 1992-3 had a diagnosis other than asthma. Thus the management of bronchitis and other obstructive airways diseases is a source of variation among practices in the prescribing of inhaled steroids.

Secondly, the cost of prescribing, as the authors acknowledge, is not a good measure of the amount of prescribing. The table shows the volume and cost of inhaled steroids prescribed in the 16 family health services authorities in the Northern and

Volume of prescribing and cost of inhaled beclomethasone, budesonide, and fluticasone in Northern and Yorkshire Regional Health Authority, 1993-4

Family health services authority	No of defined daily doses prescribed	Net ingredient cost (£)	Net ingredient cost/defined daily dose (£)
Cleveland	5 703 028	2 207 108	0.39
Cumbria	6 493 063	1 849 211	0.28
Durham	7 250 846	2 471 500	0.34
Northumberland	4 994 130	1 595 383	0.32
Gateshead	2 502 342	812 058	0.32
Newcastle	3 578 339	1 332 644	0.37
North Tyneside	2 134 631	783 882	0.37
South Tyneside	1 599 347	618 189	0.39
Sunderland	3 427 872	1 158 173	0.34
Humberside	8 873 858	3 067 972	0.35
North Yorkshire	7 577 784	2 603 121	0.34
Bradford	7 820 934	1 747 018	0.22
Calderdale	2 818 016	765 748	0.27
Kirklees	5 572 910	1 436 207	0.26
Leeds	9 058 999	2 806 670	0.31
Wakefield	3 706 345	1 373 148	0.37
Regional health authority	83 112 444	26 628 032	0.32

Yorkshire region in 1993-4. This varies almost twofold (£0.22-0.39 per defined daily dose), indicating that one of the principal determinants of total cost is the choice of product. This is not surprising since the basic NHS cost of comparable inhaled steroid treatments ranges from £0.17/day (200 µg beclomethasone twice daily by pressurised inhaler) to £0.44/day (100 µg fluticasone twice daily by Diskhaler).

The ratio of the rate of prescribing of bronchodilators to that of inhaled steroids has been suggested as a marker of prescribing quality in respiratory diseases.² Ratios, however, are influenced by changes in the numerator or the denominator, or both, so the same ratio can derive from any absolute prescribing rate, large or small. High quality prescribing in asthma therefore requires an appropriate range of prescribing of both prophylactic and bronchodilator treatments. The optimum rate of consumption of antiasthma drugs remains unknown, and the difficulties in accurately expressing drug consumption should not be underestimated.

MARK CAMPBELL
Manager, prescribing unit
D NICHOLAS BATEMAN
Medical director

SARAH J ROBERTS
Lecturer in medical statistics

JAMES M SMITH
Regional pharmaceutical adviser

NHS Executive,
Northern and Yorkshire Regional Health Authority,
Regional Drug and Therapeutics Centre,
Wolfson Unit,
Newcastle upon Tyne NE1 4LP

1 Naish J, Sturdy P, Toon P. Appropriate prescribing in asthma and its related cost in east London. *BMJ* 1995;310:97-100. (14 January.)

2 Audit Commission. *A prescription for improvement*. London: HMSO, 1994:26-7.

NHS Executive promotes prophylaxis

EDITOR,—In their paper on appropriate prescribing in asthma in east London Jeannette Naish and colleagues highlight the difficulties in defining and measuring good prescribing.¹ In both the paper and the Editor's choice, statements are made that pressure to reduce the cost of prescribing in asthma may lead to a reduction in prophylactic treatment, lower the quality of asthma care, and be cost ineffective. The implication of these statements is that the government's policy for rational prescribing has created or is intended to create such pressure.

None of the initiatives or guidance on general practitioners' prescribing with which the NHS