marker for more cautious cyclists cannot be accepted uncritically.

Keatinge and Parry quote a secondary source to the effect that cycle use in Victoria decreased by 40% after wearing a helmet became compulsory. Cycling by children and teenagers decreased by an average of 36% in the two years after the law was introduced, but cycling overall increased.⁵

FRANCIS McDERMOTT
Chairman
JOHN LANE
Member

Victorian Road Trauma Committee,
Royal Australasian College of Surgeons,
Melbourne,
Victoria,
Australia

- 1 Keatinge R, Parry R. Protection afforded by cycle helmets. *BMJ* 1995;309:1441. (26 November.)
- 2 Spaitte DW, Murphy M, Criss EA, Valenzuela TD, Meislin HW. A prospective analysis of injury severity among helmeted and non-helmeted bicyclists involved in collisions with motor vehicles. *J Trauma* 1991;31:1510-6.
- 3 McDermott FT, Lane JC, Brazenor GA, Debney EA. The effectiveness of bicyclist helmets: a study of 1710 casualties. *J Trauma* 1993;34:835-45.
- 4 McDermott F, Lane J. Protection afforded by cycle helmets. *BMJ* 1994;309:877. (1 October.)
- 5 Cameron MH, Vulcan AP, Finch CF, Newstead SW. Mandatory bicycle helmet use following a decade of voluntary promotion in Victoria, Australia—an evaluation. *Accid Anal Prev* 1994; 26:325-37.

Pain in the neck, shoulder, and arm

Terminology used is unhelpful

EDITOR,—The series title “ABC of Rheumatology” implies that the fundamentals of the subject are clearly set out. The article on pain in the neck, shoulder, and arm, however, is likely to result in more confusion than enlightenment.¹ If “mild or moderate degenerative changes [in the neck] are often seen in asymptomatic individuals” then on what evidence do the authors state that “common causes [of pain referred to the arm include]... degenerative changes”? What are the distinguishing features that allow one to conclude that “degenerative changes, including apophysial joint or ligamentous hypertrophy and osteophytes,” are among these common causes? And what is meant by mechanical disorders? Does this term refer to a prolapsed cervical disc or include degenerative changes as well?

I agree that “early mobilisation or manipulative techniques... are usually helpful,” but where does the idea that “manipulation involves moving the joint beyond normal range” come from? If one attempted to do this it could result in dislocation or fracture.

What is meant by “periarticular disorders [of the shoulder]”? Is this the same thing as disorders of the rotator cuff? What is the evidence for stating that “impingement or tendinitis of the rotator cuff is the commonest problem [causing shoulder pain]”? And what is meant by impingement?

I find it difficult to understand why, if an injection is required to treat a disorder of the rotator cuff, it is given into the subacromial bursa. Surely infiltrating the part of the rotator cuff that contains the lesion would be more effective? Furthermore, the rotator cuff consists of the fibrous capsule of the shoulder joint blended with the tendons of the subscapularis, infraspinatus, and teres minor muscles; it is therefore inaccurate to refer to the “musculotendinous rotator cuff.”

GABRIEL SYMONDS
Medical director

Tokyo British Clinic,
Tokyo 150,
Japan

- 1 Barry M, Jenner JR. Pain in neck, shoulder, and arm. *BMJ* 1995;310:183-6. (21 January.)

Long acting steroid injections are safe and effective if given correctly

EDITOR,—The article on pain in the neck, shoulder, and arm advises on the choice of steroid preparations for intra-articular and soft tissue use in the conditions mentioned.¹ I disagree with the authors that the use of long acting depot preparations should be avoided. Several studies have shown that hydrocortisone acetate is the weakest and triamcinolone hexacetonide and triamcinolone acetonide are the most potent of the steroids currently available in terms of both efficacy and duration of action (M deSilva *et al*, 15th international congress of rheumatology, Paris, 1981).^{2,3} Furthermore, relatively large volumes of hydrocortisone acetate are needed for a reasonable dose of steroid, and this is particularly relevant in soft tissue injections for medial and lateral humeral epicondylitis, in which injections have to be made into tight restricted spaces. With the more potent preparations, smaller volumes can be used.

There has been some concern about the use of depot methylprednisolone acetate in soft tissue injections for, for example, the carpal tunnel syndrome. This relates mainly to the fact that this preparation, like hydrocortisone acetate, is a microcrystalline suspension so that crystals may be retained in soft tissues long after the injection. This also explains the postinjection flare seen more commonly with these preparations⁴ and is extremely rare with triamcinolone hexacetonide. Few of my patients have complained of postinjection pain after the use of this preparation for intra-articular and soft tissue injections, including for golfer's and tennis elbow. The important factor is that these preparations must be used in the proper dosage and not repeated more than once in superficial soft tissue sites. The need to repeat injections is usually due either to poor technique or to wrong diagnosis.

I also dispute the rationale of injecting steroids and local anaesthetic into the subacromial bursa for disorders of the rotator cuff when direct injection into the shoulder joint is the standard practice. Injection into the subacromial bursa would be more appropriate in acromioclavicular arthritis as direct access to the joint is not particularly easy.

MALCOLM DESILVA
Consultant physician

Department of Rheumatology,
Prince Charles Hospital,
Merthyr Tydfil,
Mid Glamorgan CF47 9DT

- 1 Barry M, Jenner JR. Pain in neck, shoulder, and arm. *BMJ* 1995;310:183-6. (21 January.)
- 2 Bain LS, Balch HW, Jacob R. Parenteral administration of 6 α -methylprednisolone-21-acetate. Part 1. Intra-articular injection: comparison with hydrocortisone acetate. *Ann Phys Med* 1967;9(2):43-8.
- 3 Bain LS, Balch HW, Wetherley JMR. Intra-articular triamcinolone hexacetonide: double blind comparison with methylprednisolone. *Br J Clin Pract* 1972;26:12.
- 4 McCarty DJ, Hogan JM. Inflammatory reaction after intrasynovial injection of microcrystalline adrenocorticosteroid esters. *Arthritis Rheum* 1964;7:359-67.

Authors' reply

EDITOR,—Malcolm DeSilva is inaccurate in stating that we recommended that long acting steroid preparations should be avoided for intra-articular injections. In fact, we did not recommend a particular preparation for intra-articular use. Conventional practice is to use long acting preparations when injecting into joints. For soft tissue injections, long acting preparations provide maximum benefit, but this has to be weighed against the greater tendency of these compounds to cause local tissue necrosis if they are not injected into a cavity or if they are accidentally infiltrated into the skin.^{1,2} Injection into a tendon may cause it to rupture.³ For general use we recommend hydrocortisone, although experienced practitioners may prefer to use a long acting preparation in certain situations.

We disagree that intra-articular injection of the shoulder joint is the standard practice for disorders of the rotator cuff. There is a close anatomical relation between the rotator cuff and the subacromial bursa, and reactive inflammation in this bursa is often present in tendinitis of the rotator cuff. The subacromial space or bursa is the recommended site of injection for treating the commoner causes of shoulder pain—namely, impingement, tendinitis of the rotator cuff, and subacromial bursitis.⁴

With regard to degenerative changes in the cervical spine, it is accepted that there is a high prevalence of asymptomatic radiological osteoarthritis in the population. When these changes are seen in a patient presenting with neck pain it therefore does not automatically follow that the neck pain is due to the osteoarthritic changes, and other reasons should be sought. In patients with neck and radicular symptoms, however, advanced osteoarthritic changes causing entrapment of a nerve root may be seen on magnetic resonance imaging. Gabriel Symonds agrees that mobilisation or manipulative techniques aimed at restoring the full range of movement may be helpful in treating neck disorders. In such cases, when a joint is restricted in movement mobilisation entails moving the joint within its range while manipulation entails moving the restricted joint beyond its “normal” range and attempting to improve the range or restore the full range.

MAURICE BARRY
Consultant rheumatologist

James Connolly Memorial Hospital,
Blanchardstown, Dublin 15,
Republic of Ireland

JUMBO JENNER
Consultant rheumatologist

Addenbrooke's NHS Trust,
Cambridge CB2 2QQ

- 1 Fitzgerald RH. Intrasynovial injection of steroids: uses and abuses. *Mayo Clin Proc* 1976;51:655-9.
- 2 Steinbrocker O, Neustadt DH. *Aspiration and injection therapy in arthritis and musculo-skeletal disorders*. Hagerstown, MD: Harper and Row, 1972.
- 3 Sweetnam R. Corticosteroid arthropathy and tendon rupture. *J Bone Joint Surg [Br]* 1969;51:397-8.
- 4 Dalton S. The shoulder. In: Klippel J, Dieppe P, eds. *Rheumatology*. London: Mosby-Year Book Europe, 1994.

Detection of prostate cancer

Recent evidence suggests screening may be justified in high risk younger men

EDITOR,—Screening for prostate cancer has been the subject of much debate, and Fritz H Schröder considers the published data.¹ His conclusion that population based screening is not yet justified is fair. There is much anxiety that screening for prostate specific antigen will detect a large number of indolent cancers, whose detection will not decrease mortality, and because of false positive results morbidity and mortality may be increased.^{1,2} Recent data from a nested case-control study, however, are an important addition to knowledge.³

Serum samples were taken from 68% of 22 071 doctors randomised in a continuing study of β carotene in 1982. Three hundred and thirty six men who provided serum samples developed prostate cancer during 10 years of follow up. Three aged matched controls who also supplied serum samples were selected. When a cut off concentration of prostate specific antigen of 4.0 ng/ml was used, at four years of follow up the sensitivity of detection was 87% for aggressive tumours but 53% for non-aggressive cancers. Specificity was more or less unchanged over time at 91%. Nearly 80% of all aggressive prostate cancers occurring within five years would have been detected by a single measurement of prostate specific antigen. Importantly, only 32 of 80 cancers arising more than five years after the sampling time were not aggressive.