be most effective. When in New Zealand, in 1990, I was most impressed by the fact that a judge was available in the hospital every Friday afternoon and reviewed all detentions as required. Is there any likelihood of such experienced legal minds being routinely available in Britain?

T H TURNER
Consultant psychiatrist

Division of Psychiatry,
Hackney Hospital,
London E9 6BE

1 Bradley C, Marshall M, Gath D. Why do so few patients appeal against detention under section 2 of the mental health act? *BMJ* 1995;310:364-7. (11 February.)

Possible problem with section 12

EDITOR,—Caroline Bradley and colleagues studied appeals against detention under section 2 of the Mental Health Act.¹ I recently came up against a problem with section 12 of the act, which gives psychiatrists the power to detain people under the act. I was involved in a case heard by the General Medical Council after I had failed to reapply for approval under section 12 (2) of the act when renewal was due. I had relied on the previous practice of Trent Regional Health Authority and the Department of Health's guidance on the process for approval of medical practitioners under section 12 (2) of the Mental Health Act 1983, which states at paragraph 4: "and subsequently they [the regional health authority] should remind him [the practitioner] that he should apply for renewal of approval if he so wishes."² The previous practice of Trent Regional Health Authority was to remind practitioners that their licence had expired, but it did not do so in this case.

All doctors in the Trent Regional Health Authority area should be warned that they cannot rely on the authority to follow the Department of Health's guidelines. For their own protection and to avoid entrapment, medical practitioners in other regions would be wise to discover whether those regions are following the department's guidelines. It is, of course, standard procedure to remind people when their motor licence, television licence, registration with the General Medical Council, or insurance policies are due.

JOHN HARDING PRICE
Press officer, Society of Clinical Psychiatrists

Lawn Medical Centre,
Lincoln LN2 5QT

1 Bradley C, Marshall M, Gath D. Why do so few patients appeal against detention under section 2 of the mental health act? *BMJ* 1995;310:364-7. (11 February.)

2 Department of Health. *Guidance on the process for approval of medical practitioners under section 12 (2) of MHA 1983*. London: DoH, 1990. (HC(90)21.)

Authors' reply

EDITOR,—Our reported rate of appeal against detention (27%) is much closer to the national rate (roughly 25%) than that of P C Naik and L Klenka. This is because our findings were based on data from nine hospitals.

Gordon Langley raises four questions: whether our citation of the Mental Health Act is accurate; whether patients are too confused to appeal; how far the code of practice is being applied; and whether the parallel system of appeals to tribunals and hospital managers is a good idea. We will deal with each in turn.

Langley is correct on the exact wording of the act. In our paper, however, we were not quoting the act but paraphrasing it. Our purpose was to indicate to general readers that the main considerations when deciding to detain a patient are the patient's health and safety and the protection of others. We do not dispute that the patient can be detained on grounds of health or safety.

With respect to the second question, there is

an important distinction to be made between being mentally ill and being too confused to make a rational decision to exercise one's rights. As reported in our study, six of the subjects approached were too unwell to be interviewed. The 40 subjects who were interviewed, however, were sufficiently coherent to give informed consent and to complete a reasonably demanding interview. It was startling to observe the surprise of some of these patients when they were informed of their right to appeal.

Langley's question on the code of practice is interesting. Answering this question was not one of the objectives of our study, but in the light of our findings it is an obvious direction for clinical audit.

With regard to the final question, in our retrospective study 75 of 104 patients who appealed did so to the mental health review tribunal only, 15 did so to the hospital managers only, and 14 did so to both the tribunal and managers. Of the 65 appeals that reached a hearing, six of 47 appeals to the tribunal, two of eight to the managers, and none of 10 double appeals resulted in discharge. We agree that the double appeal system is absurd.

MAX MARSHALL
Wellcome health services research fellow

CAROLINE BRADLEY
Wellcome research registrar

DENNIS GATH
Clinical reader

Department of Psychiatry,
Oxford University,
Warneford Hospital,
Oxford OX3 7JX

1 Department of Health. *Mental health review tribunals for England: annual report 1993*. London: HMSO, 1993.

APACHE scoring and prediction of survival in intensive care

EDITOR,—We are disappointed that Minerva has fallen into the same trap as others before her in misconstruing the role of the acute physiology and chronic health evaluation (APACHE) score in critically ill patients.¹ Although the APACHE systems can be used to provide information on the risks of death for a group of patients suffering from a specific disease category that may require admission to an intensive care unit, they cannot be used as predictors of the risk of death in individual patients and consequently should not be used to make decisions concerning withdrawal of treatment in seriously ill patients.

For example, the APACHE score on admission for a patient with diabetic ketoacidosis may be high because of considerable physiological disturbance, but the mortality from this condition is low. On the other hand, a low score on admission may be obtained after resuscitation of a patient with a catastrophic intracranial haemorrhage, while the prognosis may remain extremely poor. Additionally, APACHE scores on admission may be altered considerably by the interventions performed before admission to the intensive care unit—the so called "lead time bias."²

Misconceptions surrounding the use of the APACHE scoring system are common. Despite the contribution that APACHE scoring has made in clarifying severity of illness and to the concept of standardised mortality ratios it was not intended to be a method of predicting outcome in individual patients. Thus curtailing costs by limiting admission or continuing treatment according to scoring is invalid.³ In addition, APACHE scores cannot be used to decide who is to be admitted to an intensive care unit and should not be used to decide on a patient's discharge from an intensive care unit to either a ward or a high dependency area, as frequently patients' scores are similar in each of these areas.⁴ Finally, recent evidence suggests

that the ability of the APACHE system to predict outcome may be no better than the clinical judgment of physicians and nurses.⁵

S N PILKINGTON
Senior registrar

Department of Intensive Care Medicine,
Queen Alexandra Hospital,
Cosham,
Portsmouth PO6 3LY

- 1 Minerva. *BMJ* 1995;310:70. (7 January.)
- 2 Dragsted L, Jorgensen J, Jensen N-H, Bonsing E, Jacobsen E, Knaus WA, et al. Interhospital comparisons of patient outcome from intensive care: importance of lead-time bias. *Crit Care Med* 1989;17:418-22.
- 3 Civetta J, Hudson-Civetta J, Nelson L. Evaluation of APACHE II for cost containment and quality assurance. *Ann Surg* 1990;212:266-75.
- 4 Franklin C, Rackow EC, Mamdani B, Burke G, Weil MH. Triage considerations in medical intensive care. *Arch Intern Med* 1990;150:1455-9.
- 5 Johnson M, Corbett M, Fitzgerald F. Evaluation of prognostic stratification in medical intensive care unit patients using clinical judgment compared with APACHE, a severity of disease classification. *Chest* 1985;88:31S.

Sexual health education interventions for young people

EDITOR,—Ann Oakley and colleagues' methodological review of the effectiveness of sex education programmes draws attention to how few well designed studies have been carried out in this important area.¹ Wellings's book review of *Sex in America* gives no indication that this self styled "definitive survey" even asked about sex education.² However, Wellings and colleagues' national survey of sexual attitudes and lifestyles did ask nearly 20 000 Britons about sex education in considerable detail (in questions 17 and 18 of the long questionnaire).³ This must have yielded the most comprehensive British data available on the effects of sex education on such matters as age at first coitus and use of contraception. It is therefore curious that these data are not discussed in either of the published reports of the survey.⁴ The official reason given for this serious omission was that there had been insufficient time to analyse the data. Yet, over a year later, these results are still not available for public scrutiny.

If the authors of the largest ever British sex survey have given such a low priority to publishing their findings on the effects of sex education it is little wonder that, as Oakley and colleagues rightly lament, we are "still knitting without a pattern."

TREVOR G STAMMERS
General practitioner

Church Lane Practice,
London SW19 3NY

- 1 Oakley A, Fullerton D, Holland J, Arnold S, France-Dawson M, Kelley P, et al. Sexual health education interventions for young people: a methodological review. *BMJ* 1995;310:155-62. (21 January.)
- 2 Wellings K. Sex in America: a definitive survey [book review]. *BMJ* 1995;310:540. (25 February.)
- 3 Wellings K, Field J, Johnson A, Wadsworth J. *Sexual behaviour in Britain*. London: Penguin, 1994:393-4.
- 4 Johnson A, Wadsworth J, Wellings K, Field J. *Sexual attitudes and lifestyles*. Oxford: Blackwell, 1994.

Peripheral vascular disease—a term to be eschewed

EDITOR,—What's in a name? Quite a lot, if it happens to be seriously misleading. The term peripheral vascular disease is widely used as synonymous with obliterative arterial disease of the limbs. This fosters a double misconception since vascular care is limited neither to arterial disease nor to the periphery.

Misconceptions such as these are not trivial, since they have far reaching and damaging effects on clinical attitudes, referral practices, the provision of resources, research funding, training, and

the allocation of teaching time in the medical curriculum.

Arterial conditions treated by vascular specialists threaten life or limb; they encompass carotid, thoracic, renal, mesenteric, aortic, pelvic, and upper limb and lower limb occlusive and aneurysmal disease. Vascular disease, including of course its cardiac and cerebral components, is by a huge margin the commonest cause of death and disability in all developed countries—far exceeding all cancers put together. In recent years there has been a progressive increase in vascular workloads, reflecting the needs of an aging population.

In the past three years, consultant surgeon posts in the United Kingdom advertised as requiring a vascular interest have exceeded all other general surgical subspecialties added together. Such is the demand that many vascular consultants have recently been appointed some time before they have achieved accreditation.

The management of venous disease is undergoing fundamental changes. It has long been neglected: witness the frequency of recurrence of varicose veins and leg ulceration. Chronic venous disease costs the NHS more than £400m a year.¹ If venous disorders are to be managed successfully and efficiently the facilities of the vascular laboratory, vascular radiology, and leg ulcer nurse specialists and the attention of surgeons with a special interest are required.

The recent working party report on vascular services in Scotland has recommended that vascular services should be centralised.² There are several reasons why this makes sense, but perhaps the most compelling relates to resources. To meet the profound changes taking place in the management of vascular disease, reorientation is required on the part of clinicians and health care planners alike. How can vascular services be resourced or strategically planned or research funded if there is no clear perception of the scope and care of vascular disease? The term peripheral vascular disease is long past its "sell by" date.

C V RUCKLEY
Professor of vascular surgery

Department of Surgery,
Royal Infirmary,
Edinburgh EH3 9YW

- 1 Laing W. *Chronic venous diseases of the leg*. London: Office of Health Economics, 1992.
- 2 National Medical Advisory Committee Working Group. *Vascular surgical services in Scotland*. Edinburgh: HMSO, 1993.

Paternal irradiation and childhood leukaemia

EDITOR,—In 1990, Gardner *et al* reported that relatively high doses of ionising radiation measured by film badges worn by men while employed at the Sellafield nuclear installation before the conception of their children were statistically associated with the incidence of leukaemia among these children.¹ The authors suggested that this association was sufficient to account for the excess of childhood leukaemia in the village of Seascale, adjacent to Sellafield.

Since 1990 several other studies of paternal preconceptional irradiation have been reported.^{2,3} We have analysed the results of these studies, and the table compares the relative risk coefficients derived from linear relative risk models based on these datasets. Full details are given elsewhere.³

The association reported by Gardner *et al* is confined to children of the Sellafield workforce born in Seascale, whereas over 90% of children of Sellafield fathers were born outside Seascale.⁴ The relative risk coefficient for Seascale is statistically incompatible ($P < 0.01$) with all the other coefficients, including that for the rest of West Cumbria.³ The association is not found for doses due to internally deposited radionuclides or for external doses received shortly before conception,

Leukaemia relative risks per 100 mSv of cumulative paternal preconceptional dose derived from various datasets using a linear relative risk model

Dataset	Relative risk per 100 mSv (95% confidence interval)
Offspring of Sellafield workforce born in Seascale ²	36.04 (14.34 to 73.01)
Offspring of Sellafield workforce born elsewhere in West Cumbria ²	1.27 (0.51 to 3.36)
Offspring of Ontario radiation workforce ³	0.63 (<0.27 to 3.40)
Offspring of Scottish radiation workforce ³	<0.51 (<0.51 to 2.95)
Offspring of Hiroshima and Nagasaki A bomb survivors ³	<0.98 (<0.98 to 1.10)

nor does the association extend to other childhood cancers.² Furthermore, Kinlen has indicated that the paternal dose association is not sufficient to account for the excess cases in Seascale.⁵

We conclude that a causal interpretation of the association between paternal preconceptional irradiation and childhood leukaemia is untenable.³

MARK LITTLE
Principal scientific officer

National Radiological Protection Board,
Chilton,
Didcot OX11 0RQ

RICHARD WAKEFORD
Principal research scientist

Health and Safety Directorate,
British Nuclear Fuels,
Risley, Warrington WA3 6AS

MONTY CHARLES
Senior research fellow

School of Physics and Space Research,
University of Birmingham,
PO Box 363, Birmingham B15 2TT

- 1 Gardner MJ, Snee MP, Hall AJ, Powell CA, Downes S, Terrell JD. Results of case-control study of leukaemia and lymphoma among young people near Sellafield nuclear plant in West Cumbria. *BMJ* 1990;300:423-9.
- 2 Health and Safety Executive. *HSE investigation of leukaemia and other cancers in the children of male workers at Sellafield*. Sudbury: HSE Books, 1993.
- 3 Little MP, Wakeford R, Charles MW. A comparison of the risks of leukaemia in the offspring of the Sellafield workforce born in Seascale and those born elsewhere in West Cumbria with the risks in the offspring of the Ontario and Scottish workforces and the Japanese bomb survivors. *J Radiol Prot* 1994;14: 187-201.
- 4 Parker L, Craft AW, Smith J, Dickinson H, Wakeford R, Binks F, *et al*. Geographical distribution of preconceptional radiation doses to fathers employed at the Sellafield nuclear installation, West Cumbria. *BMJ* 1993;307:966-71.
- 5 Kinlen LJ. Can paternal preconceptional radiation account for the increase of leukaemia and non-Hodgkin's lymphoma in Seascale? *BMJ* 1993;306:1718-21.

Preventing crime and violence

A population approach is needed

EDITOR,—Violence continues to be a cause of substantial ill health and emerging debate.^{1,2} The incidence of reported violent crime is increasing³ and results not only in appreciable mortality and morbidity but also psychological and economic costs to the community.

The response to violent crime has traditionally been through the criminal justice system and has focused on retribution and rehabilitation.³ More recently, violence has been recognised as a public health problem, and epidemiological techniques are being used in the development and implementation of preventive strategies.⁴

In addressing the problem of violence most effort has been focused on a small group of vulnerable people. This approach, termed the high risk approach, aims at reducing levels of violence in those at higher risk. Not only is it difficult to identify accurately those at highest risk, however, but most acts of violence will probably be committed by people outside these high risk groups.

Thus if efforts are restricted to the minority, most violent crime will not be prevented.

An alternative approach, the "population strategy," recognises that extreme violence reflects the range of behaviour and circumstances of society as a whole.⁵ This approach suggests that there is not a clear demarcation between those who will commit violence and those who will not but a continuum within a society that tends to follow a normal distribution. The extreme tail of the distribution is therefore a function of the society as a whole, and if that society could be shifted to the left the number of extreme incidents would diminish. Conceptualising violence in these terms suggests that societal measures aimed at reducing important influences such as poverty, alcohol consumption, access to weapons, and television violence are likely to reduce these acts.

It would be valuable to develop a measure that would help us describe the distributions of the nature and form of violent behaviour in populations. This would help in exploring the causes of violence, comparing societies, and measuring the success of interventions.

High risk approaches will continue to be necessary to help the most vulnerable people in our society, but to establish enduring change in the culture of violence in societies a broader, population based approach will be needed.

SHELLEY RIFKIN
Research fellow
ANTHONY ZWI
Senior lecturer

Department of Public Health and Policy,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

- 1 Shepherd J, Farrington DP. Preventing crime and violence. *BMJ* 1995;310:271-2. (4 February.)
- 2 Alper JS. Biological influences on criminal behaviour: how good is the evidence? *BMJ* 1995;310:272-3. (4 February.)
- 3 Shepherd JP, Farrington DP. Assault as a public health problem: discussion paper. *J R Soc Med* 1993;86:89-92.
- 4 Rosenberg ML, Fenley MA. *Violence in America: a public health approach*. New York: Oxford University Press, 1991.
- 5 Rose G. *The strategy of preventive medicine*. Oxford: Oxford University Press, 1992.

Includes victims of domestic violence

EDITOR,—In the light of the increased awareness of violence as an important public health issue,¹ we wish to draw attention to the problem of domestic violence. This is a problem that doctors in Britain have been slow to address. In 1991 an estimated 530 000 incidents of domestic violence occurred in England and Wales; victims of domestic violence are more likely to sustain injuries than those involved in other violent incidents.²

Studies undertaken in the United States show that between 22% and 35% of women presenting in emergency departments are suffering from injuries or symptoms due to continuing abuse.³ In recognition of the seriousness of the problem the American Medical Association particularly focused on it in 1992, issuing diagnostic and treatment guidelines on domestic violence.⁴

Domestic violence and its consequences tend to continue over long periods. Early identification of victims is a key part of prevention, allowing appropriate advice and intervention to be offered, which may shorten the period over which abuse recurs. We should recognise that general practitioners and other members of the primary health care team, as well as staff of accident and emergency departments, have a particular responsibility to identify victims: guidelines may help in this, and their use has been shown to increase the numbers of women identified as victims.⁵

JO RICHARDSON
Clinical research fellow
GENE FEDER
Senior lecturer

Department of General Practice,
Medical College of St Bartholomew's and
the Royal London Hospitals,
London EC1M 6BQ