resources they may considerably underestimate true service requirements.

ANDREW C BURNETT
Director of primary care development

Kent Family Health Services Authority,
Maidstone ME14 1QH

PAULA BLAND
Information analyst

Thames Cancer Registry,
Surrey SM2 5PY

- 1 Richards C, Richards H, Pheby D. Skin cancer: how accurate are local data? *BMJ* 1995;310:503. (25 February.)
- 2 Kitz DS, Sluzarsz-Ladden C, Lecky JH. Hospital resources used for inpatient and ambulatory surgery. *Anaesthesiology* 1988;69:383-6.
- 3 Evans RC, Robinson GC. Surgical day care: measurements of the economic payoff. *Can Med Assoc J* 1980;123:873-80.

Computerised data collection systems have not improved data collection

EDITOR,—It is disappointing to learn that in the 10 years from 1984 to 1993 roughly 13% of patients with malignant melanoma living in Avon were not recorded by the South Western Regional Cancer Registry.¹ The disappointment is increased by the knowledge that a previous study performed in the same hospital showed that in 1974 the same cancer registry missed 8-19% of malignant melanomas in the Avon area.² The collection of data seems not to have improved in the past 20 years despite the introduction of computerisation.

This is supported by a closer examination of the figures provided by Clive Richards and colleagues.¹ In the three years (1984-6) before the introduction of the computerised histopathology systems in 1987 the cancer registry missed nine (4%) of 222 cases. In the four years 1990-3, the registry missed 73 (19%) of 386 cases.

J L BURTON
Professor

Department of Dermatology,
Bristol Royal Infirmary,
Bristol BS2 8HW

- 1 Richards C, Richards H, Pheby D. Skin cancer: how accurate are local data? *BMJ* 1995;310:503. (25 February.)
- 2 Beadle PC, Bullock D, Bedford G, Leach JF, Webb RA, Dent NA, et al. Accuracy of skin cancer incidence data in the United Kingdom. *Clin Exp Dermatol* 1982;7:255-60.

Aid to diagnosis of melanoma in primary medical care

Doubling of excisions of genuine disease is the important issue

EDITOR,—After reading the title of Christopher B Del Mar and Adèle C Green's paper, "Aid to diagnosis of melanoma in primary medical care,"¹ I thought, "Great: something to support the budgeting of medical photography departments," which have always been an important part of any dermatology service. I was therefore disappointed when I examined the findings more closely and realised that the authors had distorted the interpretation of their results to fit their original beliefs. Their main point is that in the city in which the intervention was carried out, where clinical photography was introduced into primary care, the percentage of lesions excised that were potentially or frankly malignant diminished. This was taken as evidence of the value of their algorithm and photographs in preventing unwarranted excision of benign lesions. The observation is something of a smokescreen as the main point of interest is, are we still going to catch melanomas?

Something odd happened during the study period—namely, that excision of potentially malignant lesions in the intervention city rose from 1.8% of all excisions (78/year) to 4.2% (136/year), whereas it decreased from 3.9% (90/year) to 3.4% (94/year) in the control city. Before the

intervention 606 melanocytic lesions (in total) were excised in the intervention city—that is, 80.6% of the number (752) excised in the control city; this proportion remained unchanged at 81% after intervention. The increase in excision of potentially malignant lesions in the intervention city is the explanation for the percentage reduction in the excision of benign lesions. Before the intervention 39 potentially malignant lesions were excised in the intervention city—87% of the 45 lesions removed in the control city; after intervention the proportion rose to 146% over two years. This increase is the equivalent of 91 potentially malignant lesions, or 4.6% of the total number of excisions performed in the intervention city; it wholly accounts for the 4.8% drop in benign excisions from 93.6% to 88.8% during the study.

There are many possible explanations for this finding, one being the normal fluctuation in statistics for melanoma. The most tempting explanation is that intervention resulted in increased attention to detail by clinicians, possibly resulting in the excision of malignant lesions that would not otherwise have been removed. It is also possible to show photographs to colleagues who are not available at the time of consultation, and this could have led to excision. An alternative explanation is that something was going wrong in the control city in that the percentage of malignant lesions excised there fell and the median Breslow thickness of melanomas excised increased (from 0.4 to 0.6 mm) whereas it fell in the intervention city (from 0.6 to 0.5 mm).

Whatever the interpretation of the findings, it is inappropriate to concentrate on the spurious percentage reduction in excisions of benign lesions in the intervention city when it is the doubling of excisions of genuine disease that is the important issue.

DAVID DE BERKER
Senior registrar

Department of Dermatology,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP

- 1 Del Mar CB, Green AC. Aid to diagnosis of melanoma medical care. *BMJ* 1995;310:492-5. (25 February.)

Authors' reply

EDITOR,—David de Berker is right to point to some of the weaknesses in our study, which we discussed. We wish that we could have randomised several cities to each arm of the study to increase the likelihood that any effect we found was attributable to the intervention and not to alternative factors in one city.

It is incorrect to ascribe the relative change in potentially malignant lesions to only the absolute increase in potentially malignant lesions (from 78 to 112 a year) in the intervention city. After the cameras and algorithm were introduced the number of benign lesions that were excised decreased by 22% (270 a year) in the intervention city compared with 18% (214 a year) in the control city.

When the increase in potentially malignant lesions is looked at in more detail much of the change is attributable to a disproportionate increase in dysplastic naevi and melanotic freckles of Hutchinson. Excisions of melanomas increased by only 5.7% in the intervention city compared with 4.6% in the control city, a difference that (just) fails to reach significance.

We are interested in the alternative explanations offered for our data. The suggestion that the algorithm and cameras might have caused more potentially malignant lesions to be excised (but not affected the rate of excision of benign lesions) is an exciting one that we were at pains to refrain from suggesting. Such an effect would have important implications—far greater than a reduction in unnecessary excisions—and would probably have far greater implications for the budget of the

medical photography unit at de Berker's hospital. Unfortunately, it is unlikely. After the intervention was introduced in the intervention city significantly fewer lesions were excised because of concern about malignancy and there was a trend for doctors to feel less pressure from patients to excise lesions. Most doctors who were offered the use of the algorithm and cameras thought that they had reduced the incidence of unnecessary excisions. Demographics changed in the intervention city after the intervention was introduced, resulting in an increase in the groups of people in whom melanoma is more likely—that is, older people and men.

We can only speculate on how access to the cameras and algorithm increased the numbers of dysplastic naevi and melanotic freckles of Hutchinson that were excised. Perhaps dysplastic naevi are more unstable with time and therefore distinguished from other banal naevi more readily by follow up after their appearance has been accurately recorded.

C B DEL MAR
Reader in general practice

University of Queensland Medical School,
Brisbane,
Queensland 4006,
Australia

A C GREEN
Principal research fellow

Queensland Institute of Medical Research,
Brisbane,
Queensland 4029,
Australia

Chemotherapy for solid tumours

EDITOR,—It is easy to sympathise with some of the points on both sides of the argument over whether chemotherapy should be given for solid tumours.¹ The nihilistic approach promulgated by G M Mead, however, would lead to no advances in treatment being made. As Mead points out, treatments that might benefit only a small percentage of patients will be sought by most patients—though not by their doctors, as pointed out by Slevin *et al.*² The dramatic improvement in response rates and survival in leukaemias, lymphomas, and testicular cancer would never have happened if aggressive, highly toxic treatments had not been tried. Those of us who treated breast cancer with adjuvant chemotherapy many years ago, long before it was an accepted treatment, repeatedly questioned whether it was worth while when otherwise well women engaged in pavlovian vomiting on entering the clinic. It has taken the data from tens of thousands of patients to show that this approach is worth while in premenopausal women with positive nodes.³ Moreover, these patients can now be treated with the new generation of antiemetics, such as ondansetron, when necessary, and treatment of breast cancer with chemotherapy has been reported to increase the quality of life in its own right.⁴

I believe that when a patient desires treatment but no proved efficacious treatment exists then he or she should be offered treatment only in a clinical trial in the first instance. Giving so many patients unproved treatments "off protocol" wastes resources and represents a loss of opportunities to enhance our knowledge and optimise future treatments. It is estimated that only 5% of patients in Britain are included in clinical trials; this figure should be raised well above the 50% mark. Only then will we be in a position to judge whether we are making real advances in management. The next 10 years will see immunotherapy and gene therapy joining the therapeutic repertoire, and it will be surprising if the dramatic responses seen in animal systems are not translatable, even in part, into some clinical scenarios. Only if this change in attitudes towards clinical trials occurs will the