be invalid because of the differing prenatal environments of monozygotic and dizygotic twins.

Clearly, however, low birth weight does not have the same importance in twins as in singletons. Studies based on the results of serial ultrasound scanning show that many twins have lower growth rates than singletons in the first trimester.² The fetal origins hypothesis states that coronary artery disease and its associated conditions result from growth retardation during later pregnancy and consequent disproportionate size at birth.³ Animal studies show that the setting of a low growth trajectory during early pregnancy protects the fetus against the effects of undernutrition in later pregnancy.⁴ Hence early down regulation of fetal growth may protect some twins from the effects of undernutrition in late gestation. A balance between the protective effects of early down regulation and the detrimental effects of later undernutrition may be the reason for the unexceptional overall mortality in twins and for the similar mortality of monozygotic and dizygotic twins.

DI W PHILLIPS
Clinical scientist

Metabolic Programming Group,
University of Southampton,
Southampton General Hospital,
Southampton SO16 6YD

- Christensen K, Vaupel JW, Holm NV, Yashin AY. Mortality among twins after age 6: fetal origins hypothesis versus twin method. *BMJ* 1995;310:432-5. (18 February.)
- Leveno KJ, Santos-Ramos R, Duenhoelter JH, Teisch JS, Whalley PJ. Sonar cephalometry in twins: a tale of biparietal diameters for normal twin fetuses and a comparison with singletons. *Am J Obstet Gynecol* 1979;135:727-30.
- Barker DJP, Gluckman PD, Godfrey KM, Harding JE, Owens JA, Robinson JS. Fetal nutrition and cardiovascular disease in adult life. *Lancet* 1993;341:938-41.
- McCance RA, Widdowson EM. Nutrition and growth. *Proc R Soc Lond B Biol Sci* 1974;185:1-17.

Temperature and cardiovascular mortality

EDITOR,—Owen L Lloyd has fallen into some of the common pitfalls when discussing cold and its effects on the body.¹ The environmental cold stress (rate of heat loss) experienced by warm blooded creatures is related not just to absolute temperatures outdoors or indoors but also to movement of air (wind or draughts)—for example, a body loses more heat at 10°C with a wind of 32 km/h than at -10°C in still air.² Similarly, humidity (rain or indoor damp) increases the heat loss, and, as Peter Wilmshurst states,³ the risk of ischaemic heart disease is also related to rainfall.²

Another factor is that the physiology of the body reacts to changing stimuli, rapid change producing more response than slow change. It also adapts to a steady state—for example, 20°C can induce shivering after exposure to 30°C, whereas 0°C feels warm to someone who has left a room at -30°C with wind from a fan. It is therefore to be expected that total cold stress (temperature, wind, and rain) would have more effect on health if the severity changed frequently and rapidly.

The social class gradient in risk of ischaemic heart disease (rising from I to V) is present throughout England and Wales. This can be at least partly explained by housing.⁴ People in the higher social classes tend to have centrally heated houses with minimal internal temperature gradients, whereas people in the lower social classes, with less money, live in houses which are often damp, draughty, and difficult to heat. If they heat only one room they have frequent changes in cold stress as they move around the house. In Glasgow, people in Drumchapel have a higher incidence of ischaemic heart disease than people in affluent neighbouring Bearsden.

The risk of ischaemic heart disease in the regional health authorities of England clearly rises as the climate becomes colder, wetter, windier, and more changeable, mainly to the west and

north.⁵ This even overrides the social class factor, so that people in social class I in north and northwest England have a higher risk than those in social class V in East Anglia.

Scotland is similar to England in that the west has more wind, more rain, and a more changeable climate than the east. On this basis it is easy to understand why the west of Scotland has a higher incidence of ischaemic heart disease and stroke than the east. There is therefore no Scottish paradox related to climate. Similarly, the genetic influence is negligible since people who change their place of residence rapidly acquire the risks of ischaemic heart disease of their new area of residence.⁴

E L LLOYD
Consultant anaesthetist

Department of Anaesthetics,
Western General NHS Trust,
Edinburgh EH4 2XU

- Lloyd OL. Temperature and cardiovascular mortality. *BMJ* 1995;310:467. (18 February.)
- Lloyd EL. *Hypothermia and cold stress*. London: Croom Helm, 1986.
- Wilmshurst P. Temperature and cardiovascular mortality. *BMJ* 1994;309:1029-30. (22 October.)
- Lloyd EL. The role of cold in ischaemic heart disease: a review. *Public Health* 1991;105:205-15.
- Balarajan R, Yuen P, Machin D. Inequalities in health: changes in RHAs in the past decade. *BMJ* 1987;294:1561-4.

Undernotification of whooping cough

EDITOR,—Douglas Jenkinson is to be commended for his painstaking documentation of whooping cough in general practice.¹ He points out that the incidence observed in his practice between 1977 and 1992 was six times the rate of notifications in England and Wales over the same period and highlights an appreciable number of cases among adults. This prompted me to examine notification data on whooping cough in Cardiff.

During 1984-92 there were 1034 notifications of whooping cough in Cardiff, giving an annual notification rate of 42 cases/100 000 population compared with the 26 cases/100 000 for England and Wales. The table shows the age distribution of patients in Cardiff compared with those reported by Jenkinson in Keyworth. Although there was a

Distribution of cases of whooping cough by age in Cardiff and Keyworth.¹ Figures are numbers (percentages)

Age (years)	Cardiff* (n=1013)	Keyworth† (n=500)
0	118 (11.6)	15 (3.0)
1	113 (11.2)	35 (7.0)
2	163 (16.1)	43 (8.6)
3	143 (14.1)	86 (17.2)
4	155 (15.3)	59 (11.8)
5	133 (13.1)	53 (10.6)
6	59 (5.8)	51 (10.2)
7	31 (3.1)	30 (6.0)
8	20 (2.0)	17 (3.4)
9	13 (1.3)	21 (4.2)
10-14	23 (2.3)	33 (6.6)
15-19	8 (0.8)	7 (1.4)
≥20	34 (3.4)	50 (10.0)

*Notification in 1984-92.

†Consecutive cases in general practice in Keyworth, 1977-92.

similar excess of cases in females (47% male and 53% female v 44% male and 56% female in Keyworth), there was a higher proportion of cases in young children and a lower proportion in older children and adults among the 1013 notified cases in Cardiff in which age was known. This suggests that either selective underdiagnosis or undernotification of adult cases is occurring. Since notification rates in Cardiff were broadly comparable to those for England and Wales as a whole this is probably not just a local phenomenon.

As whooping cough becomes less common, such irregularities in the notification system may

assume increasing importance. Coverage for the primary course of immunisation against whooping cough is now 92%, but many older children and young adults have never been immunised. A situation analogous to that with measles might therefore develop, with an upward shift in the age distribution of whooping cough. This has already been documented in the United States, with an increase in reported cases in subjects aged 10 and over from 15.1% during 1977-9 to 19.8% during 1980-9 and 26.9% during 1992-3.² This is an argument in favour of a pertussis vaccine booster as the opportunity for natural boosting of immunity through exposure to the disease diminishes.^{3,4} Furthermore, the use of a clear clinical definition of whooping cough⁵ and a greater awareness among doctors of the possibility of infection in adults would considerably enhance our ability to track the epidemiology of the disease.

MERION R EVANS
Consultant in communicable disease control
South Glamorgan Health Authority,
Cardiff CF4 3QX

- Jenkinson D. Natural course of 500 consecutive cases of whooping cough: a general practice population study. *BMJ* 1995;310:299-302. (4 February.)
- Resurgence of pertussis—United States, 1993. *MMWR* 1993;42:952-60.
- Novelli V, Al-Ansari H, Mok Q, Tasker R. Pertussis vaccination: is there a need for a booster dose? *Lancet* 1994;344:1225-6.
- Miller E, White JM, Fairley CK. Pertussis vaccination. *Lancet* 1994;344:1575-6.
- Centers for Disease Control. Case definitions for public health surveillance. *MMWR* 1990;39(RR-13):1-42.

Management of severe gastroparesis diabeticorum

EDITOR,—The debate over the management of gastroparesis diabeticorum reflects how difficult this condition is to manage.¹ We have had similar difficulties in managing a 46 year old man with longstanding diabetes treated with insulin.

The patient had various long term complications, including peripheral neuropathy, diabetic nephropathy (evident on renal biopsy), and impotence as well as hypertension and angina. He was admitted recurrently with abdominal pain, nausea, and vomiting. On each occasion he received intravenous fluids and insulin and a nasogastric tube was inserted. His symptoms were so severe that he was in hospital almost continuously over six months. Gastroparesis was confirmed by a radionuclide meal, which showed delayed gastric emptying of both solids and liquids. All proved therapeutic agents and withdrawal of his anti-hypertensive and antianginal agents were tried. His glycaemic control was optimised, but the only treatment that afforded relief was nasogastric intubation.

He was discharged with a nasojejunal feeding tube for a trial of enteral feeding, which was successful. We proceeded to permanent percutaneous endoscopic gastrojejunostomy feeding, the tube being inserted under local anaesthesia without complication. The double lumen system allowed drainage of retained gastric contents while facilitating jejunal feeding through a soft biliary tube. The patient was free of symptoms and remained at home for three months. He was readmitted with a recurrence of his symptoms due to blockage of the gastrojejunostomy tube. Attempts to replace the tube were unsuccessful. A gastrojejunostomy and feeding jejunostomy were fashioned surgically under local anaesthesia through a limited laparotomy; this was also unsuccessful. He developed intractable vomiting and worsening renal failure and died of metabolic complications.

We believe that our patient would not have withstood more radical surgical options; in any case, their efficacy has not been established.² The development of percutaneous gastrostomy tubes