Prevalence and outcome of symptomatic carotid lesions in young adults

Antonio Carolei, Carmine Marini, Patrizia Nencini, Carlo Gandolfo, Cristina Motto, Elietta Zanette, Massimiliano Prencipe, Cesare Fieschi, and the National Research Council Study Group

Abstract

Objective—To estimate the prevalence and outcome of symptomatic internal carotid artery lesions in young adults.

Design—Multicentre hospital based observational study with five year follow up.

Setting—Seven neurological departments in northern and central Italy.

Subjects—240 patients (115 men) aged 15–44 with a recent transient ischaemic attack or stroke in the carotid territory.

Main outcome measures—(a) Prevalence of symptomatic internal carotid artery stenosis or occlusion detected by continuous wave Doppler ultrasonography at entry; (b) incidence rates of cerebral, cardiac, and non-vascular death; non-fatal stroke; and non-fatal myocardial infarction.

Results—Carotid stenoses of 50–99% and occlusions were found in 38 patients (15.8%). Both conditions were significantly more frequent in patients aged over 35 and in those with hypertension, diabetes mellitus, and stroke at entry. The standardised mortality ratio at five years was 10.5 (95% confidence interval 5.0 to 19.3). Survival of patients with stenoses of 0–49% and occlusions was significantly better than that of patients with stenoses of 50–99%. Carotid stenosis of 50–99% was an independent predictor of death (hazard ratio 7.9; 95% confidence interval 2.2 to 29) and non-fatal stroke (hazard ratio 7.4; 1.5 to 37.4).

Conclusions—The prevalence of carotid stenosis or occlusion in young adults after a cerebrovascular event is low. Though patients with high grade symptomatic carotid stenosis are at risk of non-fatal and fatal events, patients with internal artery occlusion apparently have a benign prognosis.

Introduction

High grade stenoses and occlusions of the internal carotid artery occur in 30–50% of patients with cerebral ischaemia in the carotid territory.¹ Both are commonly associated with age, male sex, family history of stroke, obesity, smoking, hyperlipidaemia, diabetes mellitus, and coronary heart disease.^{2,3} The prognosis of severe stenoses is worse than that of the less tight lesions, with a 14% annual recurrence rate of stroke and a 9% mortality.⁴

Patients under 45 have a low prevalence of carotid lesions.^{5–9} Early atherosclerotic disease is the cause of most such lesions, though rare non-atherosclerotic causes are seldom represented.^{1–5,8} Long term follow up studies of young adults with focal cerebral ischaemia show the good outcome of patients without atherosclerotic lesions of neck arteries.^{5,7,10–11} In the presence of carotid stenosis the stroke recurrence rate increases to 20% after an average of 5.6 years of follow up.¹²

This report describes the prevalence and outcome of symptomatic internal carotid artery stenoses and occlusions, assessed by continuous wave Doppler ultrasonography, in a prospective hospital based series of young adults (age range 15–44) with either transient ischaemic attack or stroke at entry.

Patients and methods

Between April 1984 and March 1988 all patients aged 15–44 who had their first transient ischaemic attack or stroke in the ipsilateral carotid artery territory within eight weeks before admission were investigated in seven neurological departments in northern and central Italy.⁹ The study protocol included a clinical and laboratory evaluation of vascular risk factors, brain computed tomography, cardiac evaluation with chest radiography, electrocardiography, and transthoracic echocardiography. Selective carotid angiography of the symptomatic vascular territory was performed in most cases.

The presence of internal carotid artery stenosis or occlusion was assessed by continuous wave Doppler ultrasonography (DUD 800, Delalande, France). On the basis of the results patients were divided into three categories: 0–49% stenosis (< 6 kHz) of lumen diameter, 50–99% stenosis (6–20 kHz) of lumen diameter, and occlusion.¹³ Carotid occlusion was diagnosed in the presence of low flow velocities in the common carotid artery and no detectable Doppler signals distal to the occlusion. Interrater agreement among the trained examiners (verified before the study) resulted in the substantial range (κ index 0.75; 95% confidence interval 0.48 to 1.00).

All patients were followed up for five years with an annual clinical evaluation or a complete phone interview by means of a semistructured questionnaire. Hypertension was defined as diastolic blood pressure over 90 mm Hg (when measured out of the acute phase); diabetes mellitus was noted from the medical records before or at recruitment; obesity referred to body weight exceeding 20% of ideal (Lorentz formula).⁹ Primary end points included cerebral, cardiac, and non-vascular death; non-fatal stroke; and non-fatal myocardial infarction.¹⁴ Details of all deaths were assessed from hospital records, necropsy reports, or general practitioners' reports.

Cerebral death was defined as that occurring within 30 days after a severe qualifying stroke with clinically proved deterioration. Cardiac death was defined as death occurring unwitnessed or without any preceding symptoms (sudden death) or associated with cardiac symptoms suggestive of myocardial infarction, severe congestive heart failure, and acute pulmonary oedema. Stroke was defined as a focal neurological deficit of sudden onset persisting beyond 24 hours in a surviving patient.¹⁵ Myocardial infarction was diagnosed by the presence of at least two of the following: characteristic ischaemic chest pain for more than 30 minutes, typical increase in serum enzyme activities, and new pathological Q waves on electrocardiography. The only secondary end point considered was transient ischaemic attack, defined as a focal neurological deficit resolving completely within 24 hours.¹⁵ Its recurrence was evaluated only in patients with transient ischaemic attack on admission.

During follow up no compulsory treatment was adopted. Strict medical control and optimal treatment of the recognised vascular risk factors were encouraged.

Statistical analysis—Univariate statistical analysis was by Pearson's χ^2 test. χ^2 Test for trend was also

Clinica Neurologica,
Dipartimento di Medicina
Interna e Sanità Pubblica,
Università degli Studi di
L'Aquila, Ospedale
S Maria di Collemaggio,
67100 L'Aquila-
Collemaggio, Italy
Antonio Carolei, professor of
neurology
Carmine Marini, consultant
neurologist

Clinica Neurologica,
Firenze, Italy
Patrizia Nencini, consultant
neurologist

Clinica Neurologica,
Genoa, Italy
Carlo Gandolfo, professor of
neurology

Clinica Neurologica,
Ospedale Policlinico,
Milan, Italy
Cristina Motto, research
fellow

Dipartimento di Scienze
Neurologiche, Policlinico
Umberto I, Rome, Italy
Elietta Zanette, professor of
neurology
Massimiliano Prencipe, full
professor of neurology
Cesare Fieschi, full professor
of neurology

Members of the National
Research Council Study
Group who participated in
the study are listed at the end
of this report

Correspondence to:
Professor Carolei.

BMJ 1995;310:1363-6

TABLE 1—Risk factor distribution in 240 patients with internal carotid artery stenosis 0-49% and 50-99% and occlusion. Results expressed as numbers (percentages) of patients

Risk factor	Stenosis		Occlusion (n=8)	χ^2 Test	
	0-49% (n=202)	50-99% (n=30)		P value (global)	P value (linear trend)
Age $\geq$ 35 years	118 (58.4)	23 (76.7)	7 (87.5)	0.05	0.01
Male sex	96 (47.5)	14 (46.7)	5 (62.5)	0.70	0.59
Smoking > 20 cigarettes/day	117 (57.9)	19 (63.3)	4 (50.0)	0.77	0.80
Alcohol consumption > 100 g/day	25 (12.4)	4 (13.3)	2 (25.0)	0.59	0.40
Obesity	38 (18.8)	5 (16.7)	2 (25.0)	0.86	0.90
Hypertension	25 (12.4)	8 (26.7)	5 (62.5)	0.0002	0.00006
Diabetes mellitus	4 (2.0)	4 (13.3)	0	0.004	0.06
Serum cholesterol > 5.2 mmol/l	105 (52.0)	16 (53.3)	3 (37.5)	0.71	0.63
Serum triglycerides > 1.8 mmol/l	60 (29.7)	9 (30.0)	0	0.19	0.20
Serum high density lipoprotein cholesterol < 0.8 mmol/l	27 (13.4)	9 (30.0)	1 (12.5)	0.06	0.14
Packed cell volume > 0.43 in women, > 0.49 in men	27 (13.4)	7 (23.3)	0	0.17	0.85
Ischaemic heart disease	8 (4.0)	3 (10.0)	1 (12.5)	0.22	0.09
Stroke at entry	109 (54.0)	21 (70.0)	8 (100.0)	0.012	0.003

TABLE 2—Summary of end point occurrences in 240 patients with internal carotid artery stenosis of 0-49% and 50-99% and occlusion. Results expressed as numbers (percentages) of patients

End point	Stenosis		Occlusion	Total
	0-49%	50-99%		
No subsequent event	180 (89.1)	17 (56.7)	7 (87.5)	204 (85.0)
New transient ischaemic attack	9 (4.5)	4 (13.3)	0	13 (5.4)
Stroke	3 (1.5)	3 (10.0)	0	6 (2.5)
Myocardial infarction	3 (1.5)	1 (3.3)	1 (12.5)	5 (2.1)
All deaths	5 (2.5)	5 (16.7)	0	10 (4.2)
Cerebral	3 (1.5)	0	0	3 (1.3)
Cardiac	1 (0.5)	2 (6.7)	0	3 (1.3)
Non-vascular	0	3 (10.0)	0	3 (1.3)
Unknown cause	1 (0.5)	0	0	1 (0.4)
Lost to follow up	2 (1.0)	0	0	2 (0.8)
Total	202 (84.2)	30 (12.5)	8 (3.3)	240 (100.0)

TABLE 3—Percentage annual recurrence rates (95% confidence intervals) of new transient ischaemic attack, non-fatal and fatal stroke, and cardiac event in 238 patients with internal carotid artery stenosis of 0-49% and 50-99% and occlusion

Outcome	Stenosis		Occlusion (n=8)	All (n=238)
	0-49% (n=200)	50-99% (n=30)		
New transient ischaemic attack	1.57 (0.36 to 6.67)	4.72 (0.43 to 36.1)	0 (0 to 32.4)	1.85 (0.49 to 6.74)
Non-fatal and fatal strokes	0.40 (0.06 to 2.61)	2.09 (0.26 to 14.8)	0 (0 to 32.4)	0.60 (0.13 to 2.63)
Cardiac event	0.30 (0.04 to 2.44)	1.38 (0.12 to 13.7)	2.50 (0.01 to 35.7)	0.51 (0.01 to 2.49)

FIG 1—Observed and expected survival curves

performed to emphasise the linear relation between vascular risk factors and severity of carotid disease. Patients' event free survival was assessed by the Kaplan-Meier method according to the degree of carotid disease. Comparisons between the survival distribution for the different subgroups were performed by log rank test. Survival rate was also compared with that of a standard population, matched for age and sex, according to Italian life tables,¹⁶ the standardised mortality ratio being computed with exact 95% confidence interval. Univariate hazard ratios were calculated separately for death, non-fatal stroke, and non-fatal myocardial infarction by the Cox proportional hazards regression, stenosis of 0-49% being taken as the reference category. Adjusted hazard ratios were obtained with age, sex, cardiac disease, and transient ischaemic attack or stroke at entry being included as covariates.

Results

BASELINE CHARACTERISTICS

Two hundred and forty patients were included (125 women, 115 men; mean age 36.0 (SD 7.3) years). Most (n=138) presented with stroke as the first episode. Thirty eight patients (15.8%) had ipsilateral internal carotid artery stenosis of 50-99% (n=30) or occlusion (n=8) detected by Doppler ultrasonography. Seven stenoses (23.3%) were bilateral. Selective carotid angiography was performed on 175 patients (73%). In

144 (82%) cases there was agreement between the angiographic and ultrasound findings. The coefficient of agreement adjusted for chance (κ index) was 0.62 (95% confidence interval 0.48 to 0.76), indicating substantial concordance between the two methods.

Most patients (73%) were taking antiplatelet drugs (n=153) or anticoagulants (n=22) during follow up. Five patients with transient ischaemic attack and four with stroke at entry underwent carotid endarterectomy because of symptomatic preocclusive carotid stenosis (n=2), ipsilateral stenosis of greater (n=3) or less than 70% (n=1), bilateral stenosis of greater than 70% (n=2), and post-stenotic aneurysm (n=1), all confirmed by angiography.

Table I shows the linear association of age over 35 ($P=0.05$ (χ^2 test); $P=0.01$ (χ^2 test for trend)), hypertension ($P=0.0002$; $P=0.00006$), stroke at entry ($P=0.012$; $P=0.003$), and diabetes mellitus ($P=0.004$; $P=0.06$) with the severity of carotid disease.

ANALYSIS OF OUTCOME EVENTS

The five year clinical follow up (table II), based on information reported at the annual clinical evaluation (162 patients; 68%) or collected by phone interview (76; 32%), was available in 238 patients. Eight men and two women died (4.2%). Of the nine patients who underwent endarterectomy, one had a contralateral transient ischaemic attack five years after surgery, one died in a car accident, and one with bilateral stenosis had a stroke eight months after surgery and committed suicide four months later. The observed survival rate (95.8%) was worse than expected for the age and sex matched standard Italian population (99.6%) (standardised mortality ratio 10.5 (95% confidence interval 5.04 to 19.32)) (fig 1).

Table III shows the annual recurrence rates for cerebral and cardiac events in the three subgroups of carotid lesions and in the overall series. The cumulative risk of death among patients with carotid stenoses of 50-99% was higher ($P=0.0012$ (log rank test)) than among patients with stenoses of 0-49% and occlusions (fig 2).

Carotid stenosis of 50-99% was associated with

FIG 2—Kaplan-Meier cumulative risk of death in patients with internal carotid artery stenoses of 0-49% and 50-99% and occlusion

significant hazard ratios—namely, 7.2 (95% confidence interval 2.1 to 25.0) for all deaths and 7.5 (1.5 to 37.1) for non-fatal stroke (table IV). These hazard ratios remained significant after adjusting for other potential outcome predictors.

Discussions

Doppler ultrasonography has so far been formally assessed as a diagnostic tool and is likely to remain the first line screening technique for carotid lesions, while the most recently developed ultrasound methods and angiography may clarify the diagnosis in equivocal cases. The conclusions we have drawn apply to patients with continuous wave Doppler evaluation performed according to standardised criteria.¹³ The substantial agreement ($\kappa = 0.62$) between selective carotid angiography and ultrasound in 73% of patients (175/240) allowed an unbiased estimate of both prevalence and outcome of symptomatic internal carotid artery lesions in our hospital based series.

CAROTID LESIONS AND ASSOCIATED RISK FACTORS

The 15.8% prevalence (38 cases) of ipsilateral carotid stenoses of 50-99% and occlusions agreed with other reports referring to younger case series⁵⁻⁸ but was lower than that reported in elderly patients with cerebral ischaemia.¹ Bilateral severe carotid stenoses (seven of 30 cases; 23.3%) were more frequent than reported (8.5%).¹⁷ The association between the degree of carotid stenosis and the severity of cerebral ischaemia at entry was confirmed in our series.⁴ In addition, severe carotid stenoses and occlusions were associated with age over 35, hypertension, and diabetes mellitus.

The lack of an association with cigarette smoking, hypercholesterolaemia, and hypertriglyceridaemia, found in elderly patients,² might be due to lack of sensitivity of continuous wave Doppler ultrasonography, which did not identify non-stenosing carotid plaques and could not rule out carotid dissection or fibromuscular dysplasia. Alternatively, some vascular risk factors might require longer exposure to act as true atherogenic determinants of carotid disease progression. Their association with the extent and severity of carotid disease was stronger in elderly than in young patients.³ Different criteria to define and assess vascular risk factors or to categorise carotid abnormalities might also explain discrepancies with other studies.¹⁸

LONG TERM PROGNOSIS

The standardised mortality ratio (10.5) was higher than reported in other studies (ratios between 1.5 and 2.0) with a similar follow up period and in comparisons with age and sex matched population controls, reflecting the high risk of death among young adults with cerebral ischaemia in the carotid territory.^{14 19}

Annual incidence rates of cerebral (0.6%) and cardiac events (0.5%) were lower than reported in elderly people (4.2% and 2.9% respectively)¹⁴ but close

TABLE IV—Univariate and adjusted hazard ratios for death, non-fatal stroke, and non-fatal myocardial infarction by Cox regression analysis

Outcome	Subgroup of internal carotid artery stenosis or occlusion	Hazard ratio (95% confidence interval)	
		Univariate	Adjusted for age, sex, transient ischaemic attack or stroke at entry, and cardiac disease
All deaths	Stenosis 0-49%	1.0	1.0
	Stenosis 50-99%	7.2 (2.1 to 25.0)	7.9 (2.2 to 29.0)
	Occlusion	0	0
Non-fatal stroke	Stenosis 0-49%	1.0	1.0
	Stenosis 50-99%	7.5 (1.5 to 37.1)	7.4 (1.5 to 37.4)
	Occlusion	0	0
Non-fatal myocardial infarction	Stenosis 0-49%	1.0	1.0
	Stenosis 50-99%	2.5 (0.3 to 24.0)	3.0 (0.3 to 29.0)
	Occlusion	8.1 (0.8 to 77.8)	8.9 (0.7 to 111.0)

Key messages

- Continuous wave Doppler ultrasonography is a suitable first line, low cost screening technique for carotid lesions
- Young adults with a history of transient ischaemic attack and stroke have a low prevalence of high grade symptomatic carotid artery stenosis and occlusion
- The five year prognosis is apparently benign in patients aged 15-44 with internal carotid artery occlusion but poor (in terms of fatal and non-fatal events) in the presence of high grade carotid stenosis
- Primary and secondary medical prevention of cerebral ischaemia based on routine Doppler examinations should be initiated early in subjects at risk
- Endarterectomy should be considered even in young adults with ipsilateral carotid artery stenosis of 70-99% to halt the progression of arterial lesions and improve prognosis

to those found in patients under 30 (0.7% and nil).² The five year mortality was 16.7% in patients with severe carotid stenosis. This figure was roughly comparable to the 15% reported in older patients⁴ and 20% in patients younger than 50 followed up for an average of 5.6 years.¹² The small subgroup of young adults with carotid occlusion showed the apparently favourable prognosis reported in elderly subjects with either symptomatic or asymptomatic carotid occlusion.^{5 20-22}

In our series severe carotid stenosis of 50-99% contributed independently to increased mortality and the recurrence of non-fatal stroke in young adults (table IV).

CONCLUSIONS

This study shows that young adults with high grade symptomatic carotid stenosis are at risk of non-fatal and fatal events, whereas patients with carotid occlusion apparently have a benign prognosis. Therefore, in subjects at risk primary and secondary medical prevention should be initiated early.⁹ Furthermore, other workers suggest that endarterectomy should be considered, after thorough evaluation, even in young adults with ipsilateral high grade symptomatic carotid stenosis of 70-99% to halt the progression of arterial lesions and improve prognosis.^{17 23}

This study was funded by the National Research Council (grant 94.004671.PF40-95.2.572). We thank Dr N Bornstein, Professor J D Easton, and Professor M Motolese for comments and suggestions during the preparation of the manuscript.

Members of the National Research Council Study Group who participated were: M Frontoni (University of Rome); D Inzitari (University of Florence); C Finocchi (University of Genoa); R Totaro (University of L'Aquila); G Landi (Ospedale Maggiore, Milan); L De Zanche (Padua Hospital); U Scoditti (Parma Hospital).

- 1 Zhu CZ, Norris JW. Role of carotid stenosis in ischemic stroke. *Stroke* 1990;21:1131-4.
- 2 Bogousslavsky J, Regli F, Van Melle G. Risk factors and concomitants of internal carotid artery occlusion or stenosis. A controlled study of 159 cases. *Arch Neurol* 1985;42:864-7.
- 3 Tell GS, Crouse JR, Furberg CD. Relation between blood lipids, lipoproteins, and cerebrovascular atherosclerosis. A review. *Stroke* 1988;19:423-30.
- 4 Norris JW, Zhu CZ. Stroke risk and critical carotid stenosis. *J Neurol Neurosurg Psychiatry* 1990;53:235-7.
- 5 Lisovsky F, Rousseaux P. Cerebral infarction in young people. A study of 148 patients with early cerebral angiography. *Neurol Neurosurg Psychiatry* 1991;54:576-9.
- 6 Gautier JC, Pradat-Diehl P, Loron P, Lechat P, Lascault G, Juillard JB, et al. Accidents vasculaires cérébraux des sujets jeunes. Une étude de 133 patients âgés de 9 à 45 ans. *Rev Neuro (Paris)* 1989;145:437-42.

- 7 Bogousslavsky J, Regli F. Ischemic strokes in adults younger than 30 years of age. Cause and prognosis. *Arch Neurol* 1987;44:479-82.
- 8 Klein GM, Seland PT. Occlusive cerebrovascular disease in young adults. *Can J Neurol Sci* 1984;11:302-4.
- 9 Marini C, Carolei A, Roberts RS, Prencipe M, Gandolfo C, Inzitari D, *et al*. Focal cerebral ischemia in young adults: a collaborative case-control study. *Neuroepidemiology* 1993;12:70-81.
- 10 Hindfelt B, Nilsson O. Long-term prognosis of ischemic stroke in young adults. *Acta Neurol Scand* 1992;86:440-5.
- 11 Chancellor AM, Glasgow GL, Ockelford PA, Johns A, Smith J. Etiology, prognosis and hemostatic function after cerebral infarction in young adults. *Stroke* 1989;20:477-82.
- 12 Marshall J. The cause and prognosis of stroke in people under 50 years. *J Neurol Sci* 1982;53:473-88.
- 13 Spencer MP, Reid JM. Quantitation of carotid stenosis with continuous-wave (C-W) Doppler ultrasound. *Stroke* 1979;10:326-30.
- 14 Carolei A, Candelise L, Fiorelli M, Francucci BM, Motolese M, Fieschi C, *et al*. Long-term prognosis of transient ischemic attacks and reversible ischemic neurologic deficits: a hospital-based study. *Cerebrovasc Dis* 1992;2:266-72.
- 15 Millikan CH, Ad Hoc Committee on Cerebrovascular Disease. A classification and outline of cerebrovascular diseases II. *Stroke* 1975;6:565-616.
- 16 Istituto Nazionale di Statistica (ISTAT). *Cause di morte: anno 1988*. Vol 4. Rome: ISTAT, 1991:1-400.
- 17 North American Symptomatic Carotid Endarterectomy Trial Collaborators. Beneficial effect of carotid endarterectomy in symptomatic patients with high-grade carotid stenosis. *N Engl J Med* 1991;325:445-53.
- 18 Bots ML, Breslau PJ, Briët E, de Bruyn AM, van Vliet HHDM, van den Ouweland FA, *et al*. Cardiovascular determinants of carotid artery disease. The Rotterdam elderly study. *Hypertension* 1992;19:717-20.
- 19 Furlan AJ, Whisnant JP, Baker HL Jr. Long-term prognosis after carotid artery occlusion. *Neurology* 1980;30:986-8.
- 20 Bornstein NM, Norris JW. Benign outcome of carotid occlusion. *Neurology* 1989;39:6-8.
- 21 Hennerici M, Hulsbomer HB, Rautenberg W, Heffter H. Spontaneous history of asymptomatic internal carotid occlusion. *Stroke* 1986;17:718-22.
- 22 Cote R, Barnett HJM, Taylor DW. Internal carotid occlusion: a prospective study. *Stroke* 1983;14:898-902.
- 23 European Carotid Surgery Trialists' Collaborative Group. MRC European carotid surgery trial: interim results for symptomatic patients with severe (70-99%) or mild (0-29%) carotid stenosis. *Lancet* 1991;337:1235-43.

(Accepted 24 March 1995)

Mental disorders in young and middle aged men who commit suicide

Erkki Isometsä, Markus Henriksson, Mauri Marttunen, Martti Heikkinen, Hillevi Aro, Kimmo Kuoppasalmi, Jouko Lönnqvist

Department of Mental Health, National Public Health Institute, FIN-00300, Helsinki, Finland
Erkki Isometsä, senior researcher
Martti Heikkinen, senior researcher
Hillevi Aro, professor
Jouko Lönnqvist, research professor

Department of Psychiatry, Helsinki University, Helsinki
Markus Henriksson, senior lecturer
Mauri Marttunen, senior lecturer
Kimmo Kuoppasalmi, acting associate professor

Correspondence to: Dr Isometsä.

BMJ 1995;310:1366-7

In most European countries men account for about three quarters of people who commit suicide. Suicide rates have generally risen among men under 35 but remained more stable among older men. To our knowledge, no studies specifically compare mental disorders among men of different ages who have committed suicide, although the association between mental disorders and suicide is strong and different secular trends in suicide rates have been reported. We compared the prevalences of mental disorders central to suicide—depressive syndromes, alcoholism, personality disorders, and non-affective psychotic disorders—in men of 20-34 and 35-59 who committed suicide in Finland.

Subjects, methods, and results

In the research phase of the national suicide prevention project in Finland all deaths between 1 April 1987 and 31 March 1988 that were officially classified as suicide (n = 1397) were recorded and carefully analysed retrospectively by the psychological autopsy method.¹ The definition of suicide was based on Finnish law—in every case of violent, sudden, or unexpected death the possibility of suicide is assessed by police and medico-legal examination. Data for psychological autopsies were collected from comprehensive interviews with relatives and health care staff; psychiatric, medical, and social agency records; police investigations; medicolegal examinations at necropsy and toxicological analyses; and suicide notes. Details of the methods have been published.¹

Mental disorders were retrospectively examined, according to the criteria of the *Diagnostic and Statistical Manual of Mental Disorders*, third edition, revised (DSM-III-R) in the diagnostic study of a random sample of 229 (16.4%) of the people who committed suicide.¹ The diagnostic evaluation was based on weighing and integrating all evidence and took place in two stages. Firstly, two pairs of psychiatrists independently gave provisional diagnoses, the reliability of which was tested with the κ coefficient and found to be good or at least moderate in the relevant

diagnostic categories ($\kappa = 0.52-0.94$).¹ Secondly, all cases of diagnostic disagreement were reanalysed by a third psychiatrist to achieve consensus for the final best estimate diagnoses. All men aged 20 to 59 from the random sample were included in the study, those aged 20-34 being compared with those aged 35-59. The table shows that the prevalence of psychotic disorders was significantly higher in the younger men and the prevalence of alcoholism significantly higher in the older.

Comment

Our study effectively accounted for all cases of suicide in Finland among men aged 20-59 over a year;

Comparison of retrospective best estimate diagnoses according to DSM-III-R in young and middle aged men from random sample of people who committed suicide in Finland

Diagnosis	Age 20-34 (n=44)	Age 35-59 (n=87)
<i>Axis I (clinical syndromes)</i>		
Depressive syndromes	26 (59)	56 (64)
Major depression	6 (14)	27 (31)
Depressive disorder not otherwise specified	14 (32)	18 (21)
Bipolar disorder, depressed or mixed	2 (5)	3 (3)
Dysthymia	0	3 (3)
Adjustment disorder with depressed mood	1 (2)	1 (1)
Organic mood disorder	0	2 (2)
95% Confidence interval for difference -23.0 to 12.4 ($\chi^2 = 0.35$; P = 0.55)		
Psychotic disorders*	11 (25)	7 (8)
Schizophrenia	6 (14)	4 (5)
Schizoaffective disorder,† depressive type	3 (7)	2 (2)
Other psychoses	2 (5)	1 (1)
95% Confidence interval for difference 2.9 to 31.0 ($\chi^2 = 7.09$; P = 0.008)		
Alcoholism	17 (39)	54 (62)
Alcohol dependence	7 (16)	49 (56)
Alcohol abuse	10 (23)	5 (6)
95% Confidence interval for difference 5.8 to 41.1 ($\chi^2 = 6.46$; P = 0.01)		
Other:		
Anxiety disorders	4 (9)	11 (13)
Alcohol intoxication	5 (11)	11 (13)
Anxiolytic abuse or dependence	2 (5)	5 (6)
Other organic mental disorders	0	4 (5)
Miscellaneous non-psychotic disorders	2 (5)	4 (5)
No diagnosis	1 (2)	0
Insufficient information	3 (7)	3 (3)
<i>Axis II (personality disorders)</i>		
Personality disorder	19 (43)	27 (31)
No diagnosis	15 (34)	44 (51)
Insufficient information	10 (23)	16 (18)
95% Confidence interval for difference -5.4 to 29.7 ($\chi^2 = 1.89$; P = 0.17)		
<i>Axis III (physical disorders and conditions)</i>		
Any physical disease	12 (27)	37 (43)
Comorbidity:		
No	12 (27)	12 (14)
Psychiatric	30 (68)	58 (67)
Psychiatric or somatopsychiatric	32 (73)	75 (86)

*Psychotic major depression (three cases in young men and 12 in middle aged men) not included.

†Included under depressive syndromes.

‡Cases having more than one diagnosis of mental disorder (axes I and II).

||Cases having mental disorder in conjunction with physical disease.