Recruiting ethnic minority medical students

EDITOR,—McManus and colleagues suggest that the potential for racial discrimination in recruitment to medical schools would be reduced if application forms were rendered anonymous.¹ Such approaches can easily become open to ridicule. By the time that the applicant's name, address, birth date, and place of education have been left off the application form—to address the risks of discrimination on grounds of race, sex, age, religion, and social class—selectors may find themselves reduced to interviewing candidates behind a screen, and perhaps even through an interpreter, so that their decisions cannot be affected by skin colour, appearance, voice, or accent either.

The causes of unfair discrimination do not lie with application forms. It is the selectors who are at the root of the problem. Experience in trying to ensure equal employment opportunities in both public and private sectors suggests that successful approaches to reducing unfair discrimination are likely to be based on clear, objective selection criteria free of bias, with training for selectors to eliminate the effects of any prejudice from their practice; this should be backed up by monitoring of outcomes at both shortlisting and selection stages—if necessary by individual selector—to see where there may be causes for concern. These approaches are at the heart of what the Commission for Racial Equality recommends for selection processes in employment and elsewhere.^{2,3} There has been much movement towards them across British medical education over the past 10 years or so, though more may still be needed.⁴

The selection criteria for medical education should be based on a rigorous analysis of what makes for the best possible doctors of the future. Such an analysis may or may not show that predicted A level results are the best available indicator of this, and it may or may not lead to indicators that are distributed equally across all sectors and strata of the population. But without it medical schools cannot be sure that variations in success rates among applicants from different ethnic backgrounds have a rational and justifiable explanation.

NIGEL TURNER
Director of human resources

Royal Free Hampstead NHS Trust,
London NW3 2QG

- 1 McManus IC, Richards P, Winder BC, Sproston KA, Styles V. Medical school applicants from ethnic minority groups: identifying if and when they are disadvantaged. *BMJ* 1995;310:496-500. (25 February.)
- 2 Commission for Racial Equality. *Code of practice for employers*. London: CRE, 1984.
- 3 Commission for Racial Equality. *Racial equality means business*. London: CRE, 1995.
- 4 Esmail A, Primarolo D, Nelson P, Toma T. Acceptance into medical school and racial discrimination. *BMJ* 1995;310:501-2. (25 February.)

Investigating racial discrimination in the NHS

EDITOR,—I was delighted to read that the BMA is to set up a small group looking into racial discrimination in the NHS and medical organisations.¹ As a past member of the Commission for Racial Equality for six years, I would offer a word of caution: the group should avoid the pitfalls of internal auditing and investigation and get the commission involved. This would provide the BMA with both credibility and, more importantly, the necessary skill. The Commission for Racial Equality has a large and relevant reservoir of information on discrimination in the NHS. It would also set initial guidelines for the BMA and suggest recommendations on remedial measures. An internal approach never succeeds, as has been

evident with complaints against the police. The BMA has not got the necessary skill.

DIPAK RAY
Chair, race equality committee
MSF Union, Birmingham B3 1JG

- 1 Beecham L. BMA will consider investigating racism in the NHS. *BMJ* 1995;310:64. (7 January.)

Doctors from ethnic minority groups

EDITOR,—The issues raised in the articles discussing whether doctors from ethnic minority groups should form separate associations¹ and A Akram Sayeed's pessimism about the success of such organisations in fighting against ethnic and racial discrimination² are relevant to discrimination on the grounds of caste in Indian medicine. In India doctors belonging to scheduled castes and scheduled tribes are discriminated against.³ Unlike in Britain and the United States, however, this is yet to be a matter of concern for Indian medical bodies or government agencies.⁴ Nevertheless, doctors in these categories have begun to assert themselves by forming their own associations at local or state level. Unfortunately, they have yet to develop a national organisation parallel to the Indian Medical Association. Interestingly, whereas there is provision for ethnic minorities to be represented on the General Medical Council, there is no such provision for the representation of scheduled castes and scheduled tribes on the Indian Medical Council.

I concur with A K Bose's views about the choice of names for such an organisation.¹ The organisation's name should reflect its objectives rather than its composition. Unfortunately, ethnocentric names such as Overseas Doctors Association (in Britain), Scheduled Caste and Scheduled Tribes Doctors Association (in India), etc have become a tradition. One might ask, "What is there in a name?"—but the name reflects the organisers' sociopolitical perspective with regard to changing the status quo. We have gone against tradition and have registered the name of our association as Forum for Rights and Equality. Furthermore, our membership is not restricted to people belonging to scheduled castes and scheduled tribes but is open to all who share our views.

L R MURMU
General secretary

Forum for Right and Equality,
All India Institute of Medical Sciences,
Ansari Nagar,
New Delhi-110029, India

- 1 Bose AK, Qureshi B, Kumar S, Goss B, McKenzie K, Kale R, et al. Should doctors from ethnic minority groups form separate associations? *BMJ* 1994;309:645-7. (10 September.)
- 2 Sayeed AA. Doctors from ethnic minority groups. *BMJ* 1994;309:1375. (19 November.)
- 3 Murmu LR. Caste discrimination in India. *BMJ* 1992;305:1099.
- 4 Murmu LR. Caste and medicine in India. *Natl Med J India* 1994;7:241-3.

National measles and rubella vaccination campaign

EDITOR,—The national schools measles and rubella campaign of November 1994,¹⁻³ in which most districts attained coverage of 90%, has been a success.⁴ This was despite adverse publicity and concern regarding possible adverse reactions to a new vaccine. All adverse reactions were reported to the Committee on Safety of Medicines through the yellow card system, with a copy to the immunisation coordinator. We analysed the information in St Helens and Knowsley district with EpiInfo version 5.00.

Altogether 48 000 children were immunised during November 1994, with an uptake rate of 90%. One hundred and sixteen reports (0.24%) of

Adverse reactions to Eolarix vaccine by sex, age, and batch

Batch	Boys (n=50)			Girls (n=66)		
	Aged 4-10	Aged 11-16	Age not stated	Aged 4-10	Aged 11-16	Age not stated
A	6	11	1	6	18	1
B	1	1	1	0	3	0
C	11	8	0	20	5	0
D	6	0	0	6	1	5
Not stated	4	—	—	—	1	—
Total	28	20	2	32	28	6

adverse reactions were received, 50 in boys and 66 in girls. Sixty were in children aged 10 and under (28 boys, 32 girls) and 48 in children over 10 (20 boys, 28 girls). In eight cases no age was stated. More reactions were reported in children aged 10 and under (56% of stated ages) than in the 11-16 year group (44%). There seemed to be no appreciable sex difference in the two age bands. Three quarters of the reports related to two batch numbers of the only vaccine used, Eolarix (SmithKline Beecham). For five reports the batch number was not stated.

The predominant symptoms were nausea (51 cases), dizziness (50), abdominal pain (33), headache (27), and fever (22). These were common to three of the four batch numbers used in this district and to the age bands. One report of anaphylaxis was received, in a 10 year old girl whose blood pressure was unrecordable and carotid pulse extremely weak and who had shallow breathing with severe peripheral cyanosis. She recovered completely after 0.5 ml intramuscular adrenaline and admission to hospital. Six children received outpatient treatment, while the rest recovered with reassurance.

A mass campaign of this nature was new to most staff, and there may have been some apprehension. Therefore the possibility of reactions being noted overenthusiastically must not be forgotten.

Two facts emerged concerning the manufacturer's datasheet and side effects. Firstly, the datasheet states that side effects are delayed for several days. The reports received, however, show that side effects developed in a matter of hours. Secondly, the predominant symptoms of nausea, dizziness, and abdominal pain noted do not feature in the datasheet. We are, however, aware that cases reported through the yellow card system, which has been termed a passive surveillance system and considered to be inadequate,⁵ are biased towards those occurring soon after immunisation, and the population studied (48 000) is too small for any rare adverse events to be detected.

Studies elsewhere have noted an expected 0.25% rate of mild adverse reactions after a second dose of measles, mumps, and rubella vaccine.² Our finding is in line with these. As only 0.24% of those immunised suffered from adverse reactions of a mild nature it seems that this new vaccine is free of serious side effects, but further information nationally is required.

E B P WIRATUNGA
Consultant in communicable disease control
JEANETTE O'BRIEN
Database manager

St Helens and Knowsley Health,
St Helens,
Merseyside WA10 2AP

- 1 Department of Health. *National measles and rubella immunisation campaign*. London: DoH, 1994. (PL/CMO(94)10.)
- 2 Department of Health. *Measles and rubella immunisation campaign*. London: DoH, 1994. (PL/CMO(94)12.)
- 3 Ramsay M, Gay N, Miller E, Rush M, White J, Morgan-Capner P, et al. The epidemiology of measles in England and Wales: rationale for the 1994 national vaccination campaign. *Commun Dis Rep CDR Rev* 1994;4:R141-6.
- 4 Department of Health. *Update on national measles/rubella immunisation campaign*. London: DoH, 1994. (PL/CMO(94)13.)
- 5 Farrington P, Pugh S, Colville A, Flower A, Nash J, Morgan-Capner P, et al. A new method for active surveillance of adverse events from diphtheria/tetanus/pertussis and measles/mumps/rubella vaccines. *Lancet* 1995;345:567-9.