Practical problems can be foreseen: a patient who has taken an overdose may have signed an advance directive stating "do not resuscitate," or a directive stating "I am not to be subjected to any medical treatment aimed at prolonging or sustaining life" may be found when a patient is on the operating table. Nor should it be accepted that analgesics can be used to treat thirst when simple help with nutrition will suffice.

When advance directives have been made legally enforceable, problems such as these have arisen. The doctor has been unable to treat a conscious patient who has lost the capacity to revoke the directive. As a result the patient has suffered, sometimes greatly.

The BMA's original stance on advance directives was that they should not be legally binding but should be a starting point for an improved dialogue with patients over choice of treatment. After certain judgments there has been an overhasty change. There is still, however, no mandate for the BMA's council to support the Law Commission's recommendations without widely consulting members.

There is no evidence in Britain that patients who become incompetent are being treated against their wishes or against their best interests. Nor is there any study of what the general public, as opposed to pressure groups that are seeking to advance their own agenda, really want. The original concept of a living will was introduced by Kutner in a commentary entitled "Due process of euthanasia: the living will a proposal."²

A P COLE
Consultant paediatrician

Worcester Royal Infirmary NHS Trust,
Ronswood Branch,
Worcester WR5 1HN

- 1 Doyal L. Advance directives. *BMJ* 1995;310:612-3. (11 March.)
- 2 Kutner L. Due process of euthanasia: the living will a proposal. *Indiana Law Journal* 1969;XLIX:539-54.

Incompetence develops gradually

EDITOR,—Incompetence is not an all or nothing issue (as the Law Commission recognises).¹ Dementia may cause a gradual erosion of competence, and it might be difficult to decide at what point a patient lost the ability to revoke an earlier advance directive. This can be covered in a properly framed law, but we must be vigilant to make sure that the law is properly framed and that doctors are not made to face unnecessary and impossible moral and ethical dilemmas by an oversimplistic approach.

JOHN P WATTIS
Consultant in psychiatry for older adults

Leeds Community and Mental Health Services,
Leeds LS10 2TP

- 1 Doyal L. Advance directives. *BMJ* 1995;310:612-3. (11 March.)

Keyhole surgery

EDITOR,—M J Notaras may make a case for the development of specialised (perhaps private) hernia centres but fails to make any case against laparoscopic hernia repair.¹ It is true that excellent results are reported from internationally recognised hernia repair centres (although the completeness and methods of follow up do not always stand close examination). Unfortunately, however, everyday clinical reality is different. In Britain most hernias are not repaired by experts and those of us who audit hernia repairs are aware that results are far from satisfactory.

An epidemiological study of open hernia repair in a health district in Britain reported poor results, although absolute numbers of patients were not published: recurrence had occurred by three years in 31% of cases, wound infection in 14%, persistent deep sinuses in 6%, persistent pain in 16%, and

testicular atrophy in 7%, and altogether 47% of patients were dissatisfied.² Probably this represents the worst end of the spectrum, but such results may not be as atypical as we would like to think. It is incorrect to state that organ injury and death do not occur with open elective hernia repairs. They do but are rarely reported.

Laparoscopic hernia repair can be safely carried out as a day case procedure with an entirely extraperitoneal approach. True, at present general anaesthesia is required, but in our experience most patients prefer this, and local anaesthesia is not without complications. A recently published randomised trial has indicated an overall advantage for laparoscopic repair in terms of more rapid discharge and convalescence,³ and our experience of over 700 cases supports this strongly; the advantages are even more obvious for bilateral and recurrent hernia repair.

Are such trials justified? Notaras would have us believe not, but they must take place. Perhaps the opponents of trials are frightened of the results. The mere preference of a surgeon for a particular technique does not necessarily mean that it is the best. We agree that in skilled hands the Lichtenstein repair is an excellent operation, but let it be put to the test in a randomised trial against the best laparoscopic techniques.

R STAYLOR
Consultant surgeon
T M LOOSEMORE
Consultant surgeon

St George's Hospital,
London SW17 0QT

- 1 Notaras MJ. Keyhole surgery. *BMJ* 1995;310:871-2. (1 April.)
- 2 Devlin B. In: Katkhouda N. Treatment of bilateral hernia by open surgery—the STOPPA positional repair. In: Schumpelick V, Wantz GE, eds. *Inguinal hernia repair*. Basel: Karger, 1995.
- 3 Wilson MS, Deans GT, Brough WA. Prospective trial comparing Lichtenstein with laparoscopic tension-free mesh repair of inguinal hernia. *Br J Surg* 1995;82:274-7.

Accident and emergency radiology

"Rule of two" is outmoded and unhelpful

EDITOR,—We disagree with several parts of the "rule of twos" in Robin Touquet and colleagues' article on teaching radiology in accident and emergency medicine.¹ The authors state that when a fracture is suspected the radiograph must include the joint at either end of the long bones because of the risk of an associated dislocation or subluxation. If this practice were carried out then all radiographs of twisted ankles would include the knee and all radiographs of sprained wrists would include the elbow. This is clearly not practical or necessary. It is true that in certain instances, such as a fracture of the medial malleolus, there may be an associated fracture of the upper end of the fibula. A senior house officer should know this and carefully examine the knee, obtaining radiographs only in those cases in which there is tenderness. There is also a danger that the radiographer may be tempted to include both joints and the long bone in a single film. This results in an inadequate examination of both joints because of the divergence of the x ray beam and difficulty in getting both joints square on to the film.

It is suggested that in a child with limited extension of the elbow after injury a radiograph of the normal side should be obtained for comparison at the outset. This is an outmoded practice, and few mothers nowadays will allow an x ray examination of an unaffected limb just because the doctor does not know what normality looks like. In our department we have a book showing normal elbows at various stages of development. I cannot remember the last time we obtained a radiograph of a joint for comparison.

The authors recommend that radiographs

should be obtained before and after the removal of a foreign body in a wound. It is more sensible to remove any obvious foreign bodies, clean the wound, and then obtain a radiograph to see if any foreign bodies remain in the wound. Obtaining a radiograph of something that can be seen with the naked eye is pointless.

The rule of twos is unhelpful. It results in too many radiographs, too much radiation, and too much cost.

A H TROUGHTON
Consultant radiologist

I KENDALL
Consultant in accident and emergency medicine

Princess Margaret Hospital,
Swindon SN1 4JU

- 1 Touquet R, Driscoll P, Nicolson D. Teaching in accident and emergency medicine: 10 commandments of accident and emergency radiology. *BMJ* 1995;310:642-5. (11 March.)

Comparison view radiographs are not routinely required

EDITOR,—Robin Touquet and colleagues state that "a child with limited extension of the elbow after injury should have radiographs taken of both elbow joints for comparison at the outset."¹ This suggests that comparison views should be taken as a routine in all cases, and we take issue with this.

Firstly, this guidance is directly contrary to recommendations produced by the Royal College of Radiologists.² Secondly, contralateral radiographs are superfluous if the fracture is visible in radiographs of the affected side. Thirdly, comparable radiographs of the contralateral side are difficult to obtain. Radiographs of the affected side are often poorly positioned because of pain and swelling. If contralateral views are to be used one should ensure that the exact position of the affected side is mimicked. Fourthly, contralateral views are unnecessary if good radiological opinion is readily available. Finally, the diagnosis of subtle fractures around the elbow joint will not necessarily change the clinical management. While we agree that radiographs of the elbow in children can be difficult to interpret, we believe that the correct approach is to combine the results of clinical examination with expert radiological advice. This may occasionally result in further radiological examination being advised.

T F BEATTIE
Consultant in accident and emergency

M HENDRY
Consultant paediatric radiologist

M McPHILLIPS
Consultant paediatric radiologist

S MACKENZIE
Consultant paediatric radiologist

Royal Hospital for Sick Children,
Edinburgh EH9 1LF

- 1 Touquet R, Driscoll P, Nicolson D. Teaching in accident and emergency medicine: 10 commandments of accident and emergency radiology. *BMJ* 1995;310:642-5. (11 March.)
- 2 Royal College of Radiologists. *Making the best use of the department of clinical radiology: guidelines for doctors*. 2nd ed. London: Royal College of Radiologists, 1993.

Elderly people and the 1990 act

EDITOR,—Victoria Ajayi and colleagues state that "the reduction in the length of hospital stay is consistent with the findings that after the legislation patients were more likely to be in hospital for rehabilitation than awaiting placement."¹ Our experience in Edinburgh seems to confirm this finding.

Since 1988 we have carried out five surveys to monitor the prevalence of elderly patients awaiting placements for long stay care in our general hospitals. The first two surveys were reported in 1992² and the results of the last three surveys have