indicates, however, that these reports seriously underestimate the prevalence of side effects in travellers: only rarely does a week pass in which I am not informed (at the Hospital for Tropical Diseases) by at least one traveller of his or her personal experience of side effects of mefloquine (many of them severe) or of similar symptoms in a colleague or fellow traveller. Many travellers refuse to take mefloquine in the light of their experience of its neuropsychiatric side effects.

The regimen of chloroquine plus proguanil has a low incidence of side effects and for much of sub-Saharan Africa is probably only marginally inferior in efficacy to mefloquine. Furthermore, widespread use of mefloquine for chemoprophylaxis raises the likelihood of the emergence of *P. falciparum* resistant to quinine (considerable cross resistance exists between mefloquine and quinine); this is a genuine but understated problem.

I recommend chloroquine plus proguanil for all countries of Africa in which infection with *P. falciparum* is a potential hazard. In addition, travellers should be advised to take standby treatment (usually quinine) for use in the event of a febrile illness that cannot be adequately assessed or treated at a reputable medical centre. Mefloquine should be reserved for chemotherapy of infection with *P. falciparum* that is resistant to quinine.

G C COOK
Consultant physician

Hospital for Tropical Diseases,
London NW1 0PE

- Bradley DJ, Warhurst DC. Malaria prophylaxis: guidelines for travellers from Britain. *BMJ* 1995;310:709-14. (18 March.)
- Lobel HO, Miani M, Eng T, Bernard KW, Hightower AW, Campbell CC. Long term malaria prophylaxis with weekly mefloquine. *Lancet* 1993;341:848-51.
- Steffen R, Fuchs E, Schildknecht J. Mefloquine compared with other malaria chemoprophylactic regimens in tourists visiting East Africa. *Lancet* 1993;341:1299-303.
- Westeyn JCFM, de Geus A. Comparison of three regimens for malaria prophylaxis in travellers to east, central, and southern Africa. *BMJ* 1993;307:1041-3.
- Weinke T, Trautmann M, Held T, Weber G, Eichenlaub D, Fleischer K, et al. Neuropsychiatric side effects after the use of mefloquine. *Am J Trop Med Hyg* 1991;45:86-91.

Generous prescribing advised for travelling to areas of drug shortage

EDITOR,—The guidelines on malaria prophylaxis for travellers from Britain advise that people who are unable to reach medical advice within 24 hours of becoming ill should be provided with standby treatment for malaria.¹ This point is emphasised several times.

I recently returned from working in a hospital with 210 beds in northern Malawi. In this area falciparum malaria is resistant to chloroquine in a high proportion of cases. Malawi has been using pyrimethamine-sulfadoxine as first line treatment for malaria since 1992. Resistance to pyrimethamine-sulfadoxine is increasing, and quinine is the recommended second line treatment. Supplies of quinine have been scarce in northern Malawi, and the hospital ran out of stock last February.

Medical advice may be within 24 hours' reach, but this is no guarantee that the necessary drugs will be available. Although this point is mentioned in the review, it is not highlighted. It is difficult to predict this situation from Britain, and I suggest that doctors should not be too restrictive in providing standby treatments for malaria to travellers.

GRAHAM MORRISON
General practitioner

Kersland House Surgery,
Millingavie,
Glasgow G62 8BT

- Bradley DJ, Warhurst DC, on behalf of a meeting convened by the Malaria Reference Laboratory. Malaria prophylaxis: guidelines for travellers from Britain. *BMJ* 1995;310:709-14. (18 March.)

Remember malaria, even after a year

EDITOR,—The guidelines for prophylaxis against malaria for travellers from Britain mention reports of vivax malaria resistant to chloroquine in Oceania and recommend vigilance for a year after the travellers have returned from malarious regions.¹ I worked as a locum paediatrician in Papua New Guinea for three months, leaving there in January 1990 and arriving in England (via Australia, Indonesia, Thailand, Singapore, and Egypt) in May 1990.

I developed a severe flu-like illness in August 1991, 18 months after leaving Papua New Guinea and 15 months after returning to England. I was on call at a hospital in London at the time and started to have a rigor while in the mess, although I felt well. That night I felt hot, although I was initially afebrile, and had severe nausea and vomiting. I arranged for a full blood count before going home the next day.

I remained bedbound, with my temperature regularly over 40°C and total anorexia, and was too weak to walk to the lavatory. Two days later I phoned for the results of the blood count, which showed a total white cell count of $3.8 \times 10^9/l$ and a platelet count of $69 \times 10^9/l$. The blood had been discarded before the film was looked at. I assumed that I had leukaemia.

I was driven to my local general practitioner and asked for a repeat full blood count and (rather self consciously, not having been abroad for over a year) a malarial film. The remainder is obvious. I had vivax malaria despite having taken 300 mg chloroquine weekly (for six as opposed to the recommended four weeks on my return) and not having been abroad for 15 months. By the time the diagnosis was made I had to be carried downstairs on a stretcher and required transfer to hospital by ambulance and intravenous rehydration in intensive care.

The message is clear: malaria can occasionally occur for the first time after the usually quoted 12 months. In cases in which the diagnosis is uncertain and malaria is possible it is indefensible not to look for malarial parasites.

CHARLES GODDEN
Paediatric senior registrar

Poole NHS Trust Hospital,
Poole,
Dorset BH15 2JB

- Bradley DJ, Warhurst DC. Malaria prophylaxis: guidelines for travellers from Britain. *BMJ* 1995;310:709-14. (18 March.)

Toxicity of mefloquine is similar to that of other chemoprophylaxis

EDITOR,—The revised guidelines on chemoprophylaxis against malaria perpetuate the notion that mefloquine is relatively innocuous when used for this purpose.¹ This view has been challenged by some authorities,² who argue that the side effects of mefloquine have been greatly underreported. As White recognised in an editorial on mefloquine,

Incidence of adverse events to malaria chemoprophylaxis of three to seven days' duration occurring while soldiers were in Kenya

Adverse event	Percentage (No) of respondents reporting event		Observed difference (95% confidence interval) between proportions
	Mefloquine arm (n=145)	Chloroquine-proguanil arm (n=142)	
Abdominal pain	0.15 (21)	0.19 (27)	-0.05 (-0.13 to 0.04)
Nausea	0.12 (18)	0.16 (23)	-0.04 (-0.12 to 0.04)
Anorexia	0.09 (13)	0.12 (8)	-0.04 (-0.11 to 0.03)
Fever	0.01 (1)	0.03 (5)	-0.03 (-0.06 to 0.00)
Mouth ulcers	0.08 (11)	0.07 (10)	0.01 (-0.06 to 0.06)
Pruritus	0.07 (10)	0.02 (3)	0.05 (0.0 to 0.10)
Sleeplessness	0.06 (8)	0.08 (12)	-0.03 (-0.08 to 0.03)
Abnormal dreams	0.08 (12)	0.08 (11)	0.01 (-0.06 to 0.07)
Depression	0.02 (3)	0.03 (4)	-0.01 (-0.04 to 0.03)
Anxiety attacks	0.01 (2)	0.01 (1)	0.01 (-0.02 to 0.03)
Paranoid feelings	0.03 (5)	0.0 (0)	0.03 (0.0 to 0.06)
Tremor	0.01 (1)	0.02 (3)	-0.01 (-0.04 to 0.01)

this tendency to understate the toxicity of the drug has followed from the poor scientific quality of almost all of the early studies on use of mefloquine.³

Since January the Ministry of Defence has been conducting a double blind randomised controlled trial of chemoprophylaxis with mefloquine versus chloroquine-proguanil. The subjects of the trial are British troops exercising in Kenya. Of the total trial population of 624 soldiers, 317 were randomly assigned, by means of random numbers generated by a computer, to receive mefloquine and 307 to receive chloroquine-proguanil. A questionnaire on "unusual" symptoms or illnesses was administered at eight weeks of chemoprophylaxis and was returned by 145 (46%) soldiers in the mefloquine arm of the trial and 142 (46%) in the chloroquine-proguanil arm.

The preliminary results of the trial show that both mefloquine and chloroquine-proguanil have a much higher mild toxicity than has commonly been recognised. Altogether 131 (90%) respondents given mefloquine reported some toxicity as a result of their (unknown) chemoprophylaxis, as did 126 (89%) responders given chloroquine-proguanil. The profile of adverse reactions for the two regimens has been surprisingly similar. For example, the table shows the reported incidence of adverse events of three to seven days' duration occurring at least once while the soldiers were in Kenya.

Hansford has pointed out that, whereas dosages of prophylactic antimalarial drugs for children are calculated against body weight, for adults a single standard dose is invariably prescribed, irrespective of body weight.⁴ Could it not be that much of the mild toxicity of prophylactic antimalarial drugs is due to this anomaly, coupled with the fact that some antimalarial drugs (mefloquine, for example⁵) are known to be metabolised idiosyncratically?

A more flexible regimen of mefloquine and other prophylactic antimalarial drugs is needed for adults, such that the drugs' therapeutic ratio is improved—that is, the protective action of the drug is maintained while individual toxicity is minimised. Until such regimens are developed a proportion of travellers to malarious regions will continue to feel ill as a result of their prescribed treatment and some of these will stop taking their tablets. Many cases of malaria will thereby result, which could otherwise have been avoided.

ASHLEY CROFT
Senior registrar in public health medicine

Army Medical Directorate,
Ministry of Defence,
Aldershot GU12 5RR

- Bradley DJ, Warhurst DC, on behalf of a meeting convened by the Malaria Reference Laboratory. Malaria prophylaxis: guidelines for travellers from Britain. *BMJ* 1995;310:709-14. (18 March.)
- Luzzi GA, Peto TEA. Adverse effects of antimalarials—an update. *Drug Safety* 1993;8:295-311.
- White NJ. Mefloquine. *BMJ* 1994;308:286-7.
- Hansford CH. Malaria prophylaxis dosage. *Lancet* 1995;345:1049.
- Desjardins RE, Pamplin CL, Bredow J, Barry KG, Canfield CJ. Kinetics of a new antimalarial, mefloquine. *Clin Pharmacol Ther* 1979;26:372-9.